

---

**Question 1:**

Why is reproduction essential for organisms?

Answer

Reproduction is a fundamental feature of all living organisms. It is a biological process through which living organisms produce offspring's similar to them. Reproduction ensures the continuance of various species on the Earth. In the absence of reproduction, the species will not be able to exist for a long time and may soon get extinct.

**Question 2:**

Which is a better mode of reproduction sexual or asexual? Why?

Answer

Sexual reproduction is a better mode of reproduction. It allows the formation of new variants by the combination of the DNA from two different individuals, typically one of each sex. It involves the fusion of the male and the female gamete to produce variants, which are not identical to their parents and to themselves. This variation allows the individual to adapt to constantly changing and challenging environments. Also, it leads to the evolution of better suited organisms which ensures greater survival of a species. On the contrary, asexual reproduction allows very little or no variation at all. As a result, the individuals produced are exact copies of their parents and themselves.

**Question 3:**

Why is the offspring formed by asexual reproduction referred to as clone?

Answer

A clone is a group of morphologically and genetically identical individuals.

In the process of asexual reproduction, only one parent is involved and there is no fusion of the male and the female gamete. As a result, the offsprings so produced are morphologically and genetically similar to their parents and are thus, called clones.

---

**Question 4:**

Offspring formed due to sexual reproduction have better chances of survival. Why? Is this statement always true?

Answer

Sexual reproduction involves the fusion of the male and the female gamete. This fusion allows the formation of new variants by the combination of the DNA from two (usually) different members of the species. The variations allow the individuals to adapt under varied environmental conditions for better chances of survival.


However, it is not always necessary that the offspring produced due to sexual reproduction has better chances of survival. Under some circumstances, asexual reproduction is more advantageous for certain organisms. For example, some individuals who do not move from one place to another and are well settled in their environment. Also, asexual reproduction is a fast and a quick mode of reproduction which does not consume much time and energy as compared to sexual reproduction.

**Question 5:**

How does the progeny formed from asexual reproduction differ from those formed by sexual reproduction?

Answer

Progeny formed from asexual reproduction		Progeny formed from sexual reproduction
1.	Asexual reproduction does not involve the fusion of the male and the female gamete. Organisms undergoing this kind of reproduction produce offspring's that are morphologically and genetically identical to them.	Sexual reproduction involves the fusion of the male and the female gamete of two individuals, typically one of each sex. Organisms undergoing this kind of reproduction produce offspring's that are not identical to them.



**Question 11:**

Explain (in one or two lines) the function of the followings:

**(a)** Promoter

**(b)** tRNA

**(c)** Exons

Answer

**(a)** Promoter

Promoter is a region of DNA that helps in initiating the process of transcription. It serves as the binding site for RNA polymerase.

**(b)** tRNA

tRNA or transfer RNA is a small RNA that reads the genetic code present on mRNA. It carries specific amino acid to mRNA on ribosome during translation of proteins.

**(c)** Exons

Exons are coding sequences of DNA in eukaryotes that transcribe for proteins.

**Question 12:**

Why is the Human Genome project called a mega project?

Answer

---

the fusion of the male and the female gamete and requires only one parent. Hence, vegetative reproduction is considered as a type of asexual reproduction.

**Question 7:**

What is vegetative propagation? Give two suitable examples.


Answer

Vegetative propagation is a mode of asexual reproduction in which new plants are obtained from the vegetative parts of plants. It does not involve the production of seeds or spores for the propagation of new plants. Vegetative parts of plants such as runners, rhizomes, suckers, tubers, etc. can be used as propagules for raising new plants.


Examples of vegetative reproduction are:

**1. Eyes of potato:**

The surface of a potato has several buds called eyes. Each of these buds when buried in soil develops into a new plant, which is identical to the parent plant.

**2. Leaf buds of Bryophyllum:**

The leaves of *Bryophyllum* plants bear several adventitious buds on their margins. These leaf buds have the ability to grow and develop into tiny plants when the leaves get detached from the plant and come in contact with moist soil.



**Question 8:**

Define

- (a)** Juvenile phase,
- (b)** Reproductive phase,
- (c)** Senescent phase.

Answer

- (a)** Juvenile phase:

It is the period of growth in an individual organism after its birth and before it reaches reproductive maturity.

- (b)** Reproductive phase:

It is the period when an individual organism reproduces sexually.

- (c)** Senescent phase:

It is the period when an organism grows old and loses the ability to reproduce.

**Question 9:**

Higher organisms have resorted to sexual reproduction in spite of its complexity. Why?

Answer

Although sexual reproduction involves more time and energy, higher organisms have resorted to sexual reproduction in spite of its complexity. This is because this mode of reproduction helps introduce new variations in progenies through the combination of the DNA from two (usually) different individuals. These variations allow the

---

individual to cope with various environmental conditions and thus, make the organism better suited for the environment. Variations also lead to the evolution of better organisms and therefore, provide better chances of survival. On the other hand, asexual reproduction does not provide genetic differences in the individuals produced.

**Question 10:**

Explain why meiosis and gametogenesis are always interlinked?

Answer

Meiosis is a process of reductional division in which the amount of genetic material is reduced. Gametogenesis is the process of the formation of gametes. Gametes produced by organisms are haploids (containing only one set of chromosomes), while the body of an organism is diploid. Therefore, for producing haploid gametes (gametogenesis), the germ cells of an organism undergo meiosis. During the process, the meiocytes of an organism undergo two successive nuclear and cell divisions with a single cycle of DNA replication to form the haploid gametes.

**Question 11:**

Identify each part in a flowering plant and write whether it is haploid (n) or diploid (2n).

(a) Ovary —————

(b) Anther —————

(c) Egg —————

(d) Pollen —————

(e) Male gamete —————

(f) Zygote —————

Answer

(a) Ovary Diploid (2n)

(b) Anther Diploid (2n)

(c) Egg Haploid (n)

(d) Pollen Haploid ( $n$ )

(e) Male gamete Haploid ( $n$ )

(f) Zygote Diploid ( $2n$ )

**Question 12:**

Define external fertilization. Mention its disadvantages.

Answer

External fertilization is the process in which the fusion of the male and the female gamete takes place outside the female body in an external medium, generally water. Fish, frog, starfish are some organisms that exhibit external fertilization.

**Disadvantages of external fertilization:**

In external fertilization, eggs have less chances of fertilization. This can lead to the wastage of a large number of eggs produced during the process.

Further, there is an absence of proper parental care to the offspring, which results in a low rate of survival in the progenies.

**Question 13:**

Differentiate between a zoospore and a zygote.

Answer

Zoospore		Zygote
1.	A zoospore is a motile asexual spore that utilizes the flagella for movement.	A zygote is a non-motile diploid cell formed as a result of fertilization.
2.	It is an asexual reproductive structure.	It is formed as a result of sexual reproduction.

**Question 14:**

Differentiate between gametogenesis from embryogenesis.

Answer

---

Gametogenesis	Embryogenesis
It is the process of the formation of haploid male and female gametes from diploid meiocytes through the process of meiosis.	It is the process of the development of the embryo from the repeated mitotic divisions of the diploid zygote.

**Question 15:**

Describe the post-fertilization changes in a flower.

Answer

Fertilization is the process of the fusion of the male and the female gamete to form a diploid zygote. After fertilization, the zygote divides several times to form an embryo. The fertilized ovule forms a seed. The seed contains an embryo, enclosed in a protective covering, called the seed coat. As the seed grows further, other floral parts wither and fall off. This leads to the growth of the ovary, which enlarges and ripens to become a fruit with a thick wall called the pericarp.

**Question 16:**

What is a bisexual flower? Collect five bisexual flowers from your neighborhood and with the help of your teacher find out their common and scientific names.

Answer

A flower that contains both the male and female reproductive structure (stamen and pistil) is called a bisexual flower. Examples of plants bearing bisexual flowers are:

- (1)** Water lily ( *Nymphaea odorata* )
  - (2)** Rose ( *Rosa multiflora* )
  - (3)** Hibiscus ( *Hibiscus Rosa-sinensis* )
  - (4)** Mustard ( *Brassica nigra* )
  - (5)** Petunia ( *Petunia hybrida* )
-



---

**Question 17:**

Examine a few flowers of any cucurbit plant and try to identify the staminate and pistillate flowers. Do you know any other plant that bears unisexual flowers?

Answer



Cucurbit plant bears unisexual flowers as these flowers have either the stamen or the pistil. The staminate flowers bear bright, yellow coloured petals along with stamens that represent the male reproductive structure. On the other hand, the pistillate flowers bear only the pistil that represents the female reproductive structure.

Other examples of plants that bear unisexual flowers are corn, papaya, cucumber, etc.

**Question 18:**

Why are offspring of oviparous animals at a greater risk as compared to offspring of viviparous animals?

Answer

Oviparous animals lay eggs outside their body. As a result, the eggs of these animals are under continuous threat from various environmental factors. On the other hand, in viviparous animals, the development of the egg takes place inside the body of the female. Hence, the offspring of an egg-laying or oviparous animal is at greater risk as compared to the offspring of a viviparous animal, which gives birth to its young ones.

---

---

**Question 1:**

Name the parts of an angiosperm flower in which development of male and female gametophyte take place.

Answer

The male gametophyte or the pollen grain develops inside the pollen chamber of the anther, whereas the female gametophyte (also known as the embryo sac) develops inside the nucellus of the ovule from the functional megaspore.

**Question 2:**

Differentiate between microsporogenesis and megasporogenesis. Which type of cell division occurs during these events? Name the structures formed at the end of these two events.

Answer

(a)

Microsporogenesis		Megasporogenesis
1.	It is the process of the formation of microspore tetrads from a microspore mother cell through meiosis.	It is the process of the formation of the four megaspores from a megaspore mother cell in the region of the nucellus through meiosis
2.	It occurs inside the pollen sac of the anther.	It occurs inside the ovule.

(b) Both events (microsporogenesis and megasporogenesis) involve the process of meiosis or reduction division which results in the formation of haploid gametes from the microspore and megaspore mother cells.

(c) Microsporogenesis results in the formation of haploid microspores from a diploid microspore mother cell. On the other hand, megasporogenesis results in the formation of haploid megaspores from a diploid megaspore mother cell.

---