

Exercise 10.1

1. Represent graphically a displacement of 40 km, 30° east of north.


Sol. Displacement 40 km, 30° East of North.

\Rightarrow Displacement vector \vec{OA} (say)

such that $|\vec{OA}| = 40$ (given)

and vector \vec{OA} makes an angle 30° with North in East-North quadrant.

Note. α° South of West \Rightarrow A vector in South-West quadrant making an angle of α° with West.



2. Check the following measures as scalars and vectors:

- (i) 10 kg (ii) 2 meters north-west (iii) 40°
(iv) 40 Watt (v) 10^{-19} coulomb (vi) 20 m/sec^2 .

Sol. (i) 10 kg is a measure of mass and therefore a scalar.

(\because 10 kg has no direction, it is magnitude only).

(ii) 2 meters North-West is a measure of velocity (i.e., has magnitude and direction both) and hence is a vector.

(iii) 40° is a measure of angle i.e., is magnitude only and, therefore, a scalar.

(iv) 40 Watt is a measure of power (i.e., 40 watt has no direction) and, therefore, a scalar.

(v) 10^{-19} coulomb is a measure of electric charge (i.e., is magnitude only) and, therefore, a scalar.

(vi) 20 m/sec^2 is a measure of acceleration i.e., is a measure of rate of change of velocity and hence is a vector.


3. Classify the following as scalar and vector quantities:

- (i) time period (ii) distance (iii) force
(iv) velocity (v) work done.

Sol. (i) Time-scalar (ii) Distance-scalar (iii) Force-vector
(iv) Velocity-vector (v) Work done-scalar.

4. In the adjoining figure, (a square), identify the following vectors.

- (i) Coinitial
(ii) Equal
(iii) Collinear but not equal.



Sol. (i) \vec{a} and \vec{d} have same initial point and, therefore, coinitial vectors.

(ii) \vec{b} and \vec{d} have same direction and same magnitude. Therefore,
 \vec{b} and \vec{d} are equal vectors.

(iii) \vec{a} and \vec{c} have parallel supports, so that they are collinear. Since they have opposite directions, they are not equal.

Hence \vec{a} and \vec{c} are collinear but not equal.

5. Answer the following as true or false.

- (i) \vec{a} and $-\vec{a}$ are collinear.
- (ii) Two collinear vectors are always equal in magnitude.
- (iii) Two vectors having same magnitude are collinear.
- (iv) Two collinear vectors having the same magnitude are equal.

Sol. (i) True.

(ii) False. ($\because \vec{a}$ and $2\vec{a}$ are collinear vectors but $|2\vec{a}| = 2|\vec{a}|$)

(iii) False.

($\because |\hat{i}| = |\hat{j}| = 1$ but \hat{i} and \hat{j} are vectors along x-axis (OX) and y-axis (OY) respectively.)

(iv) False.

(\because Vectors \vec{a} and $-\vec{a}$ ($= (-1)\vec{a} = m\vec{a}$) are collinear vectors and $|\vec{a}| = |-\vec{a}|$ but we know that $\vec{a} \neq -\vec{a}$ because their directions are opposite).

Note. Two vectors \vec{a} and \vec{b} are said to be equal if

- (i) $|\vec{a}| = |\vec{b}|$ (ii) \vec{a} and \vec{b} have same (like) direction.

Exercise 10.2

1. Compute the magnitude of the following vectors:

$$\vec{a} = \hat{i} + \hat{j} + \hat{k}, \quad \vec{b} = 2\hat{i} - 7\hat{j} - 3\hat{k},$$

$$\vec{c} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} - \frac{1}{\sqrt{3}}\hat{k}.$$

Sol. Given: $\vec{a} = \hat{i} + \hat{j} + \hat{k}.$

$$\text{Therefore, } |\vec{a}| = \sqrt{x^2 + y^2 + z^2} = \sqrt{1+1+1} = \sqrt{3}.$$

$$\vec{b} = 2\hat{i} - 7\hat{j} - 3\hat{k}.$$

$$\text{Therefore, } |\vec{b}| = \sqrt{4+49+9} = \sqrt{62}.$$

$$\vec{c} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} - \frac{1}{\sqrt{3}}\hat{k}.$$

$$\begin{aligned}\text{Therefore, } |\vec{c}| &= \sqrt{\left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{-1}{\sqrt{3}}\right)^2} \\ &= \sqrt{\frac{1}{3} + \frac{1}{3} + \frac{1}{3}} = \sqrt{\frac{3}{3}} = \sqrt{1} = 1.\end{aligned}$$

2. Write two different vectors having same magnitude.

Sol. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + \hat{j} - \hat{k}$.

Clearly, $\vec{a} \neq \vec{b}$. (\because Coefficients of \hat{i} and \hat{j} are same in vectors \vec{a} and \vec{b} but coefficients of \hat{k} in \vec{a} and \vec{b} are unequal as $1 \neq -1$).

$$\text{But } |\vec{a}| = \sqrt{x^2 + y^2 + z^2} = \sqrt{1+1+1} = \sqrt{3}$$

$$\text{and } |\vec{b}| = \sqrt{1+1+1} = \sqrt{3} \quad \therefore |\vec{a}| = |\vec{b}|.$$

Remark. In this way, we can construct an infinite number of possible answers.

3. Write two different vectors having same direction.

Sol. Let $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$... (i)

$$\begin{aligned}\text{and } \vec{b} &= 2(\hat{i} + 2\hat{j} + 3\hat{k}) \quad \dots (ii) \\ &= 2\vec{a} \quad \quad \quad [\text{By (i)}]\end{aligned}$$

$$\therefore \vec{b} = m\vec{a} \text{ where } m = 2 > 0.$$

\therefore Vectors \vec{a} and \vec{b} have the same direction.

$$\text{But } \vec{b} \neq \vec{a} [\because \vec{b} = 2\vec{a} \Rightarrow |\vec{b}| = |2||\vec{a}| = 2|\vec{a}| \neq |\vec{a}|]$$

Remark. In this way, we can construct an infinite number of possible answers.

4. Find the values of x and y so that the vectors $2\hat{i} + 3\hat{j}$ and $x\hat{i} + y\hat{j}$ are equal.

Sol. Given: $2\hat{i} + 3\hat{j} = x\hat{i} + y\hat{j}$.

Comparing coefficients of \hat{i} and \hat{j} on both sides, we have
 $x = 2$ and $y = 3$.

5. Find the scalar and vector components of the vector with initial point (2, 1) and terminal point (-5, 7).

Sol. Let \vec{AB} be the vector with initial point A(2, 1) and terminal point B(-5, 7).

\Rightarrow P.V. (Position Vector) of point A is $(2, 1) = 2\hat{i} + \hat{j}$ and P.V. of point B is $(-5, 7) = -5\hat{i} + 7\hat{j}$.

$$\begin{aligned}\therefore \vec{AB} &= \text{P.V. of point B} - \text{P.V. of point A} \\ &= (-5\hat{i} + 7\hat{j}) - (2\hat{i} + \hat{j}) = -5\hat{i} + 7\hat{j} - 2\hat{i} - \hat{j} \\ \Rightarrow \vec{AB} &= -7\hat{i} + 6\hat{j}.\end{aligned}$$

\therefore By definition, scalar components of the vectors \vec{AB} are coefficients of \hat{i} and \hat{j} in \vec{AB} i.e., -7 and 6 and vector components of the vector \vec{AB} are $-7\hat{i}$ and $6\hat{j}$.

6. Find the sum of the vectors:

$$\begin{aligned}\vec{a} &= \hat{i} - 2\hat{j} + \hat{k}, \vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k} \\ \text{and } \vec{c} &= \hat{i} - 6\hat{j} - 7\hat{k}.\end{aligned}$$

Sol. Given: $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$
and $\vec{c} = \hat{i} - 6\hat{j} - 7\hat{k}$.

Adding $\vec{a} + \vec{b} + \vec{c} = 0\hat{i} - 4\hat{j} - \hat{k} = -4\hat{j} - \hat{k}$.

7. Find the unit vector in the direction of the vector

$$\vec{a} = \hat{i} + \hat{j} + 2\hat{k}.$$

Sol. We know that a unit vector in the direction of the vector

$$\begin{aligned}\vec{a} = \hat{i} + \hat{j} + 2\hat{k} \text{ is } \hat{a} &= \frac{\vec{a}}{|\vec{a}|} = \frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{1+1+4}} \\ \Rightarrow \hat{a} &= \frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}} = \frac{1}{\sqrt{6}}\hat{i} + \frac{1}{\sqrt{6}}\hat{j} + \frac{2}{\sqrt{6}}\hat{k}.\end{aligned}$$

8. Find the unit vector in the direction of the vector \vec{PQ} where P and Q are the points (1, 2, 3) and (4, 5, 6) respectively.

Sol. Because points P and Q are P(1, 2, 3) and Q(4, 5, 6) (given),

therefore, position vector of point P = $\vec{OP} = 1\hat{i} + 2\hat{j} + 3\hat{k}$

and position vector of point Q = $\vec{OQ} = 4\hat{i} + 5\hat{j} + 6\hat{k}$
where O is the origin.

$$\begin{aligned}\therefore \vec{PQ} &= \text{Position vector of point Q} - \text{Position vector of point P} \\ &= \vec{OQ} - \vec{OP} = 3\hat{i} + 3\hat{j} + 3\hat{k}\end{aligned}$$

Therefore, a unit vector in the direction of vector \vec{PQ}

$$\begin{aligned}&= \frac{\vec{PQ}}{|\vec{PQ}|} = \frac{3\hat{i} + 3\hat{j} + 3\hat{k}}{\sqrt{9+9+9} = 27 = 9 \times 3}\end{aligned}$$

$$= \frac{3(\hat{i} + \hat{j} + \hat{k})}{3\sqrt{3}} = \frac{(\hat{i} + \hat{j} + \hat{k})}{\sqrt{3}} = \frac{1}{\sqrt{3}} \hat{i} + \frac{1}{\sqrt{3}} \hat{j} + \frac{1}{\sqrt{3}} \hat{k}.$$

9. For given vectors $\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + \hat{j} - \hat{k}$;
find the unit vector in the direction of $\vec{a} + \vec{b}$.

Sol. Given: Vectors $\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + \hat{j} - \hat{k}$

$$\therefore \vec{a} + \vec{b} = 2\hat{i} - \hat{j} + 2\hat{k} - \hat{i} + \hat{j} - \hat{k} = \hat{i} + 0\hat{j} + \hat{k}$$

$$\therefore |\vec{a} + \vec{b}| = \sqrt{(1)^2 + (0)^2 + (1)^2} = \sqrt{2}$$

\therefore A unit vector in the direction of $\vec{a} + \vec{b}$ is

$$\frac{\vec{a} + \vec{b}}{|\vec{a} + \vec{b}|} = \frac{\hat{i} + 0\hat{j} + \hat{k}}{\sqrt{2}} = \frac{\hat{i} + \hat{k}}{\sqrt{2}} = \frac{1}{\sqrt{2}} \hat{i} + \frac{1}{\sqrt{2}} \hat{k}.$$

10. Find a vector in the direction of vector $5\hat{i} - \hat{j} + 2\hat{k}$ which has magnitude 8 units.

Sol. Let $\vec{a} = 5\hat{i} - \hat{j} + 2\hat{k}$.

\therefore A vector in the direction of vector \vec{a} which has magnitude 8 units

$$\begin{aligned} &= 8\hat{a} = 8 \frac{\vec{a}}{|\vec{a}|} = \frac{8(5\hat{i} - \hat{j} + 2\hat{k})}{\sqrt{25 + 1 + 4}} \\ &= \frac{8}{\sqrt{30}} (5\hat{i} - \hat{j} + 2\hat{k}) = \frac{40}{\sqrt{30}} \hat{i} - \frac{8}{\sqrt{30}} \hat{j} + \frac{16}{\sqrt{30}} \hat{k}. \end{aligned}$$

11. Show that the vectors $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $-4\hat{i} + 6\hat{j} - 8\hat{k}$ are collinear.

Sol. Let $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$

$$\text{and } \vec{b} = -4\hat{i} + 6\hat{j} - 8\hat{k}$$


$$= -2(2\hat{i} - 3\hat{j} + 4\hat{k}) = -2\vec{a}$$

$$\Rightarrow \vec{b} = -2\vec{a} = m\vec{a} \text{ where } m = -2 < 0$$

\therefore Vectors \vec{a} and \vec{b} are collinear (unlike because $m = -2 < 0$).

12. Find the direction cosines of the vector $\hat{i} + 2\hat{j} + 3\hat{k}$.

Sol. The given vector is $(\vec{a}) = \hat{i} + 2\hat{j} + 3\hat{k}$



[By (i)]

$$= \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}} = \frac{1}{\sqrt{14}} \hat{i} + \frac{2}{\sqrt{14}} \hat{j} + \frac{3}{\sqrt{14}} \hat{k}$$

We know that direction cosines of a vector \vec{a} are coefficients of \hat{i} , \hat{j} , \hat{k} in \vec{a} i.e., $\frac{1}{\sqrt{14}}$, $\frac{2}{\sqrt{14}}$, $\frac{3}{\sqrt{14}}$.

- 13. Find the direction cosines of the vector joining the points A(1, 2, -3) and B(-1, -2, 1) directed from A to B.**

Sol. Given: Points A(1, 2, -3) and B(-1, -2, 1).

$$\begin{array}{ccc} \text{A} & \xrightarrow{\quad} & \text{B} \\ (1, 2, -3) & & (-1, -2, 1) \end{array}$$

\Rightarrow P.V. (Position Vector, \vec{OA}) of point A is $A(1, 2, -3) = \hat{i} + 2\hat{j} - 3\hat{k}$

and P.V. of point B is $B(-1, -2, 1) = -\hat{i} - 2\hat{j} + \hat{k}$.

\therefore Vector \vec{AB} (directed from A to B)

= P.V. of point B - P.V. of point A

$$= -\hat{i} - 2\hat{j} + \hat{k} - (\hat{i} + 2\hat{j} - 3\hat{k})$$

$$= -\hat{i} - 2\hat{j} + \hat{k} - \hat{i} - 2\hat{j} + 3\hat{k} = -2\hat{i} - 4\hat{j} + 4\hat{k}$$

$$\therefore AB = |\vec{AB}| = \sqrt{(-2)^2 + (-4)^2 + 4^2} = \sqrt{4 + 16 + 16} = 6$$

$$\therefore \text{A unit vector along } \vec{AB} = \frac{\vec{AB}}{|\vec{AB}|}$$

$$= \frac{-2\hat{i} - 4\hat{j} + 4\hat{k}}{6} = -\frac{2}{6}\hat{i} - \frac{4}{6}\hat{j} + \frac{4}{6}\hat{k} = -\frac{1}{3}\hat{i} - \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k}.$$

We know that Direction Cosines of the vector \vec{AB} are the coefficients of \hat{i} , \hat{j} , \hat{k} in a unit vector along \vec{AB} i.e., $-\frac{1}{3}$, $-\frac{2}{3}$, $\frac{2}{3}$.

- 14. Show that the vector $\hat{i} + \hat{j} + \hat{k}$ is equally inclined to the axes OX, OY and OZ.**


Sol. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$.

Let us find angle θ_1 (say) between

vector \vec{a} and OX ($\Rightarrow \hat{i}$)

($\because \hat{i}$ represents OX in vector form)

$$\therefore \cos \theta_1 = \frac{\vec{a} \cdot \hat{i}}{|\vec{a}| |\hat{i}|}$$



$$\Rightarrow \cos \theta_1 = \frac{(\hat{i} + \hat{j} + \hat{k}) \cdot (\hat{i} + 0\hat{j} + 0\hat{k})}{|\hat{i} + \hat{j} + \hat{k}| |\hat{i} + 0\hat{j} + 0\hat{k}|}$$

$$\Rightarrow \cos \theta_1 = \frac{1(1) + 1(0) + 1(0)}{\sqrt{1+1+1} \sqrt{1+0+0}} = \frac{1}{\sqrt{3}} \Rightarrow \theta_1 = \cos^{-1} \frac{1}{\sqrt{3}}$$

Similarly, angle θ_2 between vectors \vec{a} and \hat{j} (OY) is $\cos^{-1} \frac{1}{\sqrt{3}}$

and angle θ_3 between vectors \vec{a} and \hat{k} (OZ) is also $\cos^{-1} \frac{1}{\sqrt{3}}$.

$$\therefore \theta_1 = \theta_2 = \theta_3.$$

\therefore Vectors $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ is equally inclined to OX, OY and OZ

- 15. Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are**

$\hat{i} + 2\hat{j} - \hat{k}$ and $-\hat{i} + \hat{j} + \hat{k}$ respectively, in the ratio 2 : 1

(i) internally

(ii) externally.

Sol. P.V. of point P is $\vec{a} = \hat{i} + 2\hat{j} - \hat{k}$

and P.V. of point Q is $\vec{b} = -\hat{i} + \hat{j} + \hat{k}$

(i) Therefore P.V. of point R dividing PQ internally (i.e., R lies within the segment PQ) in the ratio 2 : 1 ($= m : n$) ($= PR : QR$)

is $\frac{m\vec{b} + n\vec{a}}{m + n}$

$$= \frac{2(-\hat{i} + \hat{j} + \hat{k}) + \hat{i} + 2\hat{j} - \hat{k}}{2 + 1} = \frac{-2\hat{i} + 2\hat{j} + 2\hat{k} + \hat{i} + 2\hat{j} - \hat{k}}{3}$$

$$= \frac{-\hat{i} + 4\hat{j} + \hat{k}}{3} = -\frac{1}{3} \hat{i} + \frac{4}{3} \hat{j} + \frac{1}{3} \hat{k}.$$

(ii) P.V. of point R dividing PQ externally (i.e., R lies outside PQ and to the right of point Q because ratio 2 : 1 = $\frac{2}{1} > 1$ as PR is

$$2 \text{ times } PQ \text{ i.e., } \frac{PR}{QR} = \frac{2}{1}) \text{ is } \frac{m\vec{b} - n\vec{a}}{m - n}$$

$$= \frac{2(-\hat{i} + \hat{j} + \hat{k}) - (\hat{i} + 2\hat{j} - \hat{k})}{2 - 1}$$

$$= -2\hat{i} + 2\hat{j} + 2\hat{k} - \hat{i} - 2\hat{j} + \hat{k} = -3\hat{i} + \hat{k}.$$

Remark. In the above question 15(ii), had R been dividing PQ externally in the ratio 1 : 2; then R will lie to the left of point P

$$\text{and } \frac{PR}{QR} = \frac{1}{2}.$$

16. Find the position vector of the mid-point of the vector joining the points P(2, 3, 4) and Q(4, 1, -2).

Sol. Given: Point P is (2, 3, 4) and Q is (4, 1, -2).

$$\Rightarrow \text{P.V. of point P(2, 3, 4) is } \vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\text{and P.V. of point Q(4, 1, -2) is } \vec{b} = 4\hat{i} + \hat{j} - 2\hat{k}.$$

$$\therefore \text{P.V. of mid-point R of PQ is } \frac{\vec{a} + \vec{b}}{2}.$$

[By Formula of Internal division]

$$= \frac{2\hat{i} + 3\hat{j} + 4\hat{k} + 4\hat{i} + \hat{j} - 2\hat{k}}{2} = \frac{6\hat{i} + 4\hat{j} + 2\hat{k}}{2} = 3\hat{i} + 2\hat{j} + \hat{k}.$$

17. Show that the points A, B and C with position vectors,

$$\vec{a} = 3\hat{i} - 4\hat{j} - 4\hat{k}, \vec{b} = 2\hat{i} - \hat{j} + \hat{k} \text{ and } \vec{c} = \hat{i} - 3\hat{j} - 5\hat{k},$$

respectively form the vertices of a right-angled triangle.

Sol. Given: P.V. of points A, B, C respectively are \vec{a} ($= \vec{OA}$) $= 3\hat{i} - 4\hat{j} - 4\hat{k}$,
 \vec{b} ($= \vec{OB}$) $= 2\hat{i} - \hat{j} + \hat{k}$ and \vec{c} ($= \vec{OC}$) $= \hat{i} - 3\hat{j} - 5\hat{k}$, where O is the origin.

Step I. $\therefore \vec{AB} = \text{P.V. of point B} - \text{P.V. of point A}$

$$= 2\hat{i} - \hat{j} + \hat{k} - (3\hat{i} - 4\hat{j} - 4\hat{k}) = 2\hat{i} - \hat{j} + \hat{k} - 3\hat{i} + 4\hat{j} + 4\hat{k}$$

$$\text{or } \vec{AB} = -\hat{i} + 3\hat{j} + 5\hat{k} \quad \dots(i)$$

$\vec{BC} = \text{P.V. of point C} - \text{P.V. of point B}$

$$= (\hat{i} - 3\hat{j} - 5\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = \hat{i} - 3\hat{j} - 5\hat{k} - 2\hat{i} + \hat{j} - \hat{k}$$

$$= -\hat{i} - 2\hat{j} - 6\hat{k} \quad \dots(ii)$$

$\vec{AC} = \text{P.V. of point C} - \text{P.V. of point A}$

$$= \hat{i} - 3\hat{j} - 5\hat{k} - (3\hat{i} - 4\hat{j} - 4\hat{k}) = \hat{i} - 3\hat{j} - 5\hat{k} - 3\hat{i} + 4\hat{j} + 4\hat{k}$$

$$= -2\hat{i} + \hat{j} - \hat{k} \quad \dots(iii)$$

Adding (i) and (ii),

$$\vec{AB} + \vec{BC} = -\hat{i} + 3\hat{j} + 5\hat{k} - \hat{i} - 2\hat{j} - 6\hat{k}$$

$$-2\hat{i} + \hat{j} - \hat{k} = \vec{AC} \quad [\text{By (iii)}]$$

\therefore By Triangle Law of addition of Vectors, Points A, B, C are the Vertices of a triangle or points A, B, C are collinear.

Step II.

$$\text{From (i) } AB = |\vec{AB}| = \sqrt{1 + 9 + 25} = \sqrt{35}$$

From (ii), $BC = |\vec{BC}| = \sqrt{1+4+36} = \sqrt{41}$

From (iii), $AC = |\vec{AC}| = \sqrt{4+1+1} = \sqrt{6}$

We can observe that $(\text{Longest side } BC)^2 = (\sqrt{41})^2 = 41 = 35 + 6$
 $= AB^2 + AC^2$

\therefore Points A, B, C are the vertices of a right-angled triangle.


18. In triangle ABC (Fig. below), which of the following is not true:

(A) $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$

(B) $\vec{AB} + \vec{BC} - \vec{AC} = \vec{0}$

(C) $\vec{AB} + \vec{BC} - \vec{CA} = \vec{0}$

(D) $\vec{AB} - \vec{CB} + \vec{CA} = \vec{0}$



Sol. Option (C) is not true.

Because we know by Triangle Law of Addition of vectors that

$$\begin{array}{l} \vec{AB} + \vec{BC} = \vec{AC}, \text{ i.e., } \boxed{\vec{AB} + \vec{BC} = \vec{AC}} \\ \Rightarrow \vec{AB} + \vec{BC} - \vec{AC} = \vec{0} \quad \Rightarrow \boxed{\vec{AB} + \vec{BC} - \vec{CA} = \vec{0}} \end{array}$$

But for option (C), $\vec{AB} + \vec{BC} - \vec{CA} = \vec{AC} + \vec{AC} = 2\vec{AC} \neq \vec{0}$.

Option (D) is same as option (A).


19. If \vec{a} and \vec{b} are two collinear vectors, then which of the following are incorrect:

(A) $\vec{b} = \lambda \vec{a}$, for some scalar λ . (B) $\vec{a} = \pm \vec{b}$

(C) the respective components of \vec{a} and \vec{b} are proportional

(D) both the vectors \vec{a} and \vec{b} have same direction, but different magnitudes.

Sol. Option (D) is not true because two collinear vectors can have **different** directions and also different magnitudes.



The options (A) and (C) are true by definition of collinear vectors.

Option (B) is a particular case of option (A) (taking $\lambda = \pm 1$).



Exercise 10.3

1. Find the angle between two vectors \vec{a} and \vec{b} with magnitude $\sqrt{3}$ and 2, respectively having $\vec{a} \cdot \vec{b} = \sqrt{6}$.

Sol. Given: $|\vec{a}| = \sqrt{3}$, $|\vec{b}| = 2$ and $\vec{a} \cdot \vec{b} = \sqrt{6}$

Let θ be the angle between the vectors \vec{a} and \vec{b} . We know that

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

Putting values, $\cos \theta = \frac{\sqrt{6}}{\sqrt{3}(2)}$

$$= \frac{\sqrt{6}}{\sqrt{3}\sqrt{4}} = \frac{\sqrt{6}}{\sqrt{12}} = \sqrt{\frac{6}{12}} = \sqrt{\frac{1}{2}} = \frac{1}{\sqrt{2}} = \cos \frac{\pi}{4} \quad \therefore \theta = \frac{\pi}{4}.$$

- 2. Find the angle between the vectors $\hat{i} - 2\hat{j} + 3\hat{k}$ and $3\hat{i} - 2\hat{j} + \hat{k}$.**

Sol. Given: Let $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + \hat{k}$.

$$\therefore |\vec{a}| = \sqrt{1+4+9} = \sqrt{14} \quad \because |x\hat{i} + y\hat{j} + z\hat{k}| = \sqrt{x^2 + y^2 + z^2}$$

$$\text{and } |\vec{b}| = \sqrt{9+4+1} = \sqrt{14}$$

$$\begin{aligned} \text{Also, } \vec{a} \cdot \vec{b} &= \text{Product of coefficients of } \hat{i} + \text{Product of coefficient of } \hat{j} \\ &+ \text{Product of coefficients of } \hat{k} \\ &= 1(3) + (-2)(-2) + 3(1) = 3 + 4 + 3 = 10 \end{aligned}$$

Let θ be the angle between the vectors \vec{a} and \vec{b} .

$$\text{We know that } \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{10}{\sqrt{14}\sqrt{14}} = \frac{10}{14} = \frac{5}{7}$$

$$\therefore \theta = \cos^{-1} \frac{5}{7}.$$

- 3. Find the projection of the vector $\hat{i} - \hat{j}$ on the vector $\hat{i} + \hat{j}$.**

Sol. Let $\vec{a} = \hat{i} - \hat{j} = \hat{i} - \hat{j} + 0\hat{k}$



$$\text{and } \vec{b} = \hat{i} + \hat{j} = \hat{i} + \hat{j} + 0\hat{k}$$

Projection of vector \vec{a} and \vec{b}

$$= \text{Length LM} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$

$$= \frac{(1)(1) + (-1)(1) + 0(0)}{\sqrt{(1)^2 + (1)^2 + 0^2}} = \frac{1-1+0}{\sqrt{2}} = \frac{0}{\sqrt{2}} = 0.$$

Remark. If projection of vector \vec{a} on \vec{b} is zero, then vector \vec{a} is perpendicular to vector \vec{b} .



4. Find the projection of the vector $\hat{i} + 3\hat{j} + 7\hat{k}$ on the

vector $7\hat{i} - \hat{j} + 8\hat{k}$.

Sol. Let $\vec{a} = \hat{i} + 3\hat{j} + 7\hat{k}$ and $\vec{b} = 7\hat{i} - \hat{j} + 8\hat{k}$

$$\begin{aligned} \text{We know that projection of vector } \vec{a} \text{ on vector } \vec{b} &= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \\ &= \frac{1(7) + 3(-1) + 7(8)}{\sqrt{(7)^2 + (-1)^2 + (8)^2}} = \frac{7 - 3 + 56}{\sqrt{49 + 1 + 64}} = \frac{60}{\sqrt{114}}. \end{aligned}$$

5. Show that each of the given three vectors is a unit vector: $\frac{1}{7} (2\hat{i} + 3\hat{j} + 6\hat{k})$, $\frac{1}{7} (3\hat{i} - 6\hat{j} + 2\hat{k})$,

$$\frac{1}{7} (6\hat{i} + 2\hat{j} - 3\hat{k}).$$

Also show that they are mutually perpendicular to each other.

Sol. Let $\vec{a} = \frac{1}{7} (2\hat{i} + 3\hat{j} + 6\hat{k}) = \frac{2}{7}\hat{i} + \frac{3}{7}\hat{j} + \frac{6}{7}\hat{k} \quad \dots(i)$

$$\vec{b} = \frac{1}{7} (3\hat{i} - 6\hat{j} + 2\hat{k}) = \frac{3}{7}\hat{i} - \frac{6}{7}\hat{j} + \frac{2}{7}\hat{k} \quad \dots(ii)$$

$$\vec{c} = \frac{1}{7} (6\hat{i} + 2\hat{j} - 3\hat{k}) = \frac{6}{7}\hat{i} + \frac{2}{7}\hat{j} - \frac{3}{7}\hat{k} \quad \dots(iii)$$

$$\begin{aligned} \therefore |\vec{a}| &= \sqrt{\left(\frac{2}{7}\right)^2 + \left(\frac{3}{7}\right)^2 + \left(\frac{6}{7}\right)^2} = \sqrt{\frac{4}{49} + \frac{9}{49} + \frac{36}{49}} \\ &= \sqrt{\frac{49}{49}} = \sqrt{1} = 1 \end{aligned}$$

$$\begin{aligned} |\vec{b}| &= \sqrt{\left(\frac{3}{7}\right)^2 + \left(-\frac{6}{7}\right)^2 + \left(\frac{2}{7}\right)^2} = \sqrt{\frac{9}{49} + \frac{36}{49} + \frac{4}{49}} = \sqrt{\frac{49}{49}} \\ &= \sqrt{1} = 1 \end{aligned}$$

$$\begin{aligned} |\vec{c}| &= \sqrt{\left(\frac{6}{7}\right)^2 + \left(\frac{2}{7}\right)^2 + \left(-\frac{3}{7}\right)^2} = \sqrt{\frac{36}{49} + \frac{4}{49} + \frac{9}{49}} = \sqrt{\frac{49}{49}} \\ &= \sqrt{1} = 1 \end{aligned}$$

\therefore Each of the three given vectors \vec{a} , \vec{b} , \vec{c} is a unit vector.
From (i) and (ii),

$$\begin{aligned} \vec{a} \cdot \vec{b} &= \left(\frac{2}{7}\right) \cdot \left(\frac{3}{7}\right) + \left(\frac{3}{7}\right) \left(-\frac{6}{7}\right) + \left(\frac{6}{7}\right) \left(\frac{2}{7}\right) \\ [\vec{a} \cdot \vec{b} &= a_1b_1 + a_2b_2 + a_3b_3] \end{aligned}$$

$$= \frac{6}{49} - \frac{18}{49} + \frac{12}{49} = \frac{6-18+12}{49} = \frac{0}{49} = 0$$

$\therefore \vec{a}$ and \vec{b} are perpendicular to each other.

From (ii) and (iii),

$$\begin{aligned}\vec{b} \cdot \vec{c} &= \left(\frac{3}{7}\right) \left(\frac{6}{7}\right) + \left(\frac{-6}{7}\right) \left(\frac{2}{7}\right) + \frac{2}{7} \left(\frac{-3}{7}\right) \\ &= \frac{18}{49} - \frac{12}{49} - \frac{6}{49} = \frac{18-12-6}{49} = \frac{0}{49} = 0\end{aligned}$$

$\therefore \vec{b}$ and \vec{c} are perpendicular to each other.

From (i) and (iii),

$$\begin{aligned}\vec{a} \cdot \vec{c} &= \frac{2}{7} \left(\frac{6}{7}\right) + \frac{3}{7} \left(\frac{2}{7}\right) + \left(\frac{6}{7}\right) \left(\frac{-3}{7}\right) \\ &= \frac{12}{49} + \frac{6}{49} - \frac{18}{49} = \frac{12+6-18}{49} = \frac{0}{49} = 0\end{aligned}$$

$\therefore \vec{a}$ and \vec{c} are perpendicular to each other.

Hence, \vec{a} , \vec{b} , \vec{c} are mutually perpendicular vectors.

6. Find $|\vec{a}|$ and $|\vec{b}|$, if $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 8$ and $|\vec{a}| = 8|\vec{b}|$.

Sol. Given: $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 8$ and $|\vec{a}| = 8|\vec{b}|$... (i)

$$\Rightarrow \vec{a} \cdot \vec{a} - \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} - \vec{b} \cdot \vec{b} = 8$$

$$\Rightarrow |\vec{a}|^2 - \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{b} - |\vec{b}|^2 = 8$$

$$[\because \text{We know that } \vec{a} \cdot \vec{a} = |\vec{a}|^2 \text{ and } \vec{b} \cdot \vec{b} = |\vec{b}|^2 \text{ and } \vec{b} \cdot \vec{a} = \vec{a} \cdot \vec{b}]$$

$$\Rightarrow |\vec{a}|^2 - |\vec{b}|^2 = 8 \quad \dots (ii)$$

Putting $|\vec{a}| = 8|\vec{b}|$ from (i) in (ii), $64|\vec{b}|^2 - |\vec{b}|^2 = 8$

$$\text{or } (64 - 1)|\vec{b}|^2 = 8 \quad \Rightarrow 63|\vec{b}|^2 = 8$$

$$\Rightarrow |\vec{b}|^2 = \frac{8}{63} \quad \Rightarrow |\vec{b}| = \sqrt{\frac{8}{63}} = \sqrt{\frac{4 \times 2}{9 \times 7}}$$

(\because Length i.e., modulus of a vector is never negative.)

$$\Rightarrow |\vec{b}| = \frac{2}{3} \sqrt{\frac{2}{7}}$$

Putting this value of $|\vec{b}|$ in (i),

$$|\vec{a}| = 8 \left(\frac{2}{3} \sqrt{\frac{2}{7}} \right) = \frac{16}{3} \sqrt{\frac{2}{7}}.$$

7. Evaluate the product $(3\vec{a} - 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$.

Sol. The given expression $= (3\vec{a} - 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$

$$= (3\vec{a}) \cdot (2\vec{a}) + (3\vec{a}) \cdot (7\vec{b}) - (5\vec{b}) \cdot (2\vec{a}) - (5\vec{b}) \cdot (7\vec{b})$$

$$= 6\vec{a} \cdot \vec{a} + 21\vec{a} \cdot \vec{b} - 10\vec{b} \cdot \vec{a} - 35\vec{b} \cdot \vec{b}$$

$$= 6|\vec{a}|^2 + 21\vec{a} \cdot \vec{b} - 10\vec{a} \cdot \vec{b} - 35|\vec{b}|^2$$

$$[\because \vec{a} \cdot \vec{a} = |\vec{a}|^2 \text{ and } \vec{b} \cdot \vec{b} = |\vec{b}|^2 \text{ and } \vec{b} \cdot \vec{a} = \vec{a} \cdot \vec{b}]$$

$$= 6|\vec{a}|^2 + 11\vec{a} \cdot \vec{b} - 35|\vec{b}|^2.$$

8. Find the magnitude of two vectors \vec{a} and \vec{b} , having the same magnitude such that the angle between them is 60° and their scalar product is $\frac{1}{2}$.

Sol. Given: $|\vec{a}| = |\vec{b}|$ and angle θ (say) between \vec{a} and \vec{b} is 60° and their scalar (i.e., dot) product $= \frac{1}{2}$

i.e., $\vec{a} \cdot \vec{b} = \frac{1}{2}$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta = \frac{1}{2} \quad [\because \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta]$$

Putting $|\vec{b}| = |\vec{a}|$ (given) and $\theta = 60^\circ$ (given), we have

$$|\vec{a}| |\vec{a}| \cos 60^\circ = \frac{1}{2} \Rightarrow |\vec{a}|^2 \left(\frac{1}{2} \right) = \frac{1}{2}$$

Multiplying by 2, $|\vec{a}|^2 = 1 \Rightarrow |\vec{a}| = 1$... (i)

(\because Length of a vector is never negative)

$\therefore |\vec{b}| = |\vec{a}| = 1$ [By (i)]

$\therefore |\vec{a}| = 1$ and $|\vec{b}| = 1$.

9. Find $|\vec{x}|$, if for a unit vector \vec{a} , $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$.

Sol. Given: \vec{a} is a unit vector $\Rightarrow |\vec{a}| = 1$... (i)

Also given $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$

$$\Rightarrow \vec{x} \cdot \vec{x} + \vec{x} \cdot \vec{a} - \vec{a} \cdot \vec{x} - \vec{a} \cdot \vec{a} = 12$$

$$\Rightarrow |\vec{x}|^2 + \vec{a} \cdot \vec{x} - \vec{a} \cdot \vec{x} - |\vec{a}|^2 = 12$$

$$\Rightarrow |\vec{x}|^2 - |\vec{a}|^2 = 12$$

Putting $|\vec{a}| = 1$ from (i), $|\vec{x}|^2 - 1 = 12$

$$\Rightarrow |\vec{x}|^2 = 13 \quad \Rightarrow |\vec{x}| = \sqrt{13}.$$

(\because Length of a vector is never negative.)

10. If $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 3\hat{i} + \hat{j}$ are such that $\vec{a} + \lambda\vec{b}$ is perpendicular to \vec{c} , then find the value of λ .

Sol. Given : $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$, $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$

and $\vec{c} = 3\hat{i} + \hat{j}$.

Now, $\vec{a} + \lambda\vec{b} = 2\hat{i} + 2\hat{j} + 3\hat{k} + \lambda(-\hat{i} + 2\hat{j} + \hat{k})$

$$= 2\hat{i} + 2\hat{j} + 3\hat{k} - \lambda\hat{i} + 2\lambda\hat{j} + \lambda\hat{k}$$

$$\Rightarrow \vec{a} + \lambda\vec{b} = (2 - \lambda)\hat{j} + (2 + 2\lambda)\hat{j} + (3 + \lambda)\hat{k}$$

Again given $\vec{c} = 3\hat{i} + \hat{j} = 3\hat{i} + \hat{j} + 0\hat{k}$.

Because vector $\vec{a} + \lambda\vec{b}$ is perpendicular to \vec{c} , therefore,

$$(\vec{a} + \lambda\vec{b}) \cdot \vec{c} = 0$$

i.e., Product of coefficients of $\hat{i} + \dots\dots\dots = 0$

$$\Rightarrow (2 - \lambda)3 + (2 + 2\lambda)1 + (3 + \lambda)0 = 0$$

$$\Rightarrow 6 - 3\lambda + 2 + 2\lambda = 0 \quad \Rightarrow -\lambda + 8 = 0$$

$$\Rightarrow -\lambda = -8 \quad \Rightarrow \lambda = 8.$$

11. Show that $|\vec{a}| |\vec{b}| + |\vec{b}| |\vec{a}|$ is perpendicular to $|\vec{a}| |\vec{b}| - |\vec{b}| |\vec{a}|$, for any two non-zero vectors \vec{a} and \vec{b} .

Sol. Let $\vec{c} = |\vec{a}| |\vec{b}| + |\vec{b}| |\vec{a}| = l\vec{b} + m\vec{a}$

where $l = |\vec{a}|$ and $m = |\vec{b}|$

Let $\vec{d} = |\vec{a}| |\vec{b}| - |\vec{b}| |\vec{a}| = l\vec{b} - m\vec{a}$

Now, $\vec{c} \cdot \vec{d} = (l\vec{b} + m\vec{a}) \cdot (l\vec{b} - m\vec{a})$

$$= l^2 \vec{b} \cdot \vec{b} - lm \vec{b} \cdot \vec{a} + lm \vec{a} \cdot \vec{b} - m^2 \vec{a} \cdot \vec{a}$$

$$= l^2 |\vec{b}|^2 - lm \vec{a} \cdot \vec{b} + lm \vec{a} \cdot \vec{b} - m^2 |\vec{a}|^2 = l^2 |\vec{b}|^2 - m^2 |\vec{a}|^2$$

Putting $l = |\vec{a}|$ and $m = |\vec{b}|$,

$$= |\vec{a}|^2 |\vec{b}|^2 - |\vec{b}|^2 |\vec{a}|^2 = 0$$

i.e., $\vec{c} \cdot \vec{d} = 0$

\therefore Vectors \vec{c} and \vec{d} are perpendicular to each other.

12. If $\vec{a} \cdot \vec{a} = 0$ and $\vec{a} \cdot \vec{b} = 0$, then what can be concluded about the vector \vec{b} ?

Sol. Given: $\vec{a} \cdot \vec{a} = 0 \Rightarrow |\vec{a}|^2 = 0 \Rightarrow |\vec{a}| = 0 \quad \dots(i)$

($\Rightarrow \vec{a}$ is a zero vector by definition of zero vector.)

Again given $\vec{a} \cdot \vec{b} = 0 \Rightarrow |\vec{a}| |\vec{b}| \cos \theta = 0$

Putting $|\vec{a}| = 0$ from (i), we have $0 |\vec{b}| \cos \theta = 0$

i.e., $0 = 0$ for all (any) vectors \vec{b} . $\therefore \vec{b}$ can be any vector.

Note. $(\vec{a} + \vec{b} + \vec{c})^2 = (\vec{a} + (\vec{b} + \vec{c}))^2$

$$= \vec{a}^2 + (\vec{b} + \vec{c})^2 + 2\vec{a} \cdot (\vec{b} + \vec{c})$$

$$[\because (\vec{A} + \vec{B})^2 = \vec{A}^2 + \vec{B}^2 + 2\vec{A} \cdot \vec{B}]$$

$$= \vec{a}^2 + \vec{b}^2 + \vec{c}^2 + 2\vec{b} \cdot \vec{c} + 2\vec{a} \cdot \vec{b} + 2\vec{a} \cdot \vec{c}$$

Using $\vec{a} \cdot \vec{c} = \vec{c} \cdot \vec{a}$

$$\text{or } (\vec{a} + \vec{b} + \vec{c})^2 = \vec{a}^2 + \vec{b}^2 + \vec{c}^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

13. If $\vec{a}, \vec{b}, \vec{c}$ are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, find the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$.

Sol. Because $\vec{a}, \vec{b}, \vec{c}$ are unit vectors, therefore,

$$|\vec{a}| = 1, |\vec{b}| = 1 \text{ and } |\vec{c}| = 1. \quad \dots(i)$$

Again given $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

Squaring both sides $(\vec{a} + \vec{b} + \vec{c})^2 = 0$

Using formula of **Note** above

$$\Rightarrow \vec{a}^2 + \vec{b}^2 + \vec{c}^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\text{or } |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

Putting $|\vec{a}| = 1, |\vec{b}| = 1, |\vec{c}| = 1$ from (i),

$$1 + 1 + 1 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = -3$$

Dividing both sides by 2, $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = \frac{-3}{2}$.

14. If either vector $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then $\vec{a} \cdot \vec{b} = 0$. But the converse need not be true. Justify your answer with an example.

Sol. Case I. Vector $\vec{a} = \vec{0}$. Therefore, by definition of zero vector,

$$|\vec{a}| = 0 \quad \dots(i)$$

$$\therefore \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = 0 (|\vec{b}| \cos \theta) \quad [\text{By (i)}]$$

$$= 0$$

Case II. Vector $\vec{b} = \vec{0}$. Proceeding as above we can prove that

$$\vec{a} \cdot \vec{b} = 0$$

But the converse is not true.

Let us justify it with an example.

$$\text{Let } \vec{a} = \hat{i} + \hat{j} + \hat{k}. \text{ Therefore, } |\vec{a}| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3} \neq 0.$$

Therefore $\vec{a} \neq \vec{0}$ (By definition of Zero Vector)

$$\text{Let } \vec{b} = \hat{i} + \hat{j} - 2\hat{k}.$$

$$\text{Therefore, } |\vec{b}| = \sqrt{(1)^2 + (1)^2 + (-2)^2} = \sqrt{6} \neq 0.$$


Therefore, $\vec{b} \neq \vec{0}$.

$$\text{But } \vec{a} \cdot \vec{b} = 1(1) + 1(1) + 1(-2) = 1 + 1 - 2 = 0$$

So here $\vec{a} \cdot \vec{b} = 0$ but neither $\vec{a} = \vec{0}$ nor $\vec{b} = \vec{0}$.

- 15. If the vertices A, B, C of a triangle ABC are (1, 2, 3), (-1, 0, 0) and (0, 1, 2), respectively, then find $\angle ABC$.**

Sol. Given: Vertices A, B, C of a triangle are A(1, 2, 3), B(-1, 0, 0) and C(0, 1, 2) respectively.



$$\therefore \text{Position vector (P.V.) of point A (} = \vec{OA} \text{)} = (1, 2, 3)$$

$$= \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\text{Position vector (P.V.) of point B (} = \vec{OB} \text{)} = (-1, 0, 0)$$

$$= -\hat{i} + 0\hat{j} + 0\hat{k}$$

$$\text{and position vector (P.V.) of point C (} = \vec{OC} \text{)} = (0, 1, 2)$$

$$= 0\hat{i} + \hat{j} + 2\hat{k}$$

We can see from the above figure that $\angle ABC$ is the angle between the vectors \vec{BA} and \vec{BC}

$$\begin{aligned}
 \text{Now } \vec{BA} &= \text{P.V. of terminal point A} - \text{P.V. of initial point B} \\
 &= \hat{i} + 2\hat{j} + 3\hat{k} - (-\hat{i} + 0\hat{j} + 0\hat{k}) \\
 &= \hat{i} + 2\hat{j} + 3\hat{k} + \hat{i} - 0\hat{j} - 0\hat{k} = 2\hat{i} + 2\hat{j} + 3\hat{k} \quad \dots(i)
 \end{aligned}$$

$$\begin{aligned}
 \text{and } \vec{BC} &= \text{P.V. of point C} - \text{P.V. of point B} \\
 &= 0\hat{i} + \hat{j} + 2\hat{k} - (-\hat{i} + 0\hat{j} + 0\hat{k}) \\
 &= 0\hat{i} + \hat{j} + 2\hat{k} + \hat{i} - 0\hat{j} - 0\hat{k} = \hat{i} + \hat{j} + 2\hat{k} \quad \dots(ii)
 \end{aligned}$$

$$\text{We know that } \cos \angle ABC = \frac{\vec{BA} \cdot \vec{BC}}{|\vec{BA}| |\vec{BC}|} \quad \left| \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right.$$

Using (i) and (ii)

$$\begin{aligned}
 &= \frac{2(1) + 2(1) + 3(2)}{\sqrt{4+4+9} \sqrt{1+1+4}} = \frac{10}{\sqrt{17}\sqrt{6}} = \frac{10}{\sqrt{102}} \\
 \therefore \angle ABC &= \cos^{-1} \frac{10}{\sqrt{102}}.
 \end{aligned}$$

16. Show that the points A(1, 2, 7), B(2, 6, 3) and C(3, 10, -1) are collinear.

Sol. Given points are A(1, 2, 7), B(2, 6, 3) and C(3, 10, -1).

\Rightarrow P.V.'s \vec{OA} , \vec{OB} , \vec{OC} of points A, B, C are

$$\vec{OA} = (1, 2, 7) = \hat{i} + 2\hat{j} + 7\hat{k}$$

$$\vec{OB} = (2, 6, 3) = 2\hat{i} + 6\hat{j} + 3\hat{k}$$

$$\text{and } \vec{OC} = (3, 10, -1) = 3\hat{i} + 10\hat{j} - \hat{k}$$

$\therefore \vec{AB} = \text{P.V. of terminal point B} - \text{P.V. of initial point A}$

$$\begin{aligned}
 &= 2\hat{i} + 6\hat{j} + 3\hat{k} - (\hat{i} + 2\hat{j} + 7\hat{k}) \\
 &= 2\hat{i} + 6\hat{j} + 3\hat{k} - \hat{i} - 2\hat{j} - 7\hat{k} = \hat{i} + 4\hat{j} - 4\hat{k} \quad \dots(i)
 \end{aligned}$$

and $\vec{AC} = \text{P.V. of point C} - \text{P.V. of point A}$

$$\begin{aligned}
 &= 3\hat{i} + 10\hat{j} - \hat{k} - (\hat{i} + 2\hat{j} + 7\hat{k}) \\
 &= 3\hat{i} + 10\hat{j} - \hat{k} - \hat{i} - 2\hat{j} - 7\hat{k} \\
 &= 2\hat{i} + 8\hat{j} - 8\hat{k} = 2(\hat{i} + 4\hat{j} - 4\hat{k})
 \end{aligned}$$

$$\Rightarrow \vec{AC} = 2\vec{AB}$$

[By (i)]

\Rightarrow Vectors \vec{AB} and \vec{AC} are collinear or parallel. $\because \vec{a} = m \vec{b}$
 \Rightarrow Points A, B, C are collinear.

(\because Vectors \vec{AB} and \vec{AC} have a common point A and hence can't be parallel.)

Remark. When we come to exercise 10.4 and learn that Exercise, we have a second solution for proving points A, B, C to be collinear:

Prove that $\vec{AB} \times \vec{AC} = \vec{0}$.

17. Show that the vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $3\hat{i} - 4\hat{j} - 4\hat{k}$ form the vertices of a right angled triangle.

Sol. Let the given (position) vectors be P.V.'s of the points A, B, C respectively.

P.V. of point A is $2\hat{i} - \hat{j} + \hat{k}$ and

P.V. of point B is $\hat{i} - 3\hat{j} - 5\hat{k}$ and

P.V. of point C is $3\hat{i} - 4\hat{j} - 4\hat{k}$.

$$\begin{aligned} \therefore \vec{AB} &= \text{P.V. of point B} - \text{P.V. of point A} \\ &= \hat{i} - 3\hat{j} - 5\hat{k} - (2\hat{i} - \hat{j} + \hat{k}) = \hat{i} - 3\hat{j} - 5\hat{k} - 2\hat{i} + \hat{j} - \hat{k} \\ &= -\hat{i} - 2\hat{j} - 6\hat{k} \end{aligned} \quad \dots(i)$$

$$\begin{aligned} \text{and } \vec{BC} &= \text{P.V. of point C} - \text{P.V. of point B} \\ &= 3\hat{i} - 4\hat{j} - 4\hat{k} - (\hat{i} - 3\hat{j} - 5\hat{k}) = 3\hat{i} - 4\hat{j} - 4\hat{k} - \hat{i} + 3\hat{j} + 5\hat{k} \\ &= 2\hat{i} - \hat{j} + \hat{k} \end{aligned} \quad \dots(ii)$$

$$\begin{aligned} \text{and } \vec{AC} &= \text{P.V. of point C} - \text{P.V. of point A} \\ &= 3\hat{i} - 4\hat{j} - 4\hat{k} - (2\hat{i} - \hat{j} + \hat{k}) = 3\hat{i} - 4\hat{j} - 4\hat{k} - 2\hat{i} + \hat{j} - \hat{k} \\ &= \hat{i} - 3\hat{j} - 5\hat{k} \end{aligned} \quad \dots(iii)$$

Adding (i) and (ii), we have

$$\begin{aligned} \vec{AB} + \vec{BC} &= -\hat{i} - 2\hat{j} - 6\hat{k} + 2\hat{i} - \hat{j} + \hat{k} \\ &= \hat{i} - 3\hat{j} - 5\hat{k} = \vec{AC} \end{aligned} \quad [\text{By (iii)}]$$

\therefore By Triangle Law of addition of vectors, points A, B, C are the vertices of a triangle ABC or points A, B, C are collinear.

$$\begin{aligned} \text{Now from (i) and (ii), } \vec{AB} \cdot \vec{BC} &= (-1)(2) + (-2)(-1) + (-6)(1) \\ &= -2 + 2 - 6 = -6 \neq 0 \end{aligned}$$

From (ii) and (iii), $\vec{BC} \cdot \vec{AC} = 2(1) + (-1)(-3) + 1(-5)$
 $= 2 + 3 - 5 = 0$

$\Rightarrow \vec{BC}$ is perpendicular to \vec{AC}

\Rightarrow Angle C is 90° . $\therefore \triangle ABC$ is right angled at point C.

\therefore Points A, B, C are the vertices of a right angled triangle.

18. If \vec{a} is a non-zero vector of magnitude 'a' and λ is a non-zero scalar, then $\lambda \vec{a}$ is a unit vector if

(A) $\lambda = 1$ (B) $\lambda = -1$ (C) $a = |\lambda|$ (D) $a = \frac{1}{|\lambda|}$

Sol. Given: \vec{a} is a non-zero vector of magnitude a

$\Rightarrow |\vec{a}| = 1$...(i)

Also given: $\lambda \neq 0$ and $\lambda \vec{a}$ is a unit vector.

$\Rightarrow |\lambda \vec{a}| = 1 \quad \Rightarrow |\lambda| |\vec{a}| = 1$

$\Rightarrow |\lambda| a = 1 \quad \Rightarrow a = \frac{1}{|\lambda|}$

\therefore Option (D) is the correct answer.

Kopykitab

Same textbooks, click away

Exercise 10.4

1. Find $|\vec{a} \times \vec{b}|$, if $\vec{a} = \hat{i} - 7\hat{j} + 7\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 2\hat{k}$.

Sol. Given: $\vec{a} = \hat{i} - 7\hat{j} + 7\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 2\hat{k}$.

$$\text{Therefore, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -7 & 7 \\ 3 & -2 & 2 \end{vmatrix}$$

$$\left[\because \text{ If } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k} \text{ and } \vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}; \right. \\ \left. \text{then } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \right]$$

Expanding along first row,

$$\begin{aligned} \vec{a} \times \vec{b} &= \hat{i} \begin{vmatrix} -7 & 7 \\ -2 & 2 \end{vmatrix} - \hat{j} \begin{vmatrix} 1 & 7 \\ 3 & 2 \end{vmatrix} + \hat{k} \begin{vmatrix} 1 & -7 \\ 3 & -2 \end{vmatrix} \\ \Rightarrow \vec{a} \times \vec{b} &= \hat{i}(-14 + 14) - \hat{j}(2 - 21) + \hat{k}(-2 + 21) \end{aligned}$$


$$= 0\hat{i} + 19\hat{j} + 19\hat{k}$$

$$\therefore |\vec{a} \times \vec{b}| = \sqrt{0^2 + (19)^2 + (19)^2} = \sqrt{2(19)^2} = \sqrt{2}(19) = 19\sqrt{2}.$$

Result: We know that $\vec{n} = \vec{a} \times \vec{b}$ is a vector perpendicular to both the vectors \vec{a} and \vec{b} .
Therefore, a unit vector perpendicular

to both the vectors \vec{a} and \vec{b} is

$$\hat{n} = \pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} \quad \left[\because \hat{A} = \frac{\vec{A}}{|\vec{A}|} \right]$$



2. Find a unit vector perpendicular to each of the vectors

$$\vec{a} + \vec{b} \text{ and } \vec{a} - \vec{b} \text{ where } \vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k} \text{ and } \vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}.$$

Sol. Given: $\vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$

Adding, $\vec{c} = \vec{a} + \vec{b} = 4\hat{i} + 4\hat{j} + 0\hat{k}$

Subtracting $\vec{d} = \vec{a} - \vec{b} = 2\hat{i} + 0\hat{j} + 4\hat{k}$

Therefore, $\vec{n} = \vec{c} \times \vec{d} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 4 & 0 \\ 2 & 0 & 4 \end{vmatrix}$

Expanding along first row $= \hat{i}(16 - 0) - \hat{j}(16 - 0) + \hat{k}(0 - 8)$

$$\Rightarrow \vec{n} = 16\hat{i} - 16\hat{j} - 8\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{(16)^2 + (-16)^2 + (-8)^2} = \sqrt{256 + 256 + 64} = \sqrt{576} = 24.$$

Therefore, a unit vector perpendicular to both \vec{a} and \vec{b} is

$$\begin{aligned} \hat{n} &= \pm \frac{\vec{n}}{|\vec{n}|} = \pm \frac{(16\hat{i} - 16\hat{j} - 8\hat{k})}{24} \\ &= \pm \left(\frac{16}{24}\hat{i} - \frac{16}{24}\hat{j} - \frac{8}{24}\hat{k} \right) = \pm \left(\frac{2}{3}\hat{i} - \frac{2}{3}\hat{j} - \frac{1}{3}\hat{k} \right). \end{aligned}$$

3. If a unit vector \hat{a} makes an angle $\frac{\pi}{3}$ with \hat{i} , $\frac{\pi}{4}$ with \hat{j} and an acute angle θ with \hat{k} , then find θ and hence, the components of \hat{a} .

Sol. Let $\hat{a} = x\hat{i} + y\hat{j} + z\hat{k}$ be a unit vector ... (i)

$$\Rightarrow |\hat{a}| = 1 \Rightarrow \sqrt{x^2 + y^2 + z^2} = 1$$

Squaring both sides, $x^2 + y^2 + z^2 = 1$... (ii)

Given: Angle between vectors \hat{a} and $\hat{i} = \hat{i} + 0\hat{j} + 0\hat{k}$ is $\frac{\pi}{3}$.

$$\therefore \cos \frac{\pi}{3} = \frac{\hat{a} \cdot \hat{i}}{|\hat{a}| |\hat{i}|} \quad \left[\because \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right]$$

$$\Rightarrow \frac{1}{2} = \frac{x(1) + y(0) + z(0)}{(1)(1)} \quad \text{or} \quad \frac{1}{2} = x \quad \dots (iii)$$

Again, **Given:** Angle between vectors \hat{a} and $\hat{j} = 0\hat{i} + \hat{j} + 0\hat{k}$ is $\frac{\pi}{4}$.

$$\therefore \cos \frac{\pi}{4} = \frac{\hat{a} \cdot \hat{j}}{|\hat{a}| |\hat{j}|} \Rightarrow \frac{1}{\sqrt{2}} = \frac{x(0) + y(1) + z(0)}{(1)(1)}$$

$$\Rightarrow \frac{1}{\sqrt{2}} = y \quad \dots (iv)$$

Again, **Given:** Angle between vectors \hat{a} and $\hat{k} = 0\hat{i} + 0\hat{j} + \hat{k}$ is θ where θ is acute.

$$\therefore \cos \theta = \frac{\hat{a} \cdot \hat{k}}{|\hat{a}| |\hat{k}|} = \frac{x(0) + y(0) + z(1)}{(1)(1)} = z \quad \dots (v)$$

Putting values of x , y and z from (iii), (iv) and (v) in (ii),

$$\frac{1}{4} + \frac{1}{2} + \cos^2 \theta = 1$$

$$\Rightarrow \cos^2 \theta = 1 - \frac{1}{4} - \frac{1}{2} = \frac{4-1-2}{4} = \frac{1}{4} \Rightarrow \cos \theta = \pm \frac{1}{2}$$

But θ is acute angle (given)

$$\Rightarrow \cos \theta \text{ is positive and hence } = \frac{1}{2} = \cos \frac{\pi}{3} \Rightarrow \theta = \frac{\pi}{3}$$

From (v), $z = \cos \theta = \frac{1}{2}$

Putting values of x , y , z in (i), $\hat{a} = \frac{1}{2}\hat{i} + \frac{1}{\sqrt{2}}\hat{j} + \frac{1}{2}\hat{k}$

\therefore Components of \hat{a} are coefficients of \hat{i} , \hat{j} , \hat{k} in \hat{a}

i.e., $\frac{1}{2}$, $\frac{1}{\sqrt{2}}$, $\frac{1}{2}$ and acute angle $\theta = \frac{\pi}{3}$.

4. Show that $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2\vec{a} \times \vec{b}$.

Sol. L.H.S. $= (\vec{a} - \vec{b}) \times (\vec{a} + \vec{b})$
 $= \vec{a} \times \vec{a} + \vec{a} \times \vec{b} - \vec{b} \times \vec{a} - \vec{b} \times \vec{b}$
 $= \vec{0} + \vec{a} \times \vec{b} + \vec{a} \times \vec{b} - \vec{0}$
 $[\because \vec{a} \times \vec{a} = \vec{0}, \vec{b} \times \vec{b} = \vec{0} \text{ and } \vec{b} \times \vec{a} = -\vec{a} \times \vec{b}]$
 $= 2\vec{a} \times \vec{b} = \text{R.H.S.}$

5. Find λ and μ if $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + \lambda\hat{j} + \mu\hat{k}) = \vec{0}$.

Sol. Given: $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + \lambda\hat{j} + \mu\hat{k}) = \vec{0}$

$$\Rightarrow \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 6 & 27 \\ 1 & \lambda & \mu \end{vmatrix} = \vec{0}$$

Expanding along first row,

$$\hat{i}(6\mu - 27\lambda) - \hat{j}(2\mu - 27) + \hat{k}(2\lambda - 6) = \vec{0} = 0\hat{i} + 0\hat{j} + 0\hat{k}$$

Comparing coefficients of $\hat{i}, \hat{j}, \hat{k}$ on both sides, we have

$$6\mu - 27\lambda = 0 \quad \dots(i)$$

$$2\mu - 27 = 0 \quad \dots(ii)$$

$$\text{and} \quad 2\lambda - 6 = 0 \quad \dots(iii)$$

$$\text{From (ii), } 2\mu = 27 \Rightarrow \mu = \frac{27}{2}$$

$$\text{From (iii), } 2\lambda = 6 \Rightarrow \lambda = \frac{6}{2} = 3$$

$$\text{Putting } \lambda = 3 \text{ and } \mu = \frac{27}{2} \text{ in (i), } 6\left(\frac{27}{2}\right) - 27(3) = 0$$

$$\text{or } 81 - 81 = 0 \text{ or } 0 = 0 \text{ which is true. } \therefore \lambda = 3 \text{ and } \mu = \frac{27}{2}.$$

6. Given that $\vec{a} \cdot \vec{b} = 0$ and $\vec{a} \times \vec{b} = \vec{0}$. What can you conclude about the vectors \vec{a} and \vec{b} ?

Sol. Given: $\vec{a} \cdot \vec{b} = 0 \Rightarrow |\vec{a}| |\vec{b}| \cos \theta = 0$


$$\Rightarrow \text{Either } |\vec{a}| = 0$$

$$\text{or } |\vec{b}| = 0 \text{ or } \cos \theta = 0 (\Rightarrow \theta = 90^\circ)$$

$$\Rightarrow \text{Either } \vec{a} = \vec{0} \text{ or } \vec{b} = \vec{0}$$

$$\text{or vector } \vec{a} \text{ is perpendicular to } \vec{b}. \quad \dots(i)$$

$$(\because \text{By definition, vector } \vec{a} \text{ is zero vector if and only if } |\vec{a}| = 0)$$




Again given $\vec{a} \times \vec{b} = \vec{0} \Rightarrow |\vec{a} \times \vec{b}| = 0$

$\Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 0$

[$\therefore |\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$]

\Rightarrow Either $|\vec{a}| = 0$ or $|\vec{b}| = 0$ or $\sin \theta = 0 (\Rightarrow \theta = 0)$



\Rightarrow Either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$ or vectors \vec{a} and \vec{b} are collinear (or parallel) vectors. ... (ii)

We know from common sense that vectors \vec{a} and \vec{b} are perpendicular to each other as well as are parallel (or collinear) is impossible. ... (iii)

\therefore From (i), (ii) and (iii), either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$

$\therefore \vec{a} \cdot \vec{b} = 0$ and $\vec{a} \times \vec{b} = \vec{0}$

\Rightarrow Either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$.

7. Let the vectors $\vec{a}, \vec{b}, \vec{c}$ be given as $a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}, b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}, c_1 \hat{i} + c_2 \hat{j} + c_3 \hat{k}$.

Then show that $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$.

Sol. Given: Vectors $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}, \vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k},$

$\vec{c} = c_1 \hat{i} + c_2 \hat{j} + c_3 \hat{k}$

$\therefore \vec{b} + \vec{c} = (b_1 + c_1) \hat{i} + (b_2 + c_2) \hat{j} + (b_3 + c_3) \hat{k}$

$$\begin{aligned} \text{L.H.S.} = \vec{a} \times (\vec{b} + \vec{c}) &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 + c_1 & b_2 + c_2 & b_3 + c_3 \end{vmatrix} \\ &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} + \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \end{aligned}$$

[By Property of Determinants]

$= \vec{a} \times \vec{b} + \vec{a} \times \vec{c} = \text{R.H.S.}$

8. If either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then $\vec{a} \times \vec{b} = \vec{0}$. Is the converse true? Justify your answer with an example.

Sol. Given: Either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$.

$$\therefore |\vec{a}| = |\vec{0}| = 0 \text{ or } |\vec{b}| = |\vec{0}| = 0 \quad \dots(i)$$

$$\therefore |\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta = 0 \quad (\sin \theta) = 0 \quad [\text{By (i)}]$$

$$\therefore \vec{a} \times \vec{b} = \vec{0} \quad (\text{By definition of zero vector})$$

But the converse is not true.

$$\text{Let } \vec{a} = \hat{i} + \hat{j} + \hat{k} \quad \therefore |\vec{a}| = \sqrt{1+1+1} = \sqrt{3} \neq 0.$$

$\therefore \vec{a}$ is a non-zero vector.

$$\text{Let } |\vec{b}| = 2(\hat{i} + \hat{j} + \hat{k}) = 2\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\therefore |\vec{b}| = \sqrt{4+4+4} \text{ or } |\vec{b}| = \sqrt{12} = \sqrt{4 \times 3} = 2\sqrt{3} \neq 0.$$

$\therefore \vec{b}$ is a non-zero vector.

$$\text{But } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{vmatrix}$$

$$\text{Taking 2 common from } R_3, = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix} = \vec{0}$$

($\because R_2$ and R_3 are identical)

9. Find the area of the triangle with vertices A(1, 1, 2), B(2, 3, 5) and C(1, 5, 5).

Sol. Vertices of $\triangle ABC$ are A(1, 1, 2), B(2, 3, 5) and C(1, 5, 5).


\therefore Position Vector (P.V.) of point A is $(1, 1, 2) = \hat{i} + \hat{j} + 2\hat{k}$

P.V. of point B is $(2, 3, 5)$

$$= 2\hat{i} + 3\hat{j} + 5\hat{k}$$

P.V. of point C is $(1, 5, 5)$

$$= \hat{i} + 5\hat{j} + 5\hat{k}$$



$\therefore \vec{AB} = \text{P.V. of point B} - \text{P.V. of point A}$

$$= 2\hat{i} + 3\hat{j} + 5\hat{k} - (\hat{i} + \hat{j} + 2\hat{k})$$

$$= 2\hat{i} + 3\hat{j} + 5\hat{k} - \hat{i} - \hat{j} - 2\hat{k} = \hat{i} + 2\hat{j} + 3\hat{k}$$

and $\vec{AC} = \text{P.V. of point C} - \text{P.V. of point A}$

$$= \hat{i} + 5\hat{j} + 5\hat{k} - (\hat{i} + \hat{j} + 2\hat{k}) = \hat{i} + 5\hat{j} + 5\hat{k} - \hat{i} - \hat{j} - 2\hat{k}$$

$$= 0\hat{i} + 4\hat{j} + 3\hat{k}$$

$$\therefore \vec{AB} \times \vec{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 0 & 4 & 3 \end{vmatrix}$$

$$= \hat{i}(6 - 12) - \hat{j}(3 - 0) + \hat{k}(4 - 0) = -6\hat{i} - 3\hat{j} + 4\hat{k}$$

We know that **area of triangle ABC**

$$= \frac{1}{2} |\vec{AB} \times \vec{AC}| = \frac{1}{2} \sqrt{36 + 9 + 16} = \frac{1}{2} \sqrt{61} \text{ sq. units.}$$

10. Find the area of the parallelogram whose adjacent sides are determined by the vectors $\vec{a} = \hat{i} - \hat{j} + 3\hat{k}$ and $\vec{b} = 2\hat{i} - 7\hat{j} + \hat{k}$.

Sol. Given: Vectors representing two adjacent sides of a parallelogram are

$$\vec{a} = \hat{i} - \hat{j} + 3\hat{k}$$

$$\text{and } \vec{b} = 2\hat{i} - 7\hat{j} + \hat{k}$$

$$\therefore \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 3 \\ 2 & -7 & 1 \end{vmatrix}$$

$$= \hat{i}(-1 + 21) - \hat{j}(1 - 6) + \hat{k}(-7 + 2) = 20\hat{i} + 5\hat{j} - 5\hat{k}$$

We know that **area of parallelogram** = $|\vec{a} \times \vec{b}|$

$$= \sqrt{400 + 25 + 25} = \sqrt{450} = \sqrt{25 \times 9 \times 2}$$


$$= 5(3) \sqrt{2} = 15\sqrt{2} \text{ square units.}$$

Note. Area of parallelogram whose **diagonal vectors** are $\vec{\alpha}$ and $\vec{\beta}$ is $\frac{1}{2} |\vec{\alpha} \times \vec{\beta}|$.

11. Let the vectors \vec{a} and \vec{b} be such that $|\vec{a}| = 3$, $|\vec{b}| = \frac{\sqrt{2}}{3}$, then $\vec{a} \times \vec{b}$ is a unit vector, if the angle between \vec{a} and \vec{b} is

(A) $\frac{\pi}{6}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$.

Sol. Given: $|\vec{a}| = 3$, $|\vec{b}| = \frac{\sqrt{2}}{3}$ and $\vec{a} \times \vec{b}$ is a unit vector.



$$\Rightarrow |\vec{a} \times \vec{b}| = 1 \quad \Rightarrow |\vec{a}| |\vec{b}| \sin \theta = 1$$

where θ is the angle between vectors \vec{a} and \vec{b} .

$$\text{Putting values of } |\vec{a}| \text{ and } |\vec{b}|, 3 \left(\frac{\sqrt{2}}{3} \right) \sin \theta = 1$$

$$\Rightarrow \sqrt{2} \sin \theta = 1 \Rightarrow \sin \theta = \frac{1}{\sqrt{2}} = \sin \frac{\pi}{4} \Rightarrow \theta = \frac{\pi}{4}$$

\therefore Option (B) is the correct answer.

12. Area of a rectangle having vertices A, B, C and D with position vectors $-\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k}$, $\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k}$, $\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}$ and $-\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}$, respectively, is

(A) $\frac{1}{2}$

(B) 1

(C) 2

(D) 4

Sol. Given: ABCD is a rectangle.

We know that \vec{AB} = P.V. of point B - P.V. of point A

$$= \hat{i} + \frac{1}{2}\hat{j} + 4\hat{k} - \left(-\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k} \right)$$

$$= \hat{i} + \frac{1}{2}\hat{j} + 4\hat{k} + \hat{i} - \frac{1}{2}\hat{j} - 4\hat{k} = 2\hat{i} + 0\hat{j} + 0\hat{k}$$

$$\therefore \vec{AB} = |\vec{AB}| = \sqrt{4+0+0} = \sqrt{4} = 2$$

and \vec{AD} = P.V. of point D - P.V. of point A

$$= -\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k} - \left(-\hat{i} + \frac{1}{2}\hat{j} + 4\hat{k} \right)$$

$$= -\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k} + \hat{i} - \frac{1}{2}\hat{j} - 4\hat{k} = -\hat{j} = 0\hat{i} - \hat{j} + 0\hat{k}$$


$$\therefore \vec{AD} = |\vec{AD}| = \sqrt{0+1+0} = \sqrt{1} = 1$$

$$\therefore \text{Area of rectangle ABCD} = (AB)(AD) \quad (= \text{Length} \times \text{Breadth})$$

$$= 2(1) = 2 \text{ sq. units}$$

\therefore Option (C) is the correct answer.

$$\text{or Area of rectangle ABCD} = \left| \vec{AB} \times \vec{AD} \right|.$$





MISCELLANEOUS EXERCISE

1. Write down a unit vector in XY-plane making an angle of 30° with the positive direction of x -axis.

Sol. Let \vec{OP} be the **unit** vector in XY-plane such that $\angle XOP = 30^\circ$

Therefore, $|\vec{OP}| = 1$

i.e., $OP = 1 \dots (i)$

By Triangle Law of Addition of vectors,

$$\begin{aligned} \text{In } \triangle OMP, \vec{OP} &= \vec{OM} + \vec{MP} \\ &= (OM)\hat{i} + (MP)\hat{j} \end{aligned}$$

$$[\because \hat{a} = \frac{\vec{a}}{|\vec{a}|} \Rightarrow \vec{a} = |\vec{a}| \hat{a} \text{ and unit vector along OX is } \hat{i}$$

and along OY is \hat{j}]

$$\Rightarrow \vec{OP} = OP \frac{OM}{OP} \hat{i} + OP \frac{MP}{OP} \hat{j}$$

(Dividing and multiplying by OP in R.H.S.)

$$= (1)(\cos 30^\circ) \hat{i} + (1)(\sin 30^\circ) \hat{j} \quad [\because \text{By (i), } OP = 1]$$

$$\Rightarrow \text{unit vector } \vec{OP} = (\cos 30^\circ) \hat{i} + (\sin 30^\circ) \hat{j} \dots (ii)$$

$$\Rightarrow \vec{OP} = \frac{\sqrt{3}}{2} \hat{i} + \frac{1}{2} \hat{j}.$$

Remark: From Eqn. (ii) of above solution, we can generalise the following result.

A unit vector along a line making an angle θ with positive

x-axis is $(\cos \theta) \hat{i} + (\sin \theta) \hat{j}$

2. Find the scalar components and magnitude of the vector joining the points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$.

Sol. Given points are $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$.



\Rightarrow P.V. (Position vector) of point P is

$$(x_1, y_1, z_1) = x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k}$$

and P.V. of point Q is $(x_2, y_2, z_2) = x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k}$

\therefore Vector \vec{PQ} , the vector joining the points P and Q.

= P.V. of terminal point Q - P.V. of initial point P

$$= x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k} - (x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k})$$

$$= x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k} - x_1 \hat{i} - y_1 \hat{j} - z_1 \hat{k}$$

$$\Rightarrow \vec{PQ} = (x_2 - x_1) \hat{i} + (y_2 - y_1) \hat{j} + (z_2 - z_1) \hat{k}$$

∴ Scalar components of the vector \vec{PQ} are the coefficients of \hat{i} , \hat{j} , \hat{k} in \vec{PQ} i.e., $(x_2 - x_1)$, $(y_2 - y_1)$, $(z_2 - z_1)$


and magnitude of vector \vec{PQ}

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2} \quad | \quad \sqrt{x^2 + y^2 + z^2}$$

3. A girl walks 4 km towards west, then she walks 3 km in a direction 30° east of north and stops. Determine the girl's displacement from her initial point of departure.

Sol. Let us take the initial point of departure as origin.

Let the girl walk a distance $OA = 4$ km towards west.



Through the point A draw a line AQ parallel to a line OP (which is 30° east of North i.e., in East-North quadrant making an angle of 30° with North)

Let the girl walk a distance $AB = 3$ km (given) along this direction \vec{OQ} (given). ∴ $\vec{OA} = 4(-\hat{i})$ [∵ Vector \vec{OA} is along OX']

$$= -4\hat{i} \quad \dots(i)$$

We know that (By Remark Q.N. 1 of this miscellaneous exercise) a unit vector along \vec{AQ} (or \vec{AB}) making an angle $\theta = 60^\circ$ with positive x-axis is $(\cos \theta)\hat{i} + (\sin \theta)\hat{j} = (\cos 60^\circ)\hat{i} + (\sin 60^\circ)\hat{j}$

$$= \frac{1}{2}\hat{i} + \frac{\sqrt{3}}{2}\hat{j}$$

$$\therefore \vec{AB} = |\vec{AB}| (\text{A unit vector along } \vec{AB}) \therefore \vec{a} = |\vec{a}| \hat{a}$$

$$= 3\left(\frac{1}{2}\hat{i} + \frac{\sqrt{3}}{2}\hat{j}\right) = \frac{3}{2}\hat{i} + \frac{3\sqrt{3}}{2}\hat{j} \quad \dots (ii)$$

\therefore Girl's displacement from her initial point O of departure (to final point B) = $\vec{OB} = \vec{OA} + \vec{AB}$ (By Triangle Law of Addition of vectors)

$$= -4\hat{i} + \left(\frac{3}{2}\hat{i} + \frac{3\sqrt{3}}{2}\hat{j}\right) = \left(-4 + \frac{3}{2}\right)\hat{i} + \frac{3\sqrt{3}}{2}\hat{j}$$

[By (i)] [By (ii)]

$$= \frac{-5}{2}\hat{i} + \frac{3\sqrt{3}}{2}\hat{j}.$$


4. If $\vec{a} = \vec{b} + \vec{c}$, then is it true that $|\vec{a}| = |\vec{b}| + |\vec{c}|$? Justify your answer.

Sol. The result is not true (always).

Given: $\vec{a} = \vec{b} + \vec{c}$.

\therefore Either the vectors \vec{a} , \vec{b} , \vec{c} are collinear or vectors \vec{a} , \vec{b} , \vec{c} form the sides of a triangle.

Case I. Vectors \vec{a} , \vec{b} , \vec{c} are collinear.



Let $\vec{a} = \vec{AC}$, $\vec{b} = \vec{AB}$ and $\vec{c} = \vec{BC}$,

then $\vec{a} = \vec{AC} = \vec{AB} + \vec{BC} = \vec{b} + \vec{c}$.

Also, $|\vec{a}| = AC = AB + BC = |\vec{b}| + |\vec{c}|$.

Case II. Vectors \vec{a} , \vec{b} , \vec{c} form a triangle.


Here also by Triangle Law of vectors,

$$\vec{a} = \vec{b} + \vec{c}$$

But $|\vec{a}| < |\vec{b}| + |\vec{c}|$

(\because Each side of a triangle is less than sum of the other two sides)

$\therefore |\vec{a}| = |\vec{b} + \vec{c}| = |\vec{b}| + |\vec{c}|$ is true only when vectors \vec{b} and \vec{c} are collinear vectors.



5. Find the value of x for which $x(\hat{i} + \hat{j} + \hat{k})$ is a unit vector.

Sol. Because $x(\hat{i} + \hat{j} + \hat{k}) = x\hat{i} + x\hat{j} + x\hat{k}$ is a unit vector (given)

Therefore, $|x\hat{i} + x\hat{j} + x\hat{k}| = 1$

$$\therefore \sqrt{x^2 + x^2 + x^2} = 1$$

$$[\because x\hat{i} + y\hat{j} + z\hat{k} = \sqrt{x^2 + y^2 + z^2}]$$

$$\text{Squaring both sides } 3x^2 = 1 \text{ or } x^2 = \frac{1}{3} \quad \therefore x = \pm \frac{1}{\sqrt{3}}.$$

- 6. Find a vector of magnitude 5 units and parallel to the resultant of the vectors**

$$\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k} \text{ and } \vec{b} = \hat{i} - 2\hat{j} + \hat{k}.$$

Sol. Given: Vectors $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$.

Let vector \vec{c} be the resultant of vectors \vec{a} and \vec{b} .

$$\begin{aligned} \therefore \vec{c} &= \vec{a} + \vec{b} = 2\hat{i} + 3\hat{j} - \hat{k} + \hat{i} - 2\hat{j} + \hat{k} \\ &= 3\hat{i} + \hat{j} + 0\hat{k}. \end{aligned}$$

\therefore Required vector of magnitude 5 units and parallel (or collinear) to resultant vector $\vec{c} = \vec{a} + \vec{b}$ is

$$\begin{aligned} 5\hat{c} &= 5 \frac{\vec{c}}{|\vec{c}|} = 5 \left(\frac{3\hat{i} + \hat{j} + 0\hat{k}}{\sqrt{9+1+0}} \right) \\ &= \frac{5}{\sqrt{10}} (3\hat{i} + \hat{j}) = \frac{5}{\sqrt{10}} \cdot \frac{\sqrt{10}}{\sqrt{10}} (3\hat{i} + \hat{j}) \\ &= \frac{5}{10} \sqrt{10} (3\hat{i} + \hat{j}) = \frac{\sqrt{10}}{2} (3\hat{i} + \hat{j}) = \frac{3}{2} \sqrt{10} \hat{i} + \frac{\sqrt{10}}{2} \hat{j}. \end{aligned}$$

- 7. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - \hat{j} + 3\hat{k}$ and**

$\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$, find a unit vector parallel to the vector $2\vec{a} - \vec{b} + 3\vec{c}$.

Sol. Given: Vectors $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - \hat{j} + 3\hat{k}$

and $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$.

Let $\vec{d} = 2\vec{a} - \vec{b} + 3\vec{c}$

$$\begin{aligned} &= 2(\hat{i} + \hat{j} + \hat{k}) - (2\hat{i} - \hat{j} + 3\hat{k}) + 3(\hat{i} - 2\hat{j} + \hat{k}) \\ &= 2\hat{i} + 2\hat{j} + 2\hat{k} - 2\hat{i} + \hat{j} - 3\hat{k} + 3\hat{i} - 6\hat{j} + 3\hat{k} \end{aligned}$$

$\therefore \vec{d} = 3\hat{i} - 3\hat{j} + 2\hat{k} \therefore$ A unit vector parallel to the vector

$$\vec{d} = 3\hat{i} - 3\hat{j} + 2\hat{k} \text{ is}$$

$$\hat{d} = \frac{\vec{d}}{|\vec{d}|} = \frac{3\hat{i} - 3\hat{j} + 2\hat{k}}{\sqrt{9+9+4}} = \frac{3}{\sqrt{22}} \hat{i} - \frac{3}{\sqrt{22}} \hat{j} + \frac{2}{\sqrt{22}} \hat{k}.$$

8. Show that the points A(1, -2, -8), B(5, 0, -2) and C(11, 3, 7) are collinear and find the ratio in which B divides AC.

Sol. Given: Points A(1, -2, -8), B(5, 0, -2) and C(11, 3, 7).

i.e., Position vectors of points A, B, C are

$$\vec{OA} (= A(1, -2, -8)) = \hat{i} - 2\hat{j} - 8\hat{k}$$

$$\vec{OB} (= B(5, 0, -2)) = 5\hat{i} + 0\hat{j} - 2\hat{k} = 5\hat{i} - 2\hat{k}$$

$$\text{and } \vec{OC} (= C(11, 3, 7)) = 11\hat{i} + 3\hat{j} + 7\hat{k}$$

$$\therefore \vec{AB} = \text{P.V. of point B} - \text{P.V. of point A}$$

$$= 5\hat{i} - 2\hat{k} - (\hat{i} - 2\hat{j} - 8\hat{k}) = 5\hat{i} - 2\hat{k} - \hat{i} + 2\hat{j} + 8\hat{k}$$

$$\text{or } \vec{AB} = 4\hat{i} + 2\hat{j} + 6\hat{k}$$

$$\therefore AB = |\vec{AB}| = \sqrt{16 + 4 + 36} = \sqrt{56} = \sqrt{4 \times 14} = 2\sqrt{14}$$

$$\text{and } \vec{BC} = \text{P.V. of point C} - \text{P.V. of point B}$$

$$= 11\hat{i} + 3\hat{j} + 7\hat{k} - (5\hat{i} - 2\hat{k}) = 11\hat{i} + 3\hat{j} + 7\hat{k} - 5\hat{i} + 2\hat{k}$$

$$= 6\hat{i} + 3\hat{j} + 9\hat{k}$$

$$\therefore BC = |\vec{BC}| = \sqrt{36 + 9 + 81} = \sqrt{126} = \sqrt{9 \times 14} = 3\sqrt{14}$$

$$\vec{AC} = \text{P.V. of point C} - \text{P.V. of point A}$$

$$= 11\hat{i} + 3\hat{j} + 7\hat{k} - (\hat{i} - 2\hat{j} - 8\hat{k})$$

$$= 11\hat{i} + 3\hat{j} + 7\hat{k} - \hat{i} + 2\hat{j} + 8\hat{k} = 10\hat{i} + 5\hat{j} + 15\hat{k}$$

$$\therefore AC = |\vec{AC}| = \sqrt{100 + 25 + 225} = \sqrt{350} = \sqrt{25 \times 14} = 5\sqrt{14}$$

$$\text{Now, } \vec{AB} + \vec{BC} = 4\hat{i} + 2\hat{j} + 6\hat{k} + 6\hat{i} + 3\hat{j} + 9\hat{k}$$

$$= 10\hat{i} + 5\hat{j} + 15\hat{k} = \vec{AC}$$

\therefore Points A, B, C are either collinear or are the vertices of $\triangle ABC$.

$$\text{Again } AB + BC = 2\sqrt{14} + 3\sqrt{14} = (2 + 3)\sqrt{14} = 5\sqrt{14} = AC$$

\therefore Points A, B, C are collinear.

Now to find the ratio in which B divides AC

$$\begin{array}{c} \lambda : 1 \\ \bullet \text{-----} \bullet \text{-----} \bullet \\ A(1, -2, -8) \quad B(5, 0, -2) \quad C(11, 3, 7) \\ = \vec{a} \quad \quad \quad = \vec{b} \quad \quad \quad = \vec{c} \end{array}$$

Let the point B divides AC in the ratio $\lambda : 1$.

∴ By section formula, P.V. of point B is $\frac{\lambda \vec{c} + 1 \vec{a}}{\lambda + 1}$

$$\Rightarrow (5, 0, -2) = \frac{\lambda(11, 3, 7) + (1, -2, -8)}{\lambda + 1}$$

Cross-multiplying,

$$\begin{aligned} (\lambda + 1)(5\hat{i} + 0\hat{j} - 2\hat{k}) &= \lambda(11\hat{i} + 3\hat{j} + 7\hat{k}) + (\hat{i} - 2\hat{j} - 8\hat{k}) \\ \Rightarrow 5(\lambda + 1)\hat{i} - 2(\lambda + 1)\hat{k} &= 11\lambda\hat{i} + 3\lambda\hat{j} + 7\lambda\hat{k} + \hat{i} - 2\hat{j} - 8\hat{k} \\ \Rightarrow (5\lambda + 5)\hat{i} - (2\lambda + 2)\hat{k} &= (11\lambda + 1)\hat{i} + (3\lambda - 2)\hat{j} + (7\lambda - 8)\hat{k} \end{aligned}$$

Comparing coefficients of \hat{i} , \hat{j} , \hat{k} on both sides, we have

$$\begin{aligned} 5\lambda + 5 &= 11\lambda + 1, 0 = 3\lambda - 2, -(2\lambda + 2) = 7\lambda - 8 \\ \Rightarrow -6\lambda &= -4, -3\lambda = -2, -2\lambda - 2 = 7\lambda - 8 \quad (\Rightarrow -9\lambda = -6) \end{aligned}$$

$$\Rightarrow \lambda = \frac{4}{6} = \frac{2}{3}, \lambda = \frac{2}{3}, \lambda = \frac{6}{9} = \frac{2}{3}$$

All three values of λ are same.

$$\therefore \text{Required ratio is } \lambda : 1 = \frac{2}{3} : 1 = 2 : 3.$$

9. Find the position vector of a point R which divides the line joining the two points P and Q whose position vectors are

$(2\vec{a} + \vec{b})$ and $(\vec{a} - 3\vec{b})$ externally in the ratio 1 : 2. Also, show that P is the middle point of line segment RQ.

Sol. We know that position vector of the point R dividing the join of P and Q externally in the ratio 1 : 2 = $m : n$ is given by

$$\begin{aligned} \vec{c} &= \frac{m\vec{b} - n\vec{a}}{m - n} = \frac{1(\vec{a} - 3\vec{b}) - 2(2\vec{a} + \vec{b})}{1 - 2} \\ &= \frac{\vec{a} - 3\vec{b} - 4\vec{a} - 2\vec{b}}{1 - 2} = \frac{-3\vec{a} - 5\vec{b}}{-1} = 3\vec{a} + 5\vec{b} \end{aligned}$$

Again position vector of the middle point of the line segment RQ

$$= \frac{\text{P.V. of point R} + \text{P.V. of point Q}}{2} = \frac{3\vec{a} + 5\vec{b} + \vec{a} - 3\vec{b}}{2} = \frac{4\vec{a} + 2\vec{b}}{2}$$

$$= 2\vec{a} + \vec{b} = \text{P.V. of point P. (given)}$$

∴ Point P is the middle point of the line segment RQ.

10. Two adjacent sides of a parallelogram are $2\hat{i} - 4\hat{j} + 5\hat{k}$

and $\hat{i} - 2\hat{j} - 3\hat{k}$. Find the unit vector parallel to its diagonal. Also, find its area.

Sol. Let ABCD be a parallelogram.

Given: The vectors representing two adjacent sides of this parallelogram are say

$$\vec{a} = 2\hat{i} - 4\hat{j} + 5\hat{k}$$

and $\vec{b} = \hat{i} - 2\hat{j} - 3\hat{k}$

Formula: \therefore Vectors along the diagonals \vec{AC} and \vec{DB} of the parallelogram are

$$\vec{a} + \vec{b} \text{ and } \vec{a} - \vec{b}$$

$$\begin{aligned} \text{i.e., } \vec{a} + \vec{b} &= 2\hat{i} - 4\hat{j} + 5\hat{k} + \hat{i} - 2\hat{j} - 3\hat{k} \\ &= 3\hat{i} - 6\hat{j} + 2\hat{k} \end{aligned}$$

$$\begin{aligned} \text{and } \vec{a} - \vec{b} &= 2\hat{i} - 4\hat{j} + 5\hat{k} - (\hat{i} - 2\hat{j} - 3\hat{k}) \\ &= 2\hat{i} - 4\hat{j} + 5\hat{k} - \hat{i} + 2\hat{j} + 3\hat{k} = \hat{i} - 2\hat{j} + 8\hat{k} \end{aligned}$$

\therefore Unit vectors parallel to (or along) diagonals are

$$\frac{\vec{a} + \vec{b}}{|\vec{a} + \vec{b}|} \text{ and } \frac{\vec{a} - \vec{b}}{|\vec{a} - \vec{b}|} = \frac{3\hat{i} - 6\hat{j} + 2\hat{k}}{\sqrt{9 + 36 + 4} = \sqrt{49} = 7} \text{ and } \frac{\hat{i} - 2\hat{j} + 8\hat{k}}{\sqrt{1 + 4 + 64} = \sqrt{69}}$$

Let us find area of parallelogram

$$\begin{aligned} \vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -4 & 5 \\ 1 & -2 & -3 \end{vmatrix} = \hat{i}(12 + 10) - \hat{j}(-6 - 5) + \hat{k}(-4 + 4) \\ &= 22\hat{i} + 11\hat{j} + 0\hat{k} \end{aligned}$$

We know that area of parallelogram = $|\vec{a} \times \vec{b}|$

$$= \sqrt{(22)^2 + (11)^2 + 0^2} = \sqrt{484 + 121} = \sqrt{605}$$

$$= \sqrt{5 \times 121} = \sqrt{121 \times 5} = 11\sqrt{5} \text{ sq. units.}$$

11. Show that the direction cosines of a vector equally inclined to the axes OX, OY and OZ are $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$.


Sol. Let l, m, n be the direction cosines of a vector equally inclined to the axes OX, OY, OZ.

\therefore A unit vector along the given vector is

$$\hat{a} = l\hat{i} + m\hat{j} + n\hat{k} \quad \text{and} \quad |\hat{a}| = 1$$

$$\Rightarrow \sqrt{l^2 + m^2 + n^2} = 1 \quad \therefore l^2 + m^2 + n^2 = 1 \quad \dots(i)$$

Let the given vector (for which unit vector is \hat{a}) make equal angles (given) θ, θ, θ (say) with $\mathbf{OX} (\Rightarrow \hat{i})$, $\mathbf{OY} (\Rightarrow \hat{j})$ and $\mathbf{OZ} (\Rightarrow \hat{k})$
 \therefore The given vector is in positive octant OXYZ and hence θ is acute.
...(ii)



∴ For angle θ between \hat{a} and \hat{i} ,

$$\cos \theta = \frac{\hat{a} \cdot \hat{i}}{|\hat{a}| |\hat{i}|} = \frac{(l\hat{i} + m\hat{j} + n\hat{k}) \cdot (\hat{i} + 0\hat{j} + 0\hat{k})}{(1)(1)}$$

$$\text{or } \cos \theta = l(1) + m(0) + n(0) = l$$

$$\text{or } l = \cos \theta \quad \dots(iii)$$

Similarly, for angle θ between \hat{a} and \hat{j} , $m = \cos \theta \quad \dots(iv)$

Similarly, for angle θ between \hat{a} and \hat{k} , $n = \cos \theta \quad \dots(v)$

Putting these values of l , m , n from (iii), (iv) and (v) in (i), we have

$$\cos^2 \theta + \cos^2 \theta + \cos^2 \theta = 1 \Rightarrow 3 \cos^2 \theta = 1$$

$$\Rightarrow \cos^2 \theta = \frac{1}{3} \Rightarrow \cos \theta = \pm \sqrt{\frac{1}{3}} = \pm \frac{1}{\sqrt{3}}$$

∴ $\cos \theta = \frac{1}{\sqrt{3}}$ (∵ By (ii), θ is acute and hence $\cos \theta$ is positive)

Putting $\cos \theta = \frac{1}{\sqrt{3}}$ in (ii), (iii) and (iv), direction cosines of the

required vector are l , m , $n = \frac{1}{\sqrt{3}}$, $\frac{1}{\sqrt{3}}$ and $\frac{1}{\sqrt{3}}$.

12. Let $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$, $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$. Find a vector \vec{b} which is perpendicular to both \vec{a} and \vec{b} , and $\vec{c} \cdot \vec{d} = 15$.

Sol. Given: Vectors are $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$

and $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$

By definition of cross-product of two

vectors, $\vec{a} \times \vec{b}$ is a vector

perpendicular to both \vec{a} and \vec{b} .


Hence, vector \vec{d} which is also perpendicular to both \vec{a} and \vec{b}

is $\vec{d} = \lambda(\vec{a} \times \vec{b})$ where $\lambda = 1$ or some other scalar.

$$\text{Therefore, } \vec{d} = \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 4 & 2 \\ 3 & -2 & 7 \end{vmatrix}$$

Expanding along first row, $= \lambda[\hat{i}(28 + 4) - \hat{j}(7 - 6) + \hat{k}(-2 - 12)]$

$$\text{or } \vec{d} = \lambda[32\hat{i} - \hat{j} - 14\hat{k}] \quad \dots(i)$$



$$\text{or } \vec{d} = 32\lambda \hat{i} - \lambda \hat{j} - 14\lambda \hat{k}$$

$$\text{To find } \lambda: \text{ Given: } \vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$$

$$\text{Also given } \vec{c} \cdot \vec{d} = 15$$

$$\Rightarrow 2(32\lambda) + (-1)(-\lambda) + 4(-14\lambda) = 15$$

$$\Rightarrow 64\lambda + \lambda - 56\lambda = 15 \Rightarrow 9\lambda = 15 \Rightarrow \lambda = \frac{5}{3}$$

Putting $\lambda = \frac{5}{3}$ in (i), required vector

$$\vec{d} = \frac{5}{3} (32\hat{i} - \hat{j} - 14\hat{k}) = \frac{1}{3} (160\hat{i} - 5\hat{j} - 70\hat{k}).$$

13. The scalar product of the vector $\hat{i} + \hat{j} + \hat{k}$ with a unit vector along the sum of the vectors $2\hat{i} + 4\hat{j} - 5\hat{k}$ and $\lambda\hat{i} + 2\hat{j} + 3\hat{k}$ is equal to one. Find the value of λ .

Sol. Given: Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$... (i)

$$\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k} \quad \text{and} \quad \vec{c} = \lambda\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\therefore \vec{b} + \vec{c} (= \vec{d} \text{ (say)}) = (2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}$$

$$\therefore \hat{d}, \text{ a unit vector along } \vec{b} + \vec{c} = \vec{d} \text{ is}$$

$$\hat{d} = \frac{\vec{d}}{|\vec{d}|} = \frac{(2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{(2 + \lambda)^2 + 36 + 4}} = \frac{(2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{4 + \lambda^2 + 4\lambda + 40}}$$

$$\begin{aligned} \text{or } \hat{d} &= \frac{(2 + \lambda)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{\lambda^2 + 4\lambda + 44}} \\ &= \frac{(2 + \lambda)}{\sqrt{\lambda^2 + 4\lambda + 44}} \hat{i} + \frac{6}{\sqrt{\lambda^2 + 4\lambda + 44}} \hat{j} - \frac{2}{\sqrt{\lambda^2 + 4\lambda + 44}} \hat{k} \end{aligned} \quad \dots (ii)$$

Given: Scalar (i.e., Dot) Product of \vec{a} and \hat{d} i.e., $\vec{a} \cdot \hat{d} = 1$

\therefore From (i) and (ii),

$$\frac{1(2 + \lambda)}{\sqrt{\lambda^2 + 4\lambda + 44}} + \frac{1(6)}{\sqrt{\lambda^2 + 4\lambda + 44}} + \frac{1(-2)}{\sqrt{\lambda^2 + 4\lambda + 44}} = 1$$

Multiplying by L.C.M. = $\sqrt{\lambda^2 + 4\lambda + 44}$,

$$2 + \lambda + 6 - 2 = \sqrt{\lambda^2 + 4\lambda + 44} \Rightarrow \lambda + 6 = \sqrt{\lambda^2 + 4\lambda + 44}$$

$$\text{Squaring both sides } (\lambda + 6)^2 = \lambda^2 + 4\lambda + 44$$

$$\Rightarrow \lambda^2 + 12\lambda + 36 = \lambda^2 + 4\lambda + 44$$

$$\Rightarrow 8\lambda = 8 \Rightarrow \lambda = 1.$$

14. If \vec{a} , \vec{b} , \vec{c} are mutually perpendicular vectors of equal magnitude, show that the vector $\vec{a} + \vec{b} + \vec{c}$ is equally inclined to \vec{a} , \vec{b} , \vec{c} .

Sol. Given: \vec{a} , \vec{b} , \vec{c} are mutually perpendicular vectors of equal magnitude.

$$\therefore \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a} = 0, \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{b} = 0,$$

$$\vec{a} \cdot \vec{a} = \vec{a} \cdot \vec{c} = 0 \quad \dots (i)$$

$$\text{and } |\vec{a}| = |\vec{b}| = |\vec{c}| = \lambda \text{ (say)} \quad \dots (ii)$$

Let vector $\vec{d} = \vec{a} + \vec{b} + \vec{c}$ make angles $\theta_1, \theta_2, \theta_3$ with vectors $\vec{a}, \vec{b}, \vec{c}$ respectively.

$$\therefore \cos \theta_1 = \frac{\vec{d} \cdot \vec{a}}{|\vec{d}| |\vec{a}|} = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \quad \dots (iii)$$

$$= \frac{\vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} = \frac{|\vec{a}|^2 + 0 + 0}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} \quad [\text{By (i)}]$$

$$\Rightarrow \cos \theta_1 = \frac{|\vec{a}|^2}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|} = \frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|} \quad \dots (iii)$$

Let us now find $|\vec{a} + \vec{b} + \vec{c}|$.

$$\begin{aligned} \text{We know that } |\vec{a} + \vec{b} + \vec{c}|^2 &= (\vec{a} + \vec{b} + \vec{c})^2 \\ &= \vec{a}^2 + (\vec{b} + \vec{c})^2 + 2\vec{a} \cdot (\vec{b} + \vec{c}) \\ &= \vec{a}^2 + \vec{b}^2 + \vec{c}^2 + 2\vec{a} \cdot \vec{b} + 2\vec{a} \cdot \vec{c} \quad [\because (\vec{A} + \vec{B})^2 = \vec{A}^2 + \vec{B}^2 + 2\vec{A} \cdot \vec{B}] \\ &= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2\vec{a} \cdot \vec{b} + 2\vec{a} \cdot \vec{c} \\ &= \lambda^2 + \lambda^2 + \lambda^2 + 0 + 0 + 0 = 3\lambda^2 \end{aligned}$$

Putting values from (i) and (ii)

$$|\vec{a} + \vec{b} + \vec{c}|^2 = \lambda^2 + \lambda^2 + \lambda^2 + 0 + 0 + 0 = 3\lambda^2$$

$$\therefore |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3\lambda^2} = \lambda\sqrt{3}$$

$$\text{Putting this value of } |\vec{a} + \vec{b} + \vec{c}| = \lambda\sqrt{3} \text{ and } |\vec{a}| = \lambda \text{ from (ii) in (iii), } \cos \theta_1 = \frac{\lambda}{\lambda\sqrt{3}} = \frac{1}{\sqrt{3}} \quad \therefore \theta_1 = \cos^{-1} \frac{1}{\sqrt{3}}$$

$$\text{Similarly, } \theta_2 = \cos^{-1} \frac{1}{\sqrt{3}} \text{ and } \theta_3 = \cos^{-1} \frac{1}{\sqrt{3}}$$

$$\therefore \theta_1 = \theta_2 = \theta_3 \left(= \cos^{-1} \frac{1}{\sqrt{3}} \right)$$

\therefore Vector $\vec{a} + \vec{b} + \vec{c}$ is equally inclined to the vectors \vec{a}, \vec{b} and \vec{c} .

15. Prove that $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a}|^2 + |\vec{b}|^2$, if and only if \vec{a}, \vec{b} are perpendicular, given $\vec{a} \neq \vec{0}, \vec{b} \neq \vec{0}$.

Sol. We know that $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b})$

$$= \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b}$$

$$= |\vec{a}|^2 + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{b} + |\vec{b}|^2$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} \quad \dots(i)$$

For If part: Given: \vec{a} and \vec{b} are perpendicular

$$\Rightarrow \vec{a} \cdot \vec{b} = 0$$

Putting $\vec{a} \cdot \vec{b} = 0$ in (i), we have

$$(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a}|^2 + |\vec{b}|^2$$

For Only if part:

Given: $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a}|^2 + |\vec{b}|^2$

Putting this value in L.H.S. eqn. (i), we have

$$|\vec{a}|^2 + |\vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 0 = 2\vec{a} \cdot \vec{b} \quad \Rightarrow \vec{a} \cdot \vec{b} = \frac{0}{2} = 0$$

But $\vec{a} \neq \vec{0}$ and $\vec{b} \neq \vec{0}$ (given).

\therefore Vector \vec{a} and \vec{b} are perpendicular to each other.

16. **Choose the correct answer:**

If θ is the angle between two vectors \vec{a} and \vec{b} , then $\vec{a} \cdot \vec{b} \geq 0$ only when

(A) $0 < \theta < \frac{\pi}{2}$ (B) $0 \leq \theta \leq \frac{\pi}{2}$ (C) $0 < \theta < \pi$ (D) $0 < \theta \leq \pi$

Sol. Given: $\vec{a} \cdot \vec{b} \geq 0$

$$\Rightarrow |\vec{a}| |\vec{b}| \cos \theta \geq 0 \quad \Rightarrow \cos \theta \geq 0$$

$[\because |\vec{a}|$ and $|\vec{b}|$ being lengths of vectors are always ≥ 0]
and this is true only for option (B) out of the given

options $\left(\because \text{For option (A)} \quad 0 < \theta < \frac{\pi}{2}, \cos \theta > 0 \right)$.

17. **Choose the correct answer:**

Let \vec{a} and \vec{b} be two unit vectors and θ is the angle between them. Then $\vec{a} + \vec{b}$ is a unit vector if

(A) $\theta = \frac{\pi}{4}$ (B) $\theta = \frac{\pi}{3}$ (C) $\theta = \frac{\pi}{2}$ (D) $\theta = \frac{2\pi}{3}$.

Sol. Given: \vec{a}, \vec{b} and $\vec{a} + \vec{b}$ are unit vectors

$$\Rightarrow |\vec{a}| = 1, |\vec{b}| = 1 \text{ and } |\vec{a} + \vec{b}| = 1$$

Now, squaring both sides of $|\vec{a} + \vec{b}| = 1$, we have

$$|\vec{a} + \vec{b}|^2 = 1 \Rightarrow (\vec{a} + \vec{b})^2 = 1$$

$$\Rightarrow \vec{a}^2 + \vec{b}^2 + 2\vec{a} \cdot \vec{b} = 1$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2|\vec{a}||\vec{b}|\cos\theta = 1$$

where θ is the given angle between vectors \vec{a} and \vec{b} .

Putting $|\vec{a}| = 1$ and $|\vec{b}| = 1$, we have $1 + 1 + 2\cos\theta = 1$

$$\Rightarrow 2\cos\theta = -1 \Rightarrow \cos\theta = \frac{-1}{2} = -\cos 60^\circ$$

$$\Rightarrow \cos\theta = \cos(180^\circ - 60^\circ) \Rightarrow \cos\theta = \cos 120^\circ$$

$$\Rightarrow \theta = 120^\circ = 120 \times \frac{\pi}{180} = \frac{2\pi}{3}$$

\therefore Option (D) is the correct answer.


Very Important Results

$$(1) \hat{i} \cdot \hat{i} = |\hat{i}|^2 = 1, \hat{j} \cdot \hat{j} = 1, \hat{k} \cdot \hat{k} = 1.$$

$$(2) \hat{i} \times \hat{i} = \vec{0}, \hat{j} \times \hat{j} = \vec{0} \text{ and } \hat{k} \times \hat{k} = \vec{0}.$$

$$(3) \hat{i} \cdot \hat{j} = 0 = \hat{j} \cdot \hat{i}, \hat{j} \cdot \hat{k} = 0 = \hat{k} \cdot \hat{j}, \hat{i} \cdot \hat{k} = 0 = \hat{k} \cdot \hat{i}.$$

$$(4) \hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i} \text{ and } \hat{k} \times \hat{i} = \hat{j}.$$



18. Choose the correct answer:

The value of $\hat{i} \cdot (\hat{j} \times \hat{k}) + \hat{j} \cdot (\hat{i} \times \hat{k}) + \hat{k} \cdot (\hat{i} \times \hat{j})$ is
 (A) 0 (B) -1 (C) 1 (D) 3

Sol. $\hat{i} \cdot (\hat{j} \times \hat{k}) + \hat{j} \cdot (\hat{i} \times \hat{k}) + \hat{k} \cdot (\hat{i} \times \hat{j})$

$$= \hat{i} \cdot \hat{i} + \hat{j} \cdot (-\hat{j}) + \hat{k} \cdot \hat{k}$$

$$(\because \hat{i} \times \hat{k} = -\hat{k} \times \hat{i} = -\hat{j})$$

$$= 1 - 1 + 1 = 1$$

\therefore Option (C) is the correct answer.

19. If θ be the angle between any two vectors \vec{a} and \vec{b} , then

$|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$, when θ is equal to

(A) 0 (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{2}$ (D) π

Sol. Given: $|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$

$$\Rightarrow |\vec{a}| |\vec{b}| |\cos\theta| = |\vec{a}| |\vec{b}| |\sin\theta|$$

$$(\because \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos\theta)$$

$$\Rightarrow |\vec{a} \cdot \vec{b}| = |\vec{a}| |\vec{b}| |\cos\theta|$$

Dividing both sides by $|\vec{a}| |\vec{b}|$, we have $|\cos\theta| = |\sin\theta|$

and this equation is true only for option (B) namely $\theta = \frac{\pi}{4}$ out of the given options.

$$\left[\because \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}} \text{ and also } \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}} \right]$$

\therefore Option (B) is the correct option.