



## Exercise 7.1

**Find an antiderivative (or integral) of the following functions by the method of inspection in Exercises 1 to 5.**

1.  $\sin 2x$

**Sol.** To find an anti derivative of  $\sin 2x$  by Inspection Method.

We know that  $\frac{d}{dx} (\cos 2x) = -2 \sin 2x$

Dividing by  $-2$ ,  $\frac{-1}{2} \frac{d}{dx} (\cos 2x) = \sin 2x$

$$\text{or } \frac{d}{dx} \left( \frac{-1}{2} \cos 2x \right) = \sin 2x$$

$\therefore$  By definition; **an** integral or **an** antiderivative of  $\sin 2x$  is  $\frac{-1}{2} \cos 2x$ .

**Note.** In fact anti derivative or integral of  $\sin 2x$  is  $\frac{-1}{2} \cos 2x + c$ .  
For different values of  $c$ , we get different antiderivatives. So we omitted  $c$  for writing **an** anti derivative.

## 2. $\cos 3x$

**Sol.** To find an anti derivative of  $\cos 3x$  by Inspection Method.

We know that  $\frac{d}{dx} (\sin 3x) = 3 \cos 3x$

Dividing by  $3$ ,  $\frac{1}{3} \frac{d}{dx} (\sin 3x) = \cos 3x$  or  $\frac{d}{dx} \left( \frac{1}{3} \sin 3x \right) = \cos 3x$

$\therefore$  By definition, **an** integral or **an** antiderivative of  $\cos 3x$  is  $\frac{1}{3} \sin 3x$ .

(See note after solution of Q.No.1 for not adding  $c$  to the answer.)

## 3. $e^{2x}$ .

**Sol.** To find an antiderivative of  $e^{2x}$  by Inspection Method.

We know that  $\frac{d}{dx} e^{2x} = e^{2x} \frac{d}{dx} (2x) = 2e^{2x}$

Dividing by  $2$ ,  $\frac{1}{2} \frac{d}{dx} e^{2x} = e^{2x}$  or  $\frac{d}{dx} \left( \frac{1}{2} e^{2x} \right) = e^{2x}$

$\therefore$  An antiderivative of  $e^{2x}$  is  $\frac{1}{2} e^{2x}$ .

## 4. $(ax + b)^2$ .

**Sol.** To find an anti derivative of  $(ax + b)^2$ .

We know that  $\frac{d}{dx} (ax + b)^3 = 3(ax + b)^2 \frac{d}{dx} (ax + b) = 3(ax + b)^2 a$ .

Dividing by  $3a$ ,  $\frac{1}{3a} \frac{d}{dx} (ax + b)^3 = (ax + b)^2$

or  $\frac{d}{dx} \left[ \frac{1}{3a} (ax + b)^3 \right] = (ax + b)^2$

$\therefore$  An anti derivative of  $(ax + b)^2$  is  $\frac{1}{3a} (ax + b)^3$ .

## 5. $\sin 2x - 4e^{3x}$ .

**Sol.** To find an anti derivative of  $\sin 2x - 4e^{3x}$  by Inspection Method.

We know that  $\frac{d}{dx} (\cos 2x) = -2 \sin 2x$

$$\text{Dividing by } -2, \quad \frac{d}{dx} \left( \frac{-1}{2} \cos 2x \right) = \sin 2x \quad \dots(i)$$

$$\text{Again } \frac{d}{dx} e^{3x} = 3e^{3x} \quad \therefore \quad \frac{d}{dx} \left( \frac{1}{3} e^{3x} \right) = e^{3x}$$

$$\text{Multiplying by } -4, \quad \frac{d}{dx} \left( \frac{-4}{3} e^{3x} \right) = -4e^{3x} \quad \dots(ii)$$

Adding eqns. (i) and (ii)

$$\frac{d}{dx} \left( \frac{-1}{2} \cos 2x \right) + \frac{d}{dx} \left( \frac{-4}{3} e^{3x} \right) = \sin 2x - 4e^{3x}$$

$$\text{or} \quad \frac{d}{dx} \left( \frac{-1}{2} \cos 2x - \frac{4}{3} e^{3x} \right) = \sin 2x - 4e^{3x}$$

$$\therefore \text{ An anti derivative of } \sin 2x - 4e^{3x} \text{ is } \frac{-1}{2} \cos 2x - \frac{4}{3} e^{3x}.$$

**Evaluate the following integrals in Exercises 6 to 11.**

6.  $\int (4e^{3x} + 1) dx.$

$$\begin{aligned} \text{Sol. } \int (4e^{3x} + 1) dx &= \int 4e^{3x} dx + \int 1 dx \\ &= 4 \int e^{3x} dx + x = 4 \frac{e^{3x}}{3} + x + c. \left[ \because \int e^{ax} dx = \frac{e^{ax}}{a} \text{ and } \int 1 dx = x \right] \end{aligned}$$

7.  $\int x^2 \left( 1 - \frac{1}{x^2} \right) dx.$

$$\begin{aligned} \text{Sol. } \int x^2 \left( 1 - \frac{1}{x^2} \right) dx &= \int \left( x^2 - \frac{x^2}{x^2} \right) dx = \int (x^2 - 1) dx \\ &= \int x^2 dx - \int 1 dx = \frac{x^3}{3} - x + c. \left[ \because \int x^n dx = \frac{x^{n+1}}{n+1} \text{ if } n \neq -1 \right] \end{aligned}$$

8.  $\int (ax^2 + bx + c) dx.$

$$\begin{aligned} \text{Sol. } \int (ax^2 + bx + c) dx &= \int ax^2 dx + \int bx dx + \int c dx \\ &= a \int x^2 dx + b \int x dx + c \int 1 dx = a \frac{x^3}{3} + b \frac{x^2}{2} + cx + c_1 \end{aligned}$$

where  $c_1$  is the constant of integration.

9.  $\int (2x^2 + e^x) dx.$

$$\begin{aligned} \text{Sol. } \int (2x^2 + e^x) dx &= \int 2x^2 dx + \int e^x dx \\ &= 2 \int x^2 dx + \int e^x dx = 2 \frac{x^{2+1}}{2+1} + e^x + c = 2 \frac{x^3}{3} + e^x + c. \end{aligned}$$

10.  $\int \left( \sqrt{x} - \frac{1}{\sqrt{x}} \right)^2 dx.$

**Sol.**  $\int \left( \sqrt{x} - \frac{1}{\sqrt{x}} \right)^2 dx$

Opening the square =  $\int \left( (\sqrt{x})^2 + \left( \frac{1}{\sqrt{x}} \right)^2 - 2\sqrt{x} \frac{1}{\sqrt{x}} \right) dx$

=  $\int \left( x + \frac{1}{x} - 2 \right) dx = \int x dx + \int \frac{1}{x} dx - \int 2 dx$

=  $\frac{x^2}{2} + \log |x| - 2x + c. \quad \left[ \because \int 2 dx = 2 \int 1 dx = 2x \right]$

**11.**  $\int \frac{x^3 + 5x^2 - 4}{x^2} dx.$

**Sol.**  $\int \frac{x^3 + 5x^2 - 4}{x^2} dx = \int \left( \frac{x^3}{x^2} + \frac{5x^2}{x^2} - \frac{4}{x^2} \right) dx$

$\left[ \text{Using } \frac{a+b-c}{d} = \frac{a}{d} + \frac{b}{d} - \frac{c}{d} \right]$

=  $\int (x + 5 - 4x^{-2}) dx = \int x^1 dx + \int 5 dx - \int 4x^{-2} dx$

=  $\frac{x^2}{2} + 5 \int 1 dx - 4 \int x^{-2} dx = \frac{x^2}{2} + 5x - 4 \frac{x^{-2+1}}{-2+1} + c$

=  $\frac{x^2}{2} + 5x + \frac{4}{x} + c.$

**Evaluate the following integrals in Exercises 12 to 16.**

**12.**  $\int \frac{x^3 + 3x + 4}{\sqrt{x}} dx.$

**Sol.**  $\int \frac{x^3 + 3x + 4}{\sqrt{x}} dx = \int \left( \frac{x^3}{x^{1/2}} + \frac{3x}{x^{1/2}} + \frac{4}{x^{1/2}} \right) dx$

=  $\int (x^{3-1/2} + 3x^{1-1/2} + 4x^{-1/2}) dx = \int (x^{5/2} + 3x^{1/2} + 4x^{-1/2}) dx$

=  $\int x^{5/2} dx + 3 \int x^{1/2} dx + 4 \int x^{-1/2} dx$

=  $\frac{x^{5/2+1}}{\frac{5}{2}+1} + 3 \frac{x^{1/2+1}}{\frac{1}{2}+1} + 4 \frac{x^{-1/2+1}}{\frac{-1}{2}+1} + c = \frac{x^{7/2}}{\frac{7}{2}} + 3 \frac{x^{3/2}}{\frac{3}{2}} + 4 \frac{x^{1/2}}{\frac{1}{2}} + c$

=  $\frac{2}{7} x^{7/2} + 2x^{3/2} + 8x^{1/2} + c.$

**13.**  $\int \frac{x^3 - x^2 + x - 1}{x - 1} dx.$

**Sol.**  $\int \frac{x^3 - x^2 + x - 1}{x - 1} dx = \int \frac{x^2(x-1) + (x-1)}{x-1} dx$

=  $\int \frac{(x-1)(x^2+1)}{(x-1)} dx = \int (x^2+1) dx$

$$= \int x^2 dx + \int 1 dx = \frac{x^{2+1}}{2+1} + x + c = \frac{x^3}{3} + x + c.$$

**14.**  $\int (1-x)\sqrt{x} dx.$

**Sol.**  $\int (1-x)\sqrt{x} dx = \int (\sqrt{x} - x\sqrt{x}) dx$   
 $= \int (x^{1/2} - x^1 x^{1/2}) dx = \int (x^{1/2} - x^{1+1/2}) dx$   
 $= \int (x^{1/2} - x^{3/2}) dx = \frac{x^{1/2+1}}{\frac{1}{2}+1} - \frac{x^{3/2+1}}{\frac{3}{2}+1} + c$   
 $= \frac{x^{3/2}}{\frac{3}{2}} - \frac{x^{5/2}}{\frac{5}{2}} + c = \frac{2}{3} x^{3/2} - \frac{2}{5} x^{5/2} + c.$

**15.**  $\int \sqrt{x} (3x^2 + 2x + 3) dx.$

**Sol.**  $\int \sqrt{x} (3x^2 + 2x + 3) dx = \int x^{1/2} (3x^2 + 2x + 3) dx$   
 $= \int (3x^2 x^{1/2} + 2x x^{1/2} + 3x^{1/2}) dx = \int (3x^{5/2} + 2x^{3/2} + 3x^{1/2}) dx$   
 $\left( \because 2 + \frac{1}{2} = \frac{4+1}{2} = \frac{5}{2}, 1 + \frac{1}{2} = \frac{2+1}{2} = \frac{3}{2} \right)$   
 $= 3 \int x^{5/2} dx + 2 \int x^{3/2} dx + 3 \int x^{1/2} dx$   
 $= 3 \frac{x^{5/2+1}}{\frac{5}{2}+1} + 2 \frac{x^{3/2+1}}{\frac{3}{2}+1} + 3 \frac{x^{1/2+1}}{\frac{1}{2}+1} + c = 3 \frac{x^{7/2}}{\frac{7}{2}} + 2 \frac{x^{5/2}}{\frac{5}{2}} + 3 \frac{x^{3/2}}{\frac{3}{2}} + c$   
 $= \frac{6}{7} x^{7/2} + \frac{4}{5} x^{5/2} + 2x^{3/2} + c.$

**16.**  $\int (2x - 3 \cos x + e^x) dx.$

**Sol.**  $\int (2x - 3 \cos x + e^x) dx = \int 2x dx - \int 3 \cos x dx + \int e^x dx$   
 $= 2 \int x^1 dx - 3 \int \cos x dx + \int e^x dx = 2 \frac{x^2}{2} - 3 \sin x + e^x + c$   
 $= x^2 - 3 \sin x + e^x + c.$

**Evaluate the following integrals in Exercises 17 to 20.**

**17.**  $\int (2x^2 - 3 \sin x + 5\sqrt{x}) dx.$

**Sol.**  $\int (2x^2 - 3 \sin x + 5\sqrt{x}) dx$   
 $= 2 \int x^2 dx - 3 \int \sin x dx + 5 \int x^{1/2} dx$   
 $= 2 \frac{x^{2+1}}{2+1} - 3(-\cos x) + 5 \frac{x^{1/2+1}}{\frac{1}{2}+1} + c = 2 \frac{x^3}{3} + 3 \cos x + 5 \frac{x^{3/2}}{\frac{3}{2}} + c$   
 $= 2 \frac{x^3}{3} + 3 \cos x + \frac{10}{3} x^{3/2} + c.$

18.  $\int \sec x (\sec x + \tan x) dx.$

**Sol.**  $\int \sec x (\sec x + \tan x) dx = \int (\sec^2 x + \sec x \tan x) dx$   
 $= \int \sec^2 x dx + \int \sec x \tan x dx = \tan x + \sec x + c.$

19.  $\int \frac{\sec^2 x}{\operatorname{cosec}^2 x} dx.$

**Sol.**  $\int \frac{\sec^2 x}{\operatorname{cosec}^2 x} dx = \int \frac{1}{\frac{\cos^2 x}{1}} dx = \int \frac{\sin^2 x}{\cos^2 x} dx$   
 $= \int \tan^2 x dx = \int (\sec^2 x - 1) dx$   
 $(\because \sec^2 x - \tan^2 x = 1 \Rightarrow \sec^2 x - 1 = \tan^2 x)$   
 $= \int \sec^2 x dx - \int 1 dx = \tan x - x + c.$

**Note.** Similarly  $\int \cot^2 x dx = \int (\operatorname{cosec}^2 x - 1) dx$   
 $= \int \operatorname{cosec}^2 x dx - \int 1 dx = -\cot x - x + c.$

20.  $\int \frac{2 - 3 \sin x}{\cos^2 x} dx.$

**Sol.**  $\int \frac{2 - 3 \sin x}{\cos^2 x} dx = \int \left( \frac{2}{\cos^2 x} - \frac{3 \sin x}{\cos^2 x} \right) dx$   
 $= \int \left( 2 \sec^2 x - \frac{3 \sin x}{\cos x \cos x} \right) dx = \int (2 \sec^2 x - 3 \tan x \sec x) dx$   
 $= 2 \int \sec^2 x dx - 3 \int \sec x \tan x dx = 2 \tan x - 3 \sec x + c.$

21. Choose the correct answer:

The anti derivative of  $\left( \sqrt{x} + \frac{1}{\sqrt{x}} \right)$  equals

- (A)  $\frac{1}{3} x^{1/3} + 2x^{1/2} + C$  (B)  $\frac{2}{3} x^{2/3} + \frac{1}{2} x^2 + C$   
 (C)  $\frac{2}{3} x^{3/2} + 2x^{1/2} + C$  (D)  $\frac{3}{2} x^{3/2} + \frac{1}{2} x^{1/2} + C.$

**Sol.** The anti derivative of the  $\left( \sqrt{x} + \frac{1}{\sqrt{x}} \right)$   
 $= \int \left( \sqrt{x} + \frac{1}{\sqrt{x}} \right) dx = \int (x^{1/2} + x^{-1/2}) dx$   
 $= \int x^{1/2} dx + \int x^{-1/2} dx = \frac{x^{1/2+1}}{\frac{1}{2}+1} + \frac{x^{-1/2+1}}{-\frac{1}{2}+1} + C$   
 $= \frac{x^{3/2}}{\frac{3}{2}} + \frac{x^{1/2}}{\frac{1}{2}} + C = \frac{2}{3} x^{3/2} + 2x^{1/2} + C$

$\therefore$  Option (C) is the correct answer.

22. Choose the correct answer:

If  $\frac{d}{dx} f(x) = 4x^3 - \frac{3}{x^4}$  such that  $f(2) = 0$ . Then  $f(x)$  is

(A)  $x^4 + \frac{1}{x^3} - \frac{129}{8}$

(B)  $x^3 + \frac{1}{x^4} + \frac{129}{8}$

(C)  $x^4 + \frac{1}{x^3} + \frac{129}{8}$

(D)  $x^3 + \frac{1}{x^4} - \frac{129}{8}$ .

**Sol. Given:**  $\frac{d}{dx} f(x) = 4x^3 - \frac{3}{x^4}$  and  $f(2) = 0$

$\therefore$  By definition of anti derivative (i.e., Integral),

$$\begin{aligned} f(x) &= \int \left( 4x^3 - \frac{3}{x^4} \right) dx = 4 \int x^3 dx - 3 \int \frac{1}{x^4} dx \\ &= 4 \cdot \frac{x^4}{4} - 3 \int x^{-4} dx = x^4 - 3 \frac{x^{-3}}{-3} + c \end{aligned}$$

or  $f(x) = x^4 + \frac{1}{(x^3)} + c \quad \dots(i)$

**To find c.** Let us make use of  $f(2) = 0$  (given)

Putting  $x = 2$  on both sides of (i),

$$f(2) = 16 + \frac{1}{8} + c \quad \text{or} \quad 0 = \frac{128+1}{8} + c$$

( $\because f(2) = 0$  (given))

or  $c + \frac{129}{8} = 0 \quad \text{or} \quad c = -\frac{129}{8}$

Putting  $c = -\frac{129}{8}$  in (i),  $f(x) = x^4 + \frac{1}{(x^3)} - \frac{129}{8}$

$\therefore$  Option (A) is the correct answer.

## Exercise 7.2

Integrate the functions in Exercises 1 to 8:

1.  $\frac{2x}{1+x^2}$

**Sol.** To evaluate  $\int \frac{2x}{1+x^2} dx$

**Put**  $1+x^2 = t$ . Therefore  $2x = \frac{dt}{dx}$  or  $2x dx = dt$

$$\therefore \int \frac{2x}{1+x^2} dx = \int \frac{dt}{t} = \int \frac{1}{t} dt = \log |t| + c$$

Putting  $t = 1+x^2$ ,  $= \log |1+x^2| + c = \log (1+x^2) + c$   
( $\because 1+x^2 > 0$ . Therefore  $|1+x^2| = 1+x^2$ )

2.  $\frac{(\log x)^2}{x}$ .

**Sol.** To evaluate  $\int \frac{(\log x)^2}{x} dx$

**Put  $\log x = t$ .** Therefore  $\frac{1}{x} = \frac{dt}{dx} \Rightarrow \frac{dx}{x} = dt$

$$\therefore \int \frac{(\log x)^2}{x} dx = \int t^2 dt = \frac{t^3}{3} + c$$

Putting  $t = \log x$ ,  $= \frac{1}{3} (\log x)^3 + c$ .

3.  $\frac{1}{x + x \log x}$

**Sol.** To evaluate  $\int \frac{1}{x + x \log x} dx = \int \frac{1}{x(1 + \log x)} dx$

**Put  $1 + \log x = t$ .** Therefore  $\frac{1}{x} = \frac{dt}{dx} \Rightarrow \frac{dx}{x} = dt$

$$\therefore \int \frac{1}{x + x \log x} dx = \int \frac{1}{1 + \log x} \frac{dx}{x} = \int \frac{1}{t} dt = \log |t| + c$$

Putting  $t = 1 + \log x$ ,  $\log |1 + \log x| + c$ .

4.  $\sin x \sin (\cos x)$

**Sol.** To evaluate  $\int \sin x \sin (\cos x) dx = - \int \sin (\cos x) (-\sin x) dx$

**Put  $\cos x = t$ .** Therefore  $-\sin x = \frac{dt}{dx}$

$$\therefore -\sin x dx = dt$$

$$\begin{aligned} \therefore \int \sin x \sin (\cos x) dx &= - \int \sin (\cos x) (-\sin x dx) \\ &= - \int \sin t dt = -(-\cos t) + c \\ &= \cos t + c \end{aligned}$$

Putting  $t = \cos x$ ,  $= \cos(\cos x) + c$ .

5.  $\sin(ax + b) \cos(ax + b)$

**Sol.** To evaluate  $\int \sin(ax + b) \cos(ax + b) dx$

$$= \frac{1}{2} \int 2 \sin(ax + b) \cos(ax + b) dx = \frac{1}{2} \int \sin 2(ax + b) dx$$

( $\because 2 \sin \theta \cos \theta = \sin 2\theta$ )

$$= \frac{1}{2} \int \sin (2ax + 2b) dx = \frac{1}{2} \frac{[-\cos(2ax + 2b)]}{2a \rightarrow \text{Coeff. of } x} + c$$

$$= \frac{-1}{4a} \cos 2(ax + b) + c.$$

6.  $\sqrt{ax + b}$

**Sol.** To evaluate  $\int \sqrt{ax + b} dx = \int (ax + b)^{1/2} dx$

$$= \frac{(ax + b)^{\frac{1}{2} + 1}}{\left(\frac{1}{2} + 1\right) a \rightarrow \text{Coeff. of } x} + c = \frac{(ax + b)^{\frac{3}{2}}}{\frac{3}{2}a} + c$$

$$\left[ \because \int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{a(n+1)} + c \text{ if } n \neq -1 \right]$$

$$= \frac{2}{3a} (ax + b)^{3/2} + c.$$

**7.  $x\sqrt{x+2}$**

**Sol.** To evaluate  $\int x\sqrt{x+2} \, dx$

$$\begin{aligned} &= \int x\sqrt{x+2} \, dx = \int ((x+2) - 2)\sqrt{x+2} \, dx \\ &= \int \left( (x+2)(x+2)^{\frac{1}{2}} - 2(x+2)^{\frac{1}{2}} \right) dx = \int \left( (x+2)^{\frac{3}{2}} - 2(x+2)^{\frac{1}{2}} \right) dx \\ &= \int (x+2)^{\frac{3}{2}} dx - 2 \int (x+2)^{\frac{1}{2}} dx \\ &= \frac{(x+2)^{\frac{3}{2}+1}}{\left(\frac{3}{2}+1\right)} - 2 \frac{(x+2)^{\frac{1}{2}+1}}{\left(\frac{1}{2}+1\right)} + c = \frac{(x+2)^{\frac{5}{2}}}{\frac{5}{2}} - 2 \frac{(x+2)^{\frac{3}{2}}}{\frac{3}{2}} + c \\ &= \frac{2}{5} (x+2)^{5/2} - \frac{4}{3} (x+2)^{3/2} + c. \end{aligned}$$

**OR**

To evaluate  $\int x\sqrt{x+2} \, dx$

Put  $\sqrt{\text{Linear}} = t$ , i.e.,  $\sqrt{x+2} = t$ .

Squaring  $x+2 = t^2$  ( $\Rightarrow x = t^2 - 2$ )

$$\therefore \frac{dx}{dt} = 2t, \text{ i.e., } \frac{dx}{dt} = 2t \text{ or } dx = 2t \, dt$$

$$\begin{aligned} \therefore \int x\sqrt{x+2} \, dx &= \int (t^2 - 2)t \cdot 2t \, dt = \int 2t^2(t^2 - 2) \, dt \\ &= \int 2t^2(t^2 - 2) \, dt = 2 \int t^4 \, dt - 4 \int t^2 \, dt = 2 \frac{t^5}{5} - 4 \frac{t^3}{3} + c \end{aligned}$$

$$\begin{aligned} \text{Putting } t &= \sqrt{x+2}, \quad = \frac{2}{5} (\sqrt{x+2})^5 - \frac{4}{3} (\sqrt{x+2})^3 + c \\ &= \frac{2}{5} (x+2)^{5/2} - \frac{4}{3} (x+2)^{3/2} + c = \frac{2}{5} (x+2)^{5/2} - \frac{4}{3} (x+2)^{3/2} + c. \end{aligned}$$

**8.  $x\sqrt{1+2x^2}$**

**Sol.** To evaluate  $\int x\sqrt{1+2x^2} \, dx$

$$\begin{aligned} \text{Let } I &= \int x\sqrt{1+2x^2} \, dx = \frac{1}{4} \int \sqrt{1+2x^2} (4x \, dx) \quad \dots(i) \\ &\quad \left[ \because \frac{d}{dx} (1+2x^2) = 0 + 2 \cdot 2x = 4x \right] \end{aligned}$$

Put  $1 + 2x^2 = t$ . Therefore  $4x = \frac{dt}{dx}$  or  $4x \, dx = dt$

$$\therefore \text{ From (i), } I = \frac{1}{4} \int \sqrt{t} \, dt = \frac{1}{4} \int t^{1/2} \, dt$$

$$= \frac{1}{4} \cdot \frac{t^{3/2}}{\frac{3}{2}} + c = \frac{1}{4} \cdot \frac{2}{3} t^{3/2} + c$$

Putting  $t = 1 + 2x^2$ ,  $= \frac{1}{6} (1 + 2x^2)^{3/2} + c$ .

**Integrate the functions in Exercises 9 to 17:**

**9.  $(4x + 2) \sqrt{x^2 + x + 1}$ .**

**Sol.** Let  $I = \int (4x + 2) \sqrt{x^2 + x + 1} \, dx = \int 2(2x + 1) \sqrt{x^2 + x + 1} \, dx$   
 $= \int 2\sqrt{x^2 + x + 1} (2x + 1) \, dx \quad \dots(i)$

**Put  $x^2 + x + 1 = t$ .** Therefore  $(2x + 1) = \frac{dt}{dx}$

$\therefore (2x + 1) \, dx = dt$

$\therefore$  From (i),  $I = \int 2\sqrt{t} \, dt = 2 \int t^{1/2} \, dt$

$$= 2 \cdot \frac{t^{3/2}}{\frac{3}{2}} + c = \frac{4}{3} t^{3/2} + c$$

Putting  $t = x^2 + x + 1$ ,  $I = \frac{4}{3} (x^2 + x + 1)^{3/2} + c$ .

**10.  $\frac{1}{x - \sqrt{x}}$**

**Sol.** Let  $I = \int \frac{1}{x - \sqrt{x}} \, dx \quad \dots(i)$

**Put  $\sqrt{x} = t$ , i.e.,  $x = t^2$ .** Therefore  $\frac{dx}{dt} = 2t$  or  $dx = 2t \, dt$

Squaring  $x = t^2$ . Therefore  $\frac{dx}{dt} = 2t$  or  $dx = 2t \, dt$

$\therefore$  From (i),  $I = \int \frac{1}{t^2 - t} \cdot 2t \, dt = 2 \int \frac{t}{t(t-1)} \, dt$

$$= 2 \int \frac{1}{t-1} \, dt = 2 \log |t-1| + c \left( \because \int \frac{1}{ax+b} \, dx = \frac{1}{a} \log |ax+b| \right)$$

Putting  $t = \sqrt{x}$ ,  $I = 2 \log |\sqrt{x} - 1| + c$ .

**11.  $\frac{x}{\sqrt{x+4}}$ ,  $x > 0$**

**Sol.** Let  $I = \int \frac{x}{\sqrt{x+4}} \, dx \quad \dots(i)$

$$= \int \frac{x+4-4}{\sqrt{x+4}} \, dx = \int \left( \frac{x+4}{\sqrt{x+4}} - \frac{4}{\sqrt{x+4}} \right) \, dx$$

$$= \int \sqrt{x+4} \, dx - 4 \int \frac{1}{\sqrt{x+4}} \, dx \left[ \because \frac{t}{\sqrt{t}} = \frac{t\sqrt{t}}{\sqrt{t}\sqrt{t}} = \frac{t\sqrt{t}}{t} = \sqrt{t} \right]$$

$$\begin{aligned}
&= \int (x+4)^{1/2} dx - 4 \int (x+4)^{-1/2} dx \\
&= \frac{(x+4)^{3/2}}{\frac{3}{2}(1)} - \frac{4(x+4)^{1/2}}{\frac{1}{2}(1)} + c = \frac{2}{3} (x+4)^{3/2} - 8(x+4)^{1/2} + c \\
&= \frac{2}{3} (x+4) \sqrt{x+4} - 8\sqrt{x+4} + c \\
&\quad \left[ \because t^{3/2} = t^{\frac{2}{2} + \frac{1}{2}} = t^{1 + \frac{1}{2}} = t^1 \cdot t^{1/2} = t\sqrt{t} \right] \\
&= 2\sqrt{x+4} \left( \frac{x+4}{3} - 4 \right) + c = 2\sqrt{x+4} \left( \frac{x+4-12}{3} \right) + c \\
&= \frac{2}{3} \sqrt{x+4} (x-8) + c.
\end{aligned}$$

**OR**

Put  $\sqrt{\text{Linear}} = t$ , i.e.,  $\sqrt{x+4} = t$ .  
 Squaring  $x+4 = t^2 \Rightarrow x = t^2 - 4$ .

Therefore  $\frac{dx}{dt} = 2t$  or  $dx = 2t dt$

$$\begin{aligned}
\therefore I &= \int \frac{x}{\sqrt{x+4}} dx = \int \frac{t^2 - 4}{t} \cdot 2t dt \\
&= 2 \int (t^2 - 4) dt = 2 \left[ \int t^2 dt - 4 \int 1 dt \right] \\
&= 2 \left[ \frac{t^3}{3} - 4t \right] + c = \frac{2t}{3} (t^2 - 12) + c.
\end{aligned}$$

$$\begin{aligned}
\text{Putting } t &= \sqrt{x+4}, = \frac{2}{3} \sqrt{x+4} (x+4-12) + c \\
&= \frac{2}{3} \sqrt{x+4} (x-8) + c.
\end{aligned}$$

**12.  $(x^3 - 1)^{1/3} x^5$**

**Sol.** Let  $I = \int (x^3 - 1)^{1/3} x^5 dx = \int (x^3 - 1)^{1/3} x^3 x^2 dx$

$$= \frac{1}{3} \int (x^3 - 1)^{1/3} x^3 (3x^2 dx) \quad \dots(i) \quad \left[ \because \frac{d}{dx} (x^3 - 1) = 3x^2 \right]$$

$$\text{Put } x^3 - 1 = t \quad \Rightarrow \quad x^3 = t + 1$$

$$\therefore 3x^2 = \frac{dt}{dx} \quad \Rightarrow \quad 3x^2 dx = dt$$

$$\therefore \text{ From (i), } I = \frac{1}{3} \int t^{1/3} (t+1) dt$$

$$= \frac{1}{3} \int (t^{4/3} + t^{1/3}) dt \quad \left[ \because \frac{1}{3} + 1 = \frac{1+3}{3} = \frac{4}{3} \right]$$

$$= \frac{1}{3} \left( \int t^{4/3} dt + \int t^{1/3} dt \right)$$

$$= \frac{1}{3} \left( \frac{t^{7/3}}{\frac{3}{3}} + \frac{t^{4/3}}{\frac{4}{3}} \right) + c = \frac{1}{3} \left( \frac{3}{7} t^{7/3} + \frac{3}{4} t^{4/3} \right) + c = \frac{1}{7} t^{7/3} + \frac{1}{4} t^{4/3} + c$$

$$\text{Putting } t = x^3 - 1, = \frac{1}{7} (x^3 - 1)^{7/3} + \frac{1}{4} (x^3 - 1)^{4/3} + c.$$

13.  $\frac{x^2}{(2+3x^3)^3}$

**Sol.** Let  $I = \int \frac{x^2}{(2+3x^3)^3} dx$

$$= \frac{1}{9} \int \frac{9x^2}{(2+3x^3)^3} dx \quad \dots(i) \quad \left[ \because \frac{d}{dx} (2+3x^3) = 9x^2 \right]$$

Put  $2+3x^3 = t$ . Therefore  $9x^2 = \frac{dt}{dx} \Rightarrow 9x^2 dx = dt$

$$\therefore \text{ From (i), } I = \frac{1}{9} \int t^{-3} dt = \frac{1}{9} \frac{t^{-2}}{-2} + c = \frac{-1}{18t^2} + c$$

$$\text{Putting } t = 2+3x^3; = \frac{-1}{18(2+3x^3)^2} + c.$$

14.  $\frac{1}{x(\log x)^m}, x > 0$

(Important)

**Sol.** Let  $I = \int \frac{1}{x(\log x)^m} dx \ (x > 0) \Rightarrow I = \int \frac{\frac{1}{x} dx}{(\log x)^m} \quad \dots(i)$

Put  $\log x = t$ . Therefore  $\frac{1}{x} = \frac{dt}{dx} \Rightarrow \frac{dx}{x} = dt$

$$\therefore \text{ From (i), } I = \int \frac{dt}{t^m} = \int t^{-m} dt = \frac{t^{-m+1}}{-m+1} + c$$

(Assuming  $m \neq 1$ )

$$\text{Putting } t = \log x, = \frac{(\log x)^{1-m}}{1-m} + c.$$

15.  $\frac{x}{9-4x^2}$

**Sol.** Let  $I = \int \frac{x}{9-4x^2} dx = \frac{-1}{8} \int \frac{-8x}{9-4x^2} dx \quad \dots(i)$

$$\left[ \because \frac{d}{dx} (9-4x^2) = -8x \right]$$

Put  $9-4x^2 = t$ . Therefore  $-8x = \frac{dt}{dx} \Rightarrow -8x dx = dt$

$$\therefore \text{ From (i), } I = \frac{-1}{8} \int \frac{dt}{t} = \frac{-1}{8} \int \frac{1}{t} dt = \frac{-1}{8} \log |t| + c$$

$$\text{Putting } t = 9-4x^2, = \frac{-1}{8} \log |9-4x^2| + c.$$

16.  $e^{2x+3}$

**Sol.**  $\int e^{2x+3} dx = \frac{e^{2x+3}}{2 \rightarrow \text{Coeff. of } x} + c$   $\left[ \because \int e^{ax+b} dx = \frac{e^{ax+b}}{a} \right]$   
 $= \frac{1}{2} e^{2x+3} + c.$

17.  $\frac{x}{e^{x^2}}$

**Sol.** Let  $I = \int \frac{x}{(e^{x^2})} dx = \frac{1}{2} \int \frac{2x}{(e^{x^2})} dx$  ... (i)

Put  $x^2 = t$ . Therefore  $2x = \frac{dt}{dx} \Rightarrow 2x dx = dt.$

$\therefore$  From (i),  $I = \frac{1}{2} \int \frac{dt}{(e^t)} = \frac{1}{2} \int e^{-t} dt$   
 $= \frac{1}{2} \frac{e^{-t}}{-1 \rightarrow \text{Coeff. of } t} + c = \frac{-1}{2(e^t)} + c$

Putting  $t = x^2$ ,  $I = \frac{-1}{2(e^{x^2})} + c.$

**Integrate the functions in Exercises 18 to 26:**

18.  $\frac{e^{\tan^{-1} x}}{1+x^2}$

**Sol.** Let  $I = \int \frac{e^{\tan^{-1} x}}{1+x^2} dx$  ... (i)

Put  $\tan^{-1} x = t.$

$\therefore \frac{1}{1+x^2} = \frac{dt}{dx} \Rightarrow \frac{dx}{1+x^2} = dt$

$\therefore$  From (i),  $I = \int e^t dt = e^t + c = e^{\tan^{-1} x} + c.$

19.  $\frac{e^{2x}-1}{e^{2x}+1}$

**Sol.** Let  $I = \int \frac{e^{2x}-1}{e^{2x}+1} dx$

**Multiplying every term in integrand by  $e^{-x}$ ,**

$I = \int \frac{e^x - e^{-x}}{e^x + e^{-x}} dx$  ... (i)  $[\because e^{2x} \cdot e^{-x} = e^{2x-x} = e^x]$

Put denominator  $e^x + e^{-x} = t$

$\therefore e^x + e^{-x} \frac{d}{dx} (-x) = \frac{dt}{dx} \Rightarrow (e^x - e^{-x}) dx = dt$

$\therefore$  From (i),  $I = \int \frac{dt}{t} = \int \frac{1}{t} dt = \log |t| + c$

Putting  $t = e^x + e^{-x}$ ,  $I = \log |e^x + e^{-x}| + c$  or  $I = \log (e^x + e^{-x}) + c$

$\left[ \because e^x + e^{-x} = e^x + \frac{1}{(e^x)} > 0 \text{ for all real } x \text{ and hence } |e^x + e^{-x}| = e^x + e^{-x} \right]$

20.  $\frac{e^{2x} - e^{-2x}}{e^{2x} + e^{-2x}}$

**Sol.** Let  $I = \int \frac{e^{2x} - e^{-2x}}{e^{2x} + e^{-2x}} dx = \frac{1}{2} \int \frac{2(e^{2x} - e^{-2x})}{e^{2x} + e^{-2x}} dx \quad \dots(i)$

Put denominator  $e^{2x} + e^{-2x} = t$

$\therefore e^{2x} \frac{d}{dx} 2x + e^{-2x} \frac{d}{dx} (-2x) = \frac{dt}{dx}$

$\Rightarrow e^{2x} \cdot 2 - 2e^{-2x} = \frac{dt}{dx} \Rightarrow 2(e^{2x} - e^{-2x}) dx = dt$

$\therefore$  From (i),  $I = \frac{1}{2} \int \frac{dt}{t} = \frac{1}{2} \log |t| + c$

Putting  $t = e^{2x} + e^{-2x}$ ,  $= \frac{1}{2} \log |e^{2x} + e^{-2x}| + c = \frac{1}{2} \log(e^{2x} + e^{-2x}) + c$   
 $[\because e^{2x} + e^{-2x} > 0 \Rightarrow |e^{2x} + e^{-2x}| = e^{2x} + e^{-2x}]$

21.  $\tan^2 (2x - 3)$

**Sol.**  $\int \tan^2 (2x - 3) dx = \int (\sec^2 (2x - 3) - 1) dx (\because \tan^2 \theta = \sec^2 \theta - 1)$   
 $= \int \sec^2 (2x - 3) dx - \int 1 dx$   
 $= \frac{\tan (2x - 3)}{2 \rightarrow \text{Coeff. of } x} - x + c = \frac{1}{2} \tan (2x - 3) - x + c$

$\left[ \because \int \sec^2 (ax + b) dx = \frac{1}{a} \tan (ax + b) + c \right]$

22.  $\sec^2 (7 - 4x)$

**Sol.**  $\int \sec^2 (7 - 4x) dx = \frac{\tan (7 - 4x)}{-4 \rightarrow \text{Coeff. of } x} + c$   
 $\left[ \because \int \sec^2 (ax + b) dx = \frac{1}{a} \tan (ax + b) + c \right]$   
 $= \frac{-1}{4} \tan (7 - 4x) + c.$

23.  $\frac{\sin^{-1} x}{\sqrt{1-x^2}}$

**Sol.** Let  $I = \int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx \quad \dots(i)$

Put  $\sin^{-1} x = t \quad \therefore \frac{1}{\sqrt{1-x^2}} = \frac{dt}{dx} \Rightarrow \frac{dx}{\sqrt{1-x^2}} = dt$

$\therefore$  From (i),  $I = \int t dt = \frac{t^2}{2} + c$

Putting  $t = \sin^{-1} x$ ,  $I = \frac{1}{2} (\sin^{-1} x)^2 + c.$

24.  $\frac{2 \cos x - 3 \sin x}{6 \cos x + 4 \sin x}$

**Sol.** Let  $I = \int \frac{2 \cos x - 3 \sin x}{6 \cos x + 4 \sin x} dx = \int \frac{2 \cos x - 3 \sin x}{2(2 \sin x + 3 \cos x)} dx$   
 $= \frac{1}{2} \int \frac{2 \cos x - 3 \sin x}{2 \sin x + 3 \cos x} dx \quad \dots(i)$

Put DENOMINATOR  $2 \sin x + 3 \cos x = t$

$$\therefore 2 \cos x - 3 \sin x = \frac{dt}{dx} \Rightarrow (2 \cos x - 3 \sin x) dx = dt$$

$$\therefore \text{From (i), } I = \frac{1}{2} \int \frac{dt}{t} = \frac{1}{2} \log |t| + c.$$

$$\text{Putting } t = 2 \sin x + 3 \cos x, = \frac{1}{2} \log |2 \sin x + 3 \cos x| + c.$$

**25.**  $\frac{1}{\cos^2 x (1 - \tan x)^2}$

**Sol.** Let  $I = \int \frac{1}{\cos^2 x (1 - \tan x)^2} dx = \int \frac{\sec^2 x}{(1 - \tan x)^2} dx$   
 $= - \int \frac{-\sec^2 x}{(1 - \tan x)^2} dx \quad \dots(i)$

Put  $1 - \tan x = t$ .

$$\therefore -\sec^2 x = \frac{dt}{dx} \Rightarrow -\sec^2 x dx = dt$$

$$\therefore \text{From (i), } I = - \int \frac{dt}{t^2} = - \int t^{-2} dt$$

$$= - \frac{t^{-1}}{-1} + c = \frac{1}{t} + c = \frac{1}{1 - \tan x} + c.$$

**26.**  $\frac{\cos \sqrt{x}}{\sqrt{x}}$

**Sol.** Let  $I = \int \frac{\cos \sqrt{x}}{\sqrt{x}} dx \quad \dots(i)$

Put  $\sqrt{\text{Linear}} = t$ , i.e.,  $\sqrt{x} = t$

$$\text{Squaring, } x = t^2. \text{ Therefore } \frac{dx}{dt} = 2t \quad \therefore dx = 2t dt$$

$$\therefore \text{From (i), } I = \int \frac{\cos t}{t} 2t dt = 2 \int \cos t dt = 2 \sin t + c$$

Putting  $t = \sqrt{x}$ ,  $I = 2 \sin \sqrt{x} + c$ .

**Integrate the functions in Exercises 27 to 37:**

**27.**  $\sqrt{\sin 2x} \cos 2x$

**Sol.** Let  $I = \int \sqrt{\sin 2x} \cos 2x dx = \frac{1}{2} \int \sqrt{\sin 2x} (2 \cos 2x dx) \quad \dots(i)$

Put  $\sin 2x = t$

$$\therefore \cos 2x \frac{d}{dx} (2x) = \frac{dt}{dx} \Rightarrow 2 \cos 2x dx = dt$$

$$\begin{aligned}\therefore \text{ From (i), } I &= \frac{1}{2} \int \sqrt{t} \, dt = \frac{1}{2} \int t^{1/2} \, dt \\ &= \frac{1}{2} \frac{t^{\frac{1}{2}+1}}{\frac{1}{2}+1} + c = \frac{1}{2} \frac{t^{3/2}}{\frac{3}{2}} + c = \frac{1}{3} (\sin 2x)^{3/2} + c.\end{aligned}$$

28.  $\frac{\cos x}{\sqrt{1 + \sin x}}$

**Sol.** Let  $I = \int \frac{\cos x}{\sqrt{1 + \sin x}} \, dx$  ... (i)

**Put  $1 + \sin x = t$**

$\therefore \cos x = \frac{dt}{dx} \quad \text{or} \quad \cos x \, dx = dt$

$$\begin{aligned}\therefore \text{ From (i), } I &= \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} \, dt = \frac{t^{-\frac{1}{2}+1}}{-\frac{1}{2}+1} + c \\ &= \frac{t^{1/2}}{\frac{1}{2}} + c = 2\sqrt{t} + c = 2\sqrt{1 + \sin x} + c.\end{aligned}$$

29.  $\cot x \log \sin x$

**Sol.** Let  $I = \int \cot x \log \sin x \, dx$  ... (i)

**Put  $\log \sin x = t$**

$\therefore \frac{1}{\sin x} \frac{d}{dx} (\sin x) = \frac{dt}{dx} \quad \text{or} \quad \frac{1}{\sin x} \cos x = \frac{dt}{dx}$   
or  $\cot x \, dx = dt$

$\therefore \text{ From (i), } I = \int t \, dt = \frac{t^2}{2} + c = \frac{1}{2} (\log \sin x)^2 + c.$

30.  $\frac{\sin x}{1 + \cos x}$

**Sol.** Let  $I = \int \frac{\sin x}{1 + \cos x} \, dx = - \int \frac{-\sin x}{1 + \cos x} \, dx$  ... (i)

**Put  $1 + \cos x = t$ .** Therefore  $-\sin x = \frac{dt}{dx}$

$\therefore -\sin x \, dx = dt$

$\therefore \text{ From (i), } I = - \int \frac{dt}{t} = - \log |t| + c$

Putting  $t = 1 + \cos x$ ,  $= - \log |1 + \cos x| + c.$

31.  $\frac{\sin x}{(1 + \cos x)^2}$

**Sol.** Let  $I = \int \frac{\sin x}{(1 + \cos x)^2} \, dx = - \int \frac{-\sin x \, dx}{(1 + \cos x)^2}$  ... (i)

**Put  $1 + \cos x = t$ .** Therefore  $-\sin x = \frac{dt}{dx}$

$$\Rightarrow -\sin x \, dx = dt$$

$$\begin{aligned}\therefore \text{ From (i), } I &= - \int \frac{dt}{t^2} = - \int t^{-2} \, dt = \frac{-t^{-1}}{-1} + c \\ &= \frac{1}{t} + c = \frac{1}{1 + \cos x} + c.\end{aligned}$$

**32.  $\frac{1}{1 + \cot x}$**

**Sol.** Let  $I = \int \frac{1}{1 + \cot x} \, dx = \int \frac{1}{1 + \frac{\cos x}{\sin x}} \, dx = \int \frac{1}{\left(\frac{\sin x + \cos x}{\sin x}\right)} \, dx$

$$= \int \frac{\sin x}{\sin x + \cos x} \, dx = \frac{1}{2} \int \frac{2 \sin x}{\sin x + \cos x} \, dx = \frac{1}{2} \int \frac{\sin x + \sin x}{\sin x + \cos x} \, dx$$

Adding and subtracting  $\cos x$  in the numerator of integrand,

$$\begin{aligned}I &= \frac{1}{2} \int \frac{\sin x + \cos x - \cos x + \sin x}{\sin x + \cos x} \, dx \\ &= \frac{1}{2} \int \frac{(\sin x + \cos x) - (\cos x - \sin x)}{\sin x + \cos x} \, dx \\ &= \frac{1}{2} \int \left( \frac{\sin x + \cos x}{\sin x + \cos x} - \frac{(\cos x - \sin x)}{\sin x + \cos x} \right) dx \left[ \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right] \\ &= \frac{1}{2} \int \left( 1 - \frac{(\cos x - \sin x)}{\sin x + \cos x} \right) dx \\ &= \frac{1}{2} \left[ \int 1 \, dx - \int \frac{\cos x - \sin x}{\sin x + \cos x} \, dx \right] = \frac{1}{2} [x - I_1] \quad \dots(i)\end{aligned}$$

where  $I_1 = \int \frac{\cos x - \sin x}{\sin x + \cos x} \, dx$

Put DENOMINATOR  $\sin x + \cos x = t$

$$\therefore \cos x - \sin x = \frac{dt}{dx} \Rightarrow (\cos x - \sin x) \, dx = dt$$

$$\therefore I_1 = \int \frac{dt}{t} = \log |t| = \log |\sin x + \cos x|.$$

**Note. Alternative solution for finding  $I_1$**

$$I_1 = \int \frac{\cos x - \sin x}{\sin x + \cos x} \, dx = \log |\sin x + \cos x|$$

$$\left[ \because \int \frac{f'(x)}{f(x)} \, dx = \log |f(x)| \right]$$

Putting this value of  $I_1$  in (i), required integral

$$= \frac{1}{2} [x - \log |\sin x + \cos x|] + c.$$

33.  $\frac{1}{1 - \tan x}$

**Sol.** Let  $I = \int \frac{1}{1 - \tan x} dx = \int \frac{1}{1 - \frac{\sin x}{\cos x}} dx = \int \frac{1}{\left(\frac{\cos x - \sin x}{\cos x}\right)} dx$

$$= \int \frac{\cos x}{\cos x - \sin x} dx = \frac{1}{2} \int \frac{2 \cos x}{\cos x - \sin x} dx = \frac{1}{2} \int \frac{\cos x + \cos x}{\cos x - \sin x} dx$$

Subtracting and adding  $\sin x$  in the Numerator,

$$= \frac{1}{2} \int \frac{\cos x - \sin x + \sin x + \cos x}{\cos x - \sin x} dx$$

$$= \frac{1}{2} \int \left( \frac{\cos x - \sin x}{\cos x - \sin x} + \frac{\sin x + \cos x}{\cos x - \sin x} \right) dx = \frac{1}{2} \int \left( 1 + \frac{\sin x + \cos x}{\cos x - \sin x} \right) dx$$

$$= \frac{1}{2} \left[ \int 1 dx - \int \frac{-\sin x - \cos x}{\cos x - \sin x} dx \right]$$

$$= \frac{1}{2} [x - \log |\cos x - \sin x|] + c \quad \left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right]$$

**Note.** Alternative solution for evaluating  $\int \frac{-\sin x - \cos x}{\cos x - \sin x} dx$ , put denominator  $\cos x - \sin x = t$ .

34.  $\frac{\sqrt{\tan x}}{\sin x \cos x}$

**Sol.** Let  $I = \int \frac{\sqrt{\tan x}}{\sin x \cos x} dx = \int \frac{\sqrt{\tan x}}{\frac{\sin x}{\cos x} \cos x \cos x} dx$

$$= \int \frac{\sqrt{\tan x}}{\tan x \cos^2 x} dx = \int \frac{\sec^2 x}{\sqrt{\tan x}} dx \quad \dots(i) \quad \left[ \because \frac{\sqrt{t}}{t} = \frac{1}{\sqrt{t}} \right]$$

**Put  $\tan x = t$ .**

$$\therefore \sec^2 x = \frac{dt}{dx} \Rightarrow \sec^2 x dx = dt$$

$\therefore$  From (i),

$$I = \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{\frac{1}{2}} + c = 2\sqrt{t} + c = 2\sqrt{\tan x} + c.$$

35.  $\frac{(1 + \log x)^2}{x}$

**Sol.** Let  $I = \int \frac{(1 + \log x)^2}{x} dx \quad \dots(i)$

**Put  $1 + \log x = t$**

$$\therefore \frac{1}{x} = \frac{dt}{dx} \Rightarrow \frac{dx}{x} = dt$$

$$\therefore \text{From (i), } I = \int t^2 dt = \frac{t^3}{3} + c = \frac{1}{3} (1 + \log x)^3 + c.$$

36.  $\frac{(x+1)(x+\log x)^2}{x}$

Sol. Let  $I = \int \frac{(x+1)(x+\log x)^2}{x} dx \quad \dots(i)$

Put  $x + \log x = t$

$$\therefore 1 + \frac{1}{x} = \frac{dt}{dx} \Rightarrow \frac{x+1}{x} = \frac{dt}{dx} \Rightarrow \left(\frac{x+1}{x}\right) dx = dt$$

$$\therefore \text{From (i), } I = \int t^2 dt = \frac{t^3}{3} + c$$

Putting  $t = x + \log x$ ,  $\frac{1}{3} (x + \log x)^3 + c$ .

37.  $\frac{x^3 \sin(\tan^{-1} x^4)}{1+x^8}$

Sol. Let  $I = \int \frac{x^3 \sin(\tan^{-1} x^4)}{1+x^8} dx = \frac{1}{4} \int \sin(\tan^{-1} x^4) \cdot \frac{4x^3}{1+x^8} dx \dots(i)$

Put  $(\tan^{-1} x^4) = t$

[Rule for  $\int \sin(f(x)) f'(x) dx$ ; put  $f(x) = t$ ]

$$\therefore \frac{1}{1+(x^4)^2} \frac{d}{dx} x^4 = \frac{dt}{dx} \left[ \because \frac{d}{dx} \tan^{-1} f(x) = \frac{1}{1+(f(x))^2} \frac{d}{dx} f(x) \right]$$

$$\Rightarrow \frac{4x^3}{1+x^8} dx = dt$$

$\therefore$  From (i),

$$I = \frac{1}{4} \int \sin t dt = -\frac{1}{4} \cos t + c = -\frac{1}{4} \cos(\tan^{-1} x^4) + c.$$

**Choose the correct answer in Exercises 38 and 39:**

38.  $\int \frac{10x^9 + 10^x \log_e 10 dx}{x^{10} + 10^x}$  equals

(A)  $10^x - x^{10} + C$

(C)  $(10^x - x^{10})^{-1} + C$

(B)  $10^x + x^{10} + C$

(D)  $\log(10^x + x^{10}) + C$ .

Sol. Let  $I = \int \frac{10x^9 + 10^x \log_e 10}{x^{10} + 10^x} dx \quad \dots(i)$

Put  $x^{10} + 10^x = t$

$$\therefore (10x^9 + 10^x \log_e 10) dx = dt \quad \left[ \because \frac{d}{dx} (a^x) = a^x \log_e a \right]$$

$$\therefore \text{From (i), } I = \int \frac{dt}{t} = \log |t| + c$$

Putting  $t = x^{10} + 10^x$ ,  $I = \log |x^{10} + 10^x| + c$

or  $I = \log(10^x + x^{10}) + c$ .

$\therefore$  Option (D) is the correct answer.

**OR**

$$\int \frac{10x^9 + 10^x \log_e 10}{x^{10} + 10^x} dx = \int \frac{f'(x)}{f(x)} dx = \log |f(x)| + c$$

$$= \log |x^{10} + 10^x| + c$$

∴ Option (D) is the correct answer.

39.  $\int \frac{dx}{\sin^2 x \cos^2 x}$  equals

(A)  $\tan x + \cot x + C$

(B)  $\tan x - \cot x + C$

(C)  $\tan x \cot x + C$

(D)  $\tan x - \cot 2x + C.$


**Sol.**  $\int \frac{dx}{\sin^2 x \cos^2 x} = \int \frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} dx \quad [\because 1 = \sin^2 x + \cos^2 x]$

$$= \int \left( \frac{\sin^2 x}{\sin^2 x \cos^2 x} + \frac{\cos^2 x}{\sin^2 x \cos^2 x} \right) dx \quad \left[ \because \frac{a+b}{c} = \frac{a}{c} + \frac{b}{c} \right]$$

$$= \int \left( \frac{1}{\cos^2 x} + \frac{1}{\sin^2 x} \right) dx = \int (\sec^2 x + \operatorname{cosec}^2 x) dx$$

$$= \int \sec^2 x dx + \int \operatorname{cosec}^2 x dx = \tan x - \cot x + c$$

∴ Option (B) is the correct answer.



### Exercise 7.3

Find the integrals of the following functions in Exercises 1 to 9:

1.  $\sin^2(2x + 5)$

**Sol.**  $\int \sin^2(2x + 5) \, dx = \int \frac{1}{2} (1 - \cos 2(2x + 5)) \, dx$

$$\left[ \because \sin^2 \theta = \frac{1}{2} (1 - \cos 2\theta); \text{ put } \theta = 2x + 5 \right]$$
$$= \frac{1}{2} \int (1 - \cos(4x + 10)) \, dx = \frac{1}{2} \left[ \int 1 \, dx - \int \cos(4x + 10) \, dx \right]$$
$$= \frac{1}{2} \left[ x - \frac{\sin(4x + 10)}{4 \rightarrow \text{Coeff. of } x} \right] + c = \frac{1}{2} x - \frac{1}{8} \sin(4x + 10) + c.$$

2.  $\sin 3x \cos 4x$

**Sol.**  $\int \sin 3x \cos 4x \, dx = \frac{1}{2} \int 2 \sin 3x \cos 4x \, dx$

$$= \frac{1}{2} \int (\sin(3x + 4x) + \sin(3x - 4x)) \, dx$$
$$[\because 2 \sin A \cos B = \sin(A + B) + \sin(A - B)]$$
$$= \frac{1}{2} \int (\sin 7x + \sin(-x)) \, dx = \frac{1}{2} \int (\sin 7x - \sin x) \, dx$$
$$= \frac{1}{2} \left[ \int \sin 7x \, dx - \int \sin x \, dx \right] = \frac{1}{2} \left[ \frac{-\cos 7x}{7} - (-\cos x) \right] + c$$
$$= \frac{-1}{14} \cos 7x + \frac{1}{2} \cos x + c.$$

3.  $\cos 2x \cos 4x \cos 6x$

**Sol.**  $\cos 2x \cos 4x \cos 6x = \frac{1}{2} (2 \cos 6x \cos 4x) \cos 2x$

$$= \frac{1}{2} [\cos(6x + 4x) + \cos(6x - 4x)] \cos 2x$$
$$[\because 2 \cos x \cdot \cos y = \cos(x + y) + \cos(x - y)]$$

$$\begin{aligned}
&= \frac{1}{2} (\cos 10x + \cos 2x) \cos 2x = \frac{1}{4} (2 \cos 10x \cos 2x + 2 \cos^2 2x) \\
&= \frac{1}{4} [\cos (10x + 2x) + \cos (10x - 2x) + 1 + \cos 4x] \\
&= \frac{1}{4} (\cos 12x + \cos 8x + \cos 4x + 1) \\
\therefore \int \cos 2x \cos 4x \cos 6x \, dx &= \frac{1}{4} \int (\cos 12x + \cos 8x + \cos 4x + 1) \, dx \\
&= \frac{1}{4} \left[ \int \cos 12x \, dx + \int \cos 8x \, dx + \int \cos 4x \, dx + \int 1 \, dx \right] \\
&= \frac{1}{4} \left( \frac{\sin 12x}{12} + \frac{\sin 8x}{8} + \frac{\sin 4x}{4} + x \right) + c.
\end{aligned}$$

**Note.** We know that  $\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$

$$\therefore 4 \sin^3 \theta = 3 \sin \theta - \sin 3\theta$$

$$\text{Dividing by 4, } \sin^3 \theta = \frac{3}{4} \sin \theta - \frac{1}{4} \sin 3\theta \quad \dots(i)$$

$$\text{Similarly, } \cos^3 \theta = \frac{3}{4} \cos \theta + \frac{1}{4} \cos 3\theta \quad \dots(ii)$$

$$[\because \cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta]$$

#### 4. $\sin^3 (2x + 1)$

**Sol.** To evaluate  $\int \sin^3 (2x + 1) \, dx$

We know by Eqn. (i) of above note that  $\sin^3 \theta = \frac{3}{4} \sin \theta - \frac{1}{4} \sin 3\theta$

Putting  $\theta = 2x + 1$ , we have

$$\begin{aligned}
\sin^3 (2x + 1) &= \frac{3}{4} \sin (2x + 1) - \frac{1}{4} \sin 3 (2x + 1) \\
&= \frac{3}{4} \sin (2x + 1) - \frac{1}{4} \sin (6x + 3) \\
\therefore \int \sin^3 (2x + 1) \, dx &= \frac{3}{4} \int \sin (2x + 1) \, dx - \frac{1}{4} \int \sin (6x + 3) \, dx \\
&= \frac{3}{4} \left( \frac{-\cos (2x + 1)}{2} \right) - \frac{1}{4} \left( \frac{-\cos (6x + 3)}{6 \rightarrow \text{Coeff. of } x} \right) + c \\
&= \frac{-3}{8} \cos (2x + 1) + \frac{1}{24} \cos (6x + 3) + c.
\end{aligned}$$

**OR**

**To integrate  $\sin^n x$  where  $n$  is odd, put  $\cos x = t$ .**

$$\begin{aligned}
\therefore \int \sin^3 (2x + 1) \, dx &= \int \sin^2 (2x + 1) \sin (2x + 1) \, dx \\
&= \frac{-1}{2} \int [1 - \cos^2 (2x + 1)] (-2 \sin (2x + 1)) \, dx \quad \dots(i)
\end{aligned}$$

**Put  $\cos (2x + 1) = t$**

$$\therefore -\sin (2x + 1) \frac{d}{dx} (2x + 1) = \frac{dt}{dx} \therefore -2 \sin (2x + 1) \, dx = dt$$

$$\therefore \text{From (i), the given integral} = \frac{-1}{2} \int (1 - t^2) \, dt$$

$$\begin{aligned}
 &= \frac{-1}{2} \left( t - \frac{t^3}{3} \right) + c = \frac{-1}{2} t + \frac{1}{6} t^3 + c \\
 &= \frac{-1}{2} \cos (2x + 1) + \frac{1}{6} \cos^3 (2x + 1) + c.
 \end{aligned}$$

**5.  $\sin^3 x \cos^3 x$**

**Sol.**  $\int \sin^3 x \cos^3 x \, dx = \int (\sin x \cos x)^3 \, dx$

$$\begin{aligned}
 &= \int \left( \frac{1}{2} 2 \sin x \cos x \right)^3 \, dx = \int \left( \frac{1}{2} \sin 2x \right)^3 \, dx \\
 &= \frac{1}{8} \int \sin^3 2x \, dx = \frac{1}{8} \int \left( \frac{3}{4} \sin 2x - \frac{1}{4} \sin 6x \right) \, dx \\
 &\quad \left( \text{Putting } \theta = 2x \text{ in } \sin^3 \theta = \frac{3}{4} \sin \theta - \frac{1}{4} \sin 3\theta \right) \\
 &= \frac{3}{32} \int \sin 2x \, dx - \frac{1}{32} \int \sin 6x \, dx \\
 &= \frac{-3}{32} \frac{\cos 2x}{2} - \frac{1}{32} \left( \frac{-\cos 6x}{6} \right) + c = \frac{-3}{64} \cos 2x + \frac{1}{192} \cos 6x + c.
 \end{aligned}$$

**OR**

To evaluate  $\int \sin^3 x \cos^3 x \, dx$ , Put either  $\sin x = t$  or  $\cos x = t$ .  
(The form of answer given in N.C.E.R.T. book II can be obtained by putting  $\cos x = t$ )

**6.  $\sin x \sin 2x \sin 3x$**

**Sol.**  $\sin x \sin 2x \sin 3x = \frac{1}{2} (2 \sin 3x \sin 2x) \sin x$

$$\begin{aligned}
 &= \frac{1}{2} [\cos (3x - 2x) - \cos (3x + 2x)] \sin x \\
 &\quad [\because 2 \sin x \sin y = \cos (x - y) - \cos (x + y)] \\
 &= \frac{1}{2} (\cos x - \cos 5x) \sin x = \frac{1}{4} (2 \cos x \sin x - 2 \cos 5x \sin x) \\
 &= \frac{1}{4} [\sin 2x - \{\sin (5x + x) - \sin (5x - x)\}] \\
 &\quad [\because 2 \cos x \sin y = \sin (x + y) - \sin (x - y)] \\
 &= \frac{1}{4} (\sin 2x - \sin 6x + \sin 4x) \\
 \therefore \int \sin x \sin 2x \sin 3x \, dx &= \frac{1}{4} \int (\sin 2x + \sin 4x - \sin 6x) \, dx \\
 &= \frac{1}{4} \left[ \int \sin 2x \, dx + \int \sin 4x \, dx - \int \sin 6x \, dx \right] \\
 &= \frac{1}{4} \left( -\frac{\cos 2x}{2} - \frac{\cos 4x}{4} + \frac{\cos 6x}{6} \right) + c.
 \end{aligned}$$

**7.  $\sin 4x \sin 8x$**

**Sol.**  $\int \sin 4x \sin 8x \, dx = \frac{1}{2} \int 2 \sin 4x \sin 8x \, dx$

$$\begin{aligned}
 &= \frac{1}{2} \int [\cos (4x - 8x) - \cos (4x + 8x)] \, dx \\
 &\quad [\because 2 \sin A \sin B = \cos (A - B) - \cos (A + B)]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \int (\cos(-4x) - \cos 12x) dx = \frac{1}{2} \int (\cos 4x - \cos 12x) dx \\
 &\quad [\because \cos(-\theta) = \cos \theta] \\
 &= \frac{1}{2} \left[ \int \cos 4x dx - \int \cos 12x dx \right] = \frac{1}{2} \left[ \frac{\sin 4x}{4} - \frac{\sin 12x}{12} \right] + c.
 \end{aligned}$$

8.  $\frac{1 - \cos x}{1 + \cos x}$

**Sol.**  $\int \frac{1 - \cos x}{1 + \cos x} dx = \int \frac{2 \sin^2 \frac{x}{2}}{2 \cos^2 \frac{x}{2}} dx = \int \tan^2 \frac{x}{2} dx$

$$\left( \because 1 - \cos \theta = 2 \sin^2 \frac{\theta}{2} \text{ and } 1 + \cos \theta = 2 \cos^2 \frac{\theta}{2} \right)$$

$$= \int \left( \sec^2 \frac{x}{2} - 1 \right) dx \quad (\because \tan^2 \theta = \sec^2 \theta - 1)$$

$$= \int \sec^2 \frac{x}{2} dx - \int 1 dx = \frac{\tan \frac{x}{2}}{\frac{1}{2} \rightarrow \text{Coeff. of } x} - x + c = 2 \tan \frac{x}{2} - x + c.$$

9.  $\frac{\cos x}{1 + \cos x}$

**Sol.**  $\int \frac{\cos x}{1 + \cos x} dx$

Adding and subtracting 1 in the numerator of integrand,

$$\begin{aligned}
 &= \int \frac{1 + \cos x - 1}{1 + \cos x} dx = \int \left( \frac{1 + \cos x}{1 + \cos x} - \frac{1}{1 + \cos x} \right) dx \left( \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right) \\
 &= \int \left( 1 - \frac{1}{2 \cos^2 \frac{x}{2}} \right) dx = \int 1 dx - \frac{1}{2} \int \sec^2 \frac{x}{2} dx \\
 &= x - \frac{1}{2} \frac{\tan \frac{x}{2}}{\frac{1}{2}} + c = x - \tan \frac{x}{2} + c.
 \end{aligned}$$

**Find the integrals of the functions in Exercises 10 to 18:**

10.  $\sin^4 x$

**Sol.**  $\int \sin^4 x dx = \int (\sin^2 x)^2 dx = \int \left( \frac{1 - \cos 2x}{2} \right)^2 dx$

$$\begin{aligned}
 &= \int \frac{(1 - \cos 2x)^2}{4} dx = \frac{1}{4} \int (1 + \cos^2 2x - 2 \cos 2x) dx \\
 &= \frac{1}{4} \int \left( 1 + \left( \frac{1 + \cos 4x}{2} \right) - 2 \cos 2x \right) dx \left[ \because \cos^2 \theta = \frac{1 + \cos 2\theta}{2} \right]
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{4} \int \left( \frac{2 + 1 + \cos 4x - 4 \cos 2x}{2} \right) dx = \frac{1}{8} \int (3 + \cos 4x - 4 \cos 2x) dx \\
&= \frac{1}{8} \left[ 3 \int 1 dx + \int \cos 4x dx - 4 \int \cos 2x dx \right] \\
&= \frac{1}{8} \left[ 3x + \frac{\sin 4x}{4} - \frac{4 \sin 2x}{2} \right] + c = \frac{3}{8}x + \frac{1}{32} \sin 4x - \frac{1}{4} \sin 2x + c
\end{aligned}$$

**11.  $\cos^4 2x$**

**Sol.**  $\int \cos^4 2x dx = \int (\cos^2 2x)^2 dx$

$$\begin{aligned}
&= \int \left( \frac{1 + \cos 4x}{2} \right)^2 dx = \int \frac{1}{4} (1 + \cos 4x)^2 dx \\
&= \frac{1}{4} \int (1 + \cos^2 4x + 2 \cos 4x) dx \\
&= \frac{1}{4} \int \left( 1 + \frac{1 + \cos 8x}{2} + 2 \cos 4x \right) dx \quad \left[ \because \cos^2 \theta = \frac{1 + \cos 2\theta}{2} \right] \\
&= \frac{1}{4} \int \left( \frac{2 + 1 + \cos 8x + 4 \cos 4x}{2} \right) dx = \frac{1}{8} \int (3 + \cos 8x + 4 \cos 4x) dx \\
&= \frac{1}{8} \left[ 3 \int 1 dx + \int \cos 8x dx + 4 \int \cos 4x dx \right] \\
&= \frac{1}{8} \left[ 3x + \frac{\sin 8x}{8} + \frac{4 \sin 4x}{4} \right] + c = \frac{3}{8}x + \frac{1}{64} \sin 8x + \frac{1}{8} \sin 4x + c
\end{aligned}$$

**12.  $\frac{\sin^2 x}{1 + \cos x}$**

**Sol.**  $\int \frac{\sin^2 x}{1 + \cos x} dx = \int \frac{1 - \cos^2 x}{1 + \cos x} dx = \int \frac{(1 - \cos x)(1 + \cos x)}{1 + \cos x} dx$

$$= \int (1 - \cos x) dx = \int 1 dx - \int \cos x dx = x - \sin x + c.$$

**Note.** It may be noted that letters  $a, b, c, d, \dots, q$  of English Alphabet and letters  $\alpha, \beta, \gamma, \delta$  of Greek Alphabet are generally treated as constants.

**13.  $\frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha}$**

**Sol.**  $\int \frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha} dx = \int \frac{(2 \cos^2 x - 1) - (2 \cos^2 \alpha - 1)}{\cos x - \cos \alpha} dx$

$$\begin{aligned}
&= \int \frac{2 \cos^2 x - 1 - 2 \cos^2 \alpha + 1}{\cos x - \cos \alpha} dx = \int \frac{2 \cos^2 x - 2 \cos^2 \alpha}{\cos x - \cos \alpha} dx \\
&= 2 \int \frac{\cos^2 x - \cos^2 \alpha}{\cos x - \cos \alpha} dx = 2 \int \frac{(\cos x - \cos \alpha)(\cos x + \cos \alpha)}{(\cos x - \cos \alpha)} dx \\
&= 2 \int (\cos x + \cos \alpha) dx = 2 \left[ \int \cos x dx + \int \cos \alpha dx \right] \\
&= 2 [\sin x + \cos \alpha \int 1 dx] = 2 [\sin x + (\cos \alpha) x] + c \\
&= 2 \sin x + 2x \cos \alpha + c.
\end{aligned}$$

**Remark.**  $\int \sin a \, dx = \sin a \int 1 \, dx = x \sin a$ .

Please note that  $\int \sin a \, dx \neq -\cos a$ .

**14.  $\frac{\cos x - \sin x}{1 + \sin 2x}$**

**Sol.** Let  $I = \int \frac{\cos x - \sin x}{1 + \sin 2x} dx = \int \frac{\cos x - \sin x}{\cos^2 x + \sin^2 x + 2 \sin x \cos x} dx$   
$$= \int \frac{\cos x - \sin x}{(\cos x + \sin x)^2} dx \quad \dots(i)$$

**Put  $\cos x + \sin x = t$ .**

$\therefore -\sin x + \cos x = \frac{dt}{dx}$ . Therefore  $(\cos x - \sin x) dx = dt$ .

$\therefore$  From (i),  $I = \int \frac{dt}{t^2} = \int t^{-2} dt = \frac{t^{-1}}{-1} + c$

$\Rightarrow I = \frac{-1}{t} + c = \frac{-1}{\cos x + \sin x} + c$ .

**15.  $\tan^3 2x \sec 2x$**

**Sol.** Let  $I = \int \tan^3 2x \sec 2x \, dx = \int \tan^2 2x \tan 2x \sec 2x \, dx$   
$$= \int (\sec^2 2x - 1) \sec 2x \tan 2x \, dx \quad [\because \tan^2 \theta = \sec^2 \theta - 1]$$
  
$$= \frac{1}{2} \int (\sec^2 2x - 1)(2 \sec 2x \tan 2x) \, dx \quad \dots(i)$$

**Put  $\sec 2x = t$ .** Therefore  $\sec 2x \tan 2x \frac{d}{dx} (2x) = \frac{dt}{dx}$

$\therefore 2 \sec 2x \tan 2x \, dx = dt$

$\therefore$  From (i),  $I = \frac{1}{2} \int (t^2 - 1) dt = \frac{1}{2} \left( \int t^2 dt - \int 1 dt \right)$   
$$= \frac{1}{2} \left( \frac{t^3}{3} - t \right) + c = \frac{1}{6} t^3 - \frac{1}{2} t + c$$

Putting  $t = \sec 2x$ ,  $= \frac{1}{6} \sec^3 2x - \frac{1}{2} \sec 2x + c$ .

**16.  $\tan^4 x$**

**Sol.**  $\int \tan^4 x \, dx = \int \tan^2 x \tan^2 x \, dx = \int \tan^2 x (\sec^2 x - 1) \, dx$   
$$= \int (\tan^2 x \sec^2 x - \tan^2 x) \, dx = \int \tan^2 x \sec^2 x \, dx - \int \tan^2 x \, dx$$
  
$$= \int \tan^2 x \sec^2 x \, dx - \int (\sec^2 x - 1) \, dx$$
  
$$= \int \tan^2 x \sec^2 x \, dx - \int \sec^2 x \, dx + \int 1 \, dx$$
  
$$\downarrow$$

For this integral, put  **$\tan x = t$** .

$\therefore \sec^2 x = \frac{dt}{dx}$  or  $\sec^2 x \, dx = dt$

$$= \int t^2 dt - \tan x + x + c = \frac{t^3}{3} - \tan x + x + c$$

Put  $t = \tan x$ ,  $= \frac{1}{3} \tan^3 x - \tan x + x + c$ .

17.  $\frac{\sin^3 x + \cos^3 x}{\sin^2 x \cos^2 x}$

**Sol.**  $\int \frac{\sin^3 x + \cos^3 x}{\sin^2 x \cos^2 x} dx = \int \left( \frac{\sin^3 x}{\sin^2 x \cos^2 x} + \frac{\cos^3 x}{\sin^2 x \cos^2 x} \right) dx$

$$\left( \because \frac{a+b}{c} = \frac{a}{c} + \frac{b}{c} \right)$$

$$= \int \left( \frac{\sin x}{\cos^2 x} + \frac{\cos x}{\sin^2 x} \right) dx = \int \left( \frac{\sin x}{\cos x \cos x} + \frac{\cos x}{\sin x \sin x} \right) dx$$

$$= \int (\tan x \sec x + \cot x \operatorname{cosec} x) dx$$

$$= \int \sec x \tan x dx + \int \operatorname{cosec} x \cot x dx = \sec x - \operatorname{cosec} x + c.$$

18.  $\frac{\cos 2x + 2 \sin^2 x}{\cos^2 x}$

**Sol.**  $\int \frac{\cos 2x + 2 \sin^2 x}{\cos^2 x} dx = \int \frac{(1 - 2 \sin^2 x) + 2 \sin^2 x}{\cos^2 x} dx$

$$= \int \frac{1}{\cos^2 x} dx = \int \sec^2 x dx = \tan x + c.$$

**Integrate the functions in Exercises 19 to 22:**

**Note.** Method to evaluate  $\int \frac{1}{\sin^p x \cos^q x} dx$  if  $(p + q)$  is a negative even integer ( $= -n$  (say)); then multiply Numerator and Denominator of integrand by  $\sec^n x$ .

19.  $\frac{1}{\sin x \cos^3 x}$

**Sol.** Let  $I = \int \frac{1}{\sin x \cos^3 x} dx$  ... (i)

Here  $p + q = -1 - 3 = -4$  is a negative even integer.

So multiplying both Numerator and Denominator of integrand of (i) by  $\sec^4 x$ ,

$$I = \int \frac{\sec^4 x}{\sin x \cos^3 x \sec^4 x} dx = \int \frac{\sec^4 x}{\tan x} dx$$

$$\left( \because \sin x \cos^3 x \sec^4 x = \sin x \cos^3 x \cdot \frac{1}{\cos^4 x} = \frac{\sin x}{\cos x} = \tan x \right)$$

or  $I = \int \frac{\sec^2 x \sec^2 x}{\tan x} dx = \int \frac{(1 + \tan^2 x) \sec^2 x}{\tan x} dx$  ... (ii)

**Put  $\tan x = t$**

$$\therefore \sec^2 x = \frac{dt}{dx} \Rightarrow \sec^2 x \, dx = dt$$

$$\begin{aligned} \therefore \text{From (ii), } I &= \int \frac{(1+t^2)}{t} \, dt = \int \left( \frac{1}{t} + \frac{t^2}{t} \right) dt \\ &= \int \left( \frac{1}{t} + t \right) dt = \int \frac{1}{t} \, dt + \int t \, dt = \log |t| + \frac{t^2}{2} + c \\ \text{Putting } t &= \tan x, = \log |\tan x| + \frac{1}{2} \tan^2 x + c. \end{aligned}$$

**20.  $\frac{\cos 2x}{(\cos x + \sin x)^2}$**

**Sol.** Let  $I = \int \frac{\cos 2x}{(\cos x + \sin x)^2} \, dx = \int \frac{\cos^2 x - \sin^2 x}{(\cos x + \sin x)^2} \, dx$   
 $= \int \frac{(\cos x + \sin x)(\cos x - \sin x)}{(\cos x + \sin x)(\cos x + \sin x)} \, dx = \int \frac{\cos x - \sin x}{\cos x + \sin x} \, dx \quad \dots(i)$

Put DENOMINATOR  $\cos x + \sin x = t$

$$\therefore -\sin x + \cos x = \frac{dt}{dx} \Rightarrow (\cos x - \sin x) \, dx = dt$$

$$\therefore \text{From (i), } I = \int \frac{dt}{t} = \log |t| + c = \log |\cos x + \sin x| + c$$

**Note.** Another method to evaluate integral (i) is, apply

$$\int \frac{f'(x)}{f(x)} \, dx = \log |f(x)|.$$

**21.  $\sin^{-1}(\cos x)$**

**Sol.**  $\int \sin^{-1}(\cos x) \, dx = \int \sin^{-1} \sin \left( \frac{\pi}{2} - x \right) \, dx$   
 $= \int \left( \frac{\pi}{2} - x \right) \, dx = \int \frac{\pi}{2} \, dx - \int x \, dx$   
 $= \frac{\pi}{2} \int 1 \, dx - \int x^1 \, dx = \frac{\pi}{2} x - \frac{x^2}{2} + c.$

**22.  $\frac{1}{\cos(x-a)\cos(x-b)}$**

**Sol.** Let  $I = \int \frac{1}{\cos(x-a)\cos(x-b)} \, dx \quad \dots(i)$

Here  $(x-a) - (x-b) = x-a-x+b = b-a \quad \dots(ii)$

By looking at Eqn. (ii), dividing and multiplying the integrand in (i) by  $\sin(b-a)$ ,

$$\begin{aligned} I &= \frac{1}{\sin(b-a)} \int \frac{\sin(b-a)}{\cos(x-a)\cos(x-b)} \, dx \\ &= \frac{1}{\sin(b-a)} \int \frac{\sin[(x-a)-(x-b)]}{\cos(x-a)\cos(x-b)} \, dx \quad [\text{By (ii)}] \\ &= \frac{1}{\sin(b-a)} \int \frac{\sin(x-a)\cos(x-b) - \cos(x-a)\sin(x-b)}{\cos(x-a)\cos(x-b)} \, dx \\ &\quad [\because \sin(A-B) = \sin A \cos B - \cos A \sin B] \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\sin(b-a)} \int \left[ \frac{\sin(x-a) \cos(x-b)}{\cos(x-a) \cos(x-b)} - \frac{\cos(x-a) \sin(x-b)}{\cos(x-a) \cos(x-b)} \right] dx \\
 &\quad \left( \because \frac{A-B}{C} = \frac{A}{C} - \frac{B}{C} \right) \\
 &= \frac{1}{\sin(b-a)} \int [\tan(x-a) - \tan(x-b)] dx \\
 &= \frac{1}{\sin(b-a)} [-\log |\cos(x-a)| + \log |\cos(x-b)|] + c \\
 &\quad \left( \because \int \tan x \, dx = -\log |\cos x| \right) \\
 &= \frac{1}{\sin(b-a)} \log \left| \frac{\cos(x-b)}{\cos(x-a)} \right| + c. \left( \because \log m - \log n = \log \frac{m}{n} \right)
 \end{aligned}$$

Choose the correct answer in Exercises 23 and 24:

23.  $\int \frac{\sin^2 x - \cos^2 x}{\sin^2 x \cos^2 x} dx$  is equal to

(A)  $\tan x + \cot x + C$

(B)  $\tan x + \operatorname{cosec} x + C$

(C)  $-\tan x + \cot x + C$

(D)  $\tan x + \sec x + C$

Sol.  $\int \frac{\sin^2 x - \cos^2 x}{\sin^2 x \cos^2 x} dx$

$$= \int \left( \frac{\sin^2 x}{\sin^2 x \cos^2 x} - \frac{\cos^2 x}{\sin^2 x \cos^2 x} \right) dx \quad \left[ \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right]$$

$$= \int \left( \frac{1}{\cos^2 x} - \frac{1}{\sin^2 x} \right) dx = \int (\sec^2 x - \operatorname{cosec}^2 x) dx$$

$$= \int \sec^2 x \, dx - \int \operatorname{cosec}^2 x \, dx = \tan x - (-\cot x) + C$$

$$= \tan x + \cot x + C \quad \therefore \text{Option (A) is the correct answer.}$$

24.  $\int \frac{e^x(1+x)}{\cos^2(e^x x)} dx$  equals

(A)  $-\cot(e^x) + C$

(B)  $\tan(xe^x) + C$

(C)  $\tan(e^x) + C$

(D)  $\cot(e^x) + C$

Sol. Let  $I = \int \frac{e^x(1+x)}{\cos^2(e^x x)} dx \quad \dots(i)$

Put  $e^x \cdot x = t$

[To evaluate  $\int$  (T-function or Inverse T-function  $f(x)) f'(x) dx$ , put  $f(x) = t$ ]

$$\text{Applying Product Rule, } e^x \cdot 1 + xe^x = \frac{dt}{dx}$$

$$\text{or } e^x(1+x) dx = dt$$

$$\therefore \text{ From (i), } I = \int \frac{dt}{\cos^2 t} = \int \sec^2 t \, dt$$

$$= \tan t + C = \tan(xe^x) + C \therefore \text{Option (B) is the correct answer.}$$

## Exercise 7.4

Integrate the following functions in Exercises 1 to 9:

1.  $\frac{3x^2}{x^6 + 1}$

**Sol.** Let  $I = \int \frac{3x^2}{x^6 + 1} dx = \int \frac{3x^2}{(x^3)^2 + 1^2} dx \quad \dots(i)$

**Put**  $x^3 = t$

$$\therefore 3x^2 = \frac{dt}{dx} \Rightarrow 3x^2 dx = dt$$

$$\therefore \text{From (i), } I = \int \frac{dt}{t^2 + 1^2} = \frac{1}{1} \tan^{-1} \frac{t}{1} + C$$

$$\left[ \because \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right]$$

Putting  $t = x^3$ ;  $= \tan^{-1} (x^3) + C$ .

**Note.**  $ax^2 + b$  ( $a \neq 0$ ) is called a **pure quadratic**.

2.  $\frac{1}{\sqrt{1 + 4x^2}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{1 + 4x^2}} dx = \int \frac{1}{\sqrt{(2x)^2 + 1^2}} dx$

Using  $\int \frac{1}{\sqrt{x^2 + a^2}} dx = \log \left| x + \sqrt{x^2 + a^2} \right|$ ,

$$I = \frac{\log \left| (2x) + \sqrt{(2x)^2 + 1^2} \right|}{2 \rightarrow \text{Coeff. of } x} + C = \frac{1}{2} \log \left| 2x + \sqrt{4x^2 + 1} \right| + C.$$

3.  $\frac{1}{\sqrt{(2-x)^2 + 1}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{(2-x)^2 + 1}} dx = \int \frac{1}{\sqrt{(2-x)^2 + 1}} dx$

Using  $\int \frac{1}{\sqrt{x^2 + a^2}} dx = \log \left| x + \sqrt{x^2 + a^2} \right|$ ,

$$= \frac{\log \left| (2-x) + \sqrt{(2-x)^2 + 1^2} \right|}{-1 \rightarrow \text{Coeff. of } x} + C$$

$$= -\log \left| 2-x + \sqrt{4+x^2-4x+1} \right| + C$$

$$= \log \left| \frac{1}{2-x + \sqrt{x^2-4x+5}} \right| + C.$$

$$\left[ \because -\log \frac{m}{n} = -(\log m - \log n) = \log n - \log m = \log \frac{n}{m} \right]$$

4.  $\frac{1}{\sqrt{9-25x^2}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{9-25x^2}} dx = \int \frac{1}{\sqrt{3^2-(5x)^2}} dx$

$$= \frac{\sin^{-1} \frac{5x}{3}}{5 \rightarrow \text{Coeff. of } x} + C \quad \left[ \because \int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1} \frac{x}{a} \right]$$

$$= \frac{1}{5} \sin^{-1} \left( \frac{5x}{3} \right) + C.$$

5.  $\frac{3x}{1+2x^4}$

**Sol.** Let  $I = \int \frac{3x}{1+2x^4} dx = \frac{3}{2} \int \frac{2x}{1+2(x^2)^2} dx \quad \dots(i)$

**Put  $x^2 = t$ .**  $\therefore 2x = \frac{dt}{dx} \Rightarrow 2x dx = dt$

$\therefore$  From (i),  $I = \frac{3}{2} \int \frac{dt}{1+2t^2} = \frac{3}{2} \int \frac{1}{(\sqrt{2}t)^2+1^2} dt$

$$= \frac{3}{2} \frac{\frac{1}{\sqrt{2}} \tan^{-1} \frac{\sqrt{2}t}{1}}{\sqrt{2} \rightarrow \text{Coeff. of } t} + C = \frac{3}{2\sqrt{2}} \tan^{-1} (\sqrt{2}t) + C$$

Putting  $t = x^2$ ,  $= \frac{3}{2\sqrt{2}} \tan^{-1} (\sqrt{2}x^2) + C.$

6.  $\frac{x^2}{1-x^6}$

**Sol.** Let  $I = \int \frac{x^2}{1-x^6} dx = \int \frac{x^2}{1-(x^3)^2} dx = \frac{1}{3} \int \frac{3x^2}{1-(x^3)^2} dx$

**Put  $x^3 = t$ .** Therefore  $3x^2 = \frac{dt}{dx} \Rightarrow 3x^2 dx = dt.$

$\therefore I = \frac{1}{3} \int \frac{dt}{1-t^2} = \frac{1}{3} \int \frac{1}{1^2-t^2} dt = \frac{1}{3} \frac{1}{2 \times 1} \log \left| \frac{1+t}{1-t} \right| + C$

$$\left[ \because \int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| \right]$$

Putting  $t = x^3$ ,  $= \frac{1}{6} \log \left| \frac{1+x^3}{1-x^3} \right| + C.$

7.  $\frac{x-1}{\sqrt{x^2-1}}$

**Sol.** Let  $I = \int \frac{x-1}{\sqrt{x^2-1}} dx = \int \left( \frac{x}{\sqrt{x^2-1}} - \frac{1}{\sqrt{x^2-1}} \right) dx$

$$\begin{aligned}
 &= \int \frac{x}{\sqrt{x^2-1}} dx - \int \frac{1}{\sqrt{x^2-1^2}} dx \\
 &= \frac{1}{2} \int \frac{2x}{\sqrt{x^2-1}} dx - \log \left| x + \sqrt{x^2-1^2} \right| \quad \dots(i) \\
 &\left( \because \int \frac{1}{\sqrt{x^2-a^2}} dx = \log \left| x + \sqrt{x^2-a^2} \right| \right)
 \end{aligned}$$

$$\text{Let } I_1 = \int \frac{2x}{\sqrt{x^2-1}} dx$$

$$\text{Put } x^2 - 1 = t. \text{ Therefore } 2x = \frac{dt}{dx} \Rightarrow 2x dx = dt$$

$$\therefore I_1 = \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{\frac{1}{2}} = 2\sqrt{t} = 2\sqrt{x^2-1} + C$$

$$\text{Putting this value of } I_1 = \int \frac{2x}{\sqrt{x^2-1}} dx \text{ in (i),}$$

$$\begin{aligned}
 I &= \frac{1}{2} (2\sqrt{x^2-1} + C) - \log \left| x + \sqrt{x^2-1} \right| \\
 &= \sqrt{x^2-1} + \frac{C}{2} - \log \left| x + \sqrt{x^2-1} \right| \\
 &= \sqrt{x^2-1} - \log \left| x + \sqrt{x^2-1} \right| + C_1 \text{ where } C_1 = \frac{C}{2}.
 \end{aligned}$$

$$8. \frac{x^2}{\sqrt{x^6+a^6}}$$

$$\text{Sol. Let } I = \int \frac{x^2}{\sqrt{x^6+a^6}} dx = \frac{1}{3} \int \frac{3x^2}{\sqrt{(x^3)^2+a^6}} dx \quad \dots(i)$$

$$\text{Put } x^3 = t. \text{ Therefore } 3x^2 = \frac{dt}{dx} \Rightarrow 3x^2 dx = dt.$$

$$\therefore \text{ From (i), } I = \frac{1}{3} \int \frac{dt}{\sqrt{t^2+a^6}} = \frac{1}{3} \int \frac{1}{\sqrt{t^2+(a^3)^2}} dt$$

$$= \frac{1}{3} \log \left| t + \sqrt{t^2+(a^3)^2} \right| + C \left[ \because \int \frac{1}{\sqrt{x^2+a^2}} dx = \log \left| x + \sqrt{x^2+a^2} \right| \right]$$

$$\text{Putting } t = x^3, = \frac{1}{3} \log \left| x^3 + \sqrt{x^6+a^6} \right| + C.$$

$$9. \frac{\sec^2 x}{\sqrt{\tan^2 x + 4}}$$

$$\text{Sol. Let } I = \int \frac{\sec^2 x}{\sqrt{\tan^2 x + 4}} dx \quad \dots(i)$$

**Put  $\tan x = t$ .**  $\therefore \sec^2 x = \frac{dt}{dx} \Rightarrow \sec^2 x \, dx = dt$

$\therefore$  From (i),  $I = \int \frac{dt}{\sqrt{t^2 + 4}} = \int \frac{1}{\sqrt{t^2 + 2^2}} \, dt$

$$= \log \left| t + \sqrt{t^2 + 2^2} \right| + C \left[ \because \int \frac{1}{\sqrt{x^2 + a^2}} \, dx = \log \left| x + \sqrt{x^2 + a^2} \right| \right]$$

Putting  $t = \tan x$ ,  $I = \log \left| \tan x + \sqrt{\tan^2 x + 4} \right| + C$ .

**Integrate the following functions in Exercises 10 to 18:**

**Note. Rule to evaluate**

$$\int \frac{1}{\text{Quadratic}} \, dx \text{ or } \int \frac{1}{\sqrt{\text{Quadratic}}} \, dx \text{ or } \int \sqrt{\text{Quadratic}} \, dx$$

Write Quadratic. Take coefficient of  $x^2$  common to make it unity. Then complete squares by adding and subtracting

$$\left( \frac{1}{2} \text{ coefficient of } x \right)^2$$

10.  $\frac{1}{\sqrt{x^2 + 2x + 2}}$

**Sol.**  $\int \frac{1}{\sqrt{x^2 + 2x + 2}} \, dx = \int \frac{1}{\sqrt{x^2 + 2x + 1 + 1}} \, dx = \int \frac{1}{\sqrt{(x+1)^2 + 1^2}} \, dx$

Using  $\int \frac{1}{\sqrt{x^2 + a^2}} \, dx = \log \left| x + \sqrt{x^2 + a^2} \right|$

$$= \log \left| x + 1 + \sqrt{(x+1)^2 + 1^2} \right| + c = \log \left| x + 1 + \sqrt{x^2 + 2x + 2} \right| + c.$$

11.  $\frac{1}{9x^2 + 6x + 5}$

**Sol.** Let  $I = \int \frac{1}{9x^2 + 6x + 5} \, dx$  ... (i)

$$\int \frac{1}{\text{Quadratic}} \, dx$$

Here Quadratic expression =  $9x^2 + 6x + 5$

Making coefficient of  $x^2$  unity, =  $9 \left( x^2 + \frac{6x}{9} + \frac{5}{9} \right)$

$$= 9 \left( x^2 + \frac{2x}{3} + \frac{5}{9} \right)$$

To complete squares, adding and subtracting  $\left( \frac{1}{2} \text{ Coefficient of } x \right)^2$

$$= \left( \left( \frac{1}{2} \times \frac{2}{3} \right)^2 = \left( \frac{1}{3} \right)^2 = \frac{1}{9} \right) = 9 \left( x^2 + \frac{2x}{3} + \left( \frac{1}{3} \right)^2 - \frac{1}{9} + \frac{5}{9} \right)$$

$$= 9 \left[ \left( x + \frac{1}{3} \right)^2 + \frac{4}{9} \right] \Rightarrow 9x^2 + 6x + 5 = 9 \left[ \left( x + \frac{1}{3} \right)^2 + \left( \frac{2}{3} \right)^2 \right]$$

Putting this value in (i),  $I = \int \frac{1}{9 \left[ \left( x + \frac{1}{3} \right)^2 + \left( \frac{2}{3} \right)^2 \right]} dx$

$$= \frac{1}{9} \int \frac{1}{\left( x + \frac{1}{3} \right)^2 + \left( \frac{2}{3} \right)^2} dx$$

$$= \frac{1}{9} \cdot \frac{1}{\left( \frac{2}{3} \right)} \tan^{-1} \frac{x + \frac{1}{3}}{\frac{2}{3}} + c \quad \left( \because \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right)$$

$$= \frac{1}{9} \cdot \frac{3}{2} \tan^{-1} \left( \frac{\frac{3x+1}{2}}{\frac{2}{3}} \right) + c = \frac{1}{6} \tan^{-1} \left( \frac{3x+1}{2} \right) + c.$$

12.  $\frac{1}{\sqrt{7-6x-x^2}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{7-6x-x^2}} dx$  ... (i)  $\left| \begin{array}{l} \text{Type } \int \frac{1}{\text{Quadratic}} dx \end{array} \right.$

Here Quadratic expression is  $7-6x-x^2 = -x^2-6x+7$ .

Making coefficient of  $x^2$  unity,  $= -(x^2+6x-7)$ .

To complete squares, adding and subtracting  $\left( \frac{1}{2} \text{ coefficient of } x \right)^2$

$$= \left( \frac{1}{2} \times 6 \right)^2 = 9$$

$$= -[x^2+6x+9-9-7] = -[(x+3)^2-16] \quad \dots (ii)$$

$$= -(x+3)^2+16 = 4^2-(x+3)^2 \quad \dots (iii)$$

**(Note.** Must adjust negative sign outside Eqn. (ii) in the bracket as shown above because otherwise we shall get  $\sqrt{-1} = i$  on taking square roots.)

Putting the value of quadratic expression from (iii) in (i),

$$I = \int \frac{1}{\sqrt{4^2-(x+3)^2}} dx = \sin^{-1} \left( \frac{x+3}{4} \right) + c$$

$$\left[ \because \int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1} \frac{x}{a} \right]$$

13.  $\frac{1}{\sqrt{(x-1)(x-2)}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{(x-1)(x-2)}} dx = \int \frac{1}{\sqrt{x^2-2x-x+2}} dx$

$$= \int \frac{1}{\sqrt{x^2 - 3x + 2}} \quad \dots(i)$$

Here quadratic expression is  $x^2 - 3x + 2$ . Coefficient of  $x^2$  is already unity. To complete squares, adding and subtracting

$$\begin{aligned} & \left( \frac{1}{2} \text{ coefficient of } x \right)^2, \text{ i.e., } \left( -\frac{3}{2} \right)^2 = \left( \frac{3}{2} \right)^2 \\ x^2 - 3x + 2 &= x^2 - 3x + \left( \frac{3}{2} \right)^2 - \frac{9}{4} + 2 \\ &= \left( x - \frac{3}{2} \right)^2 - \frac{1}{4} \quad \left[ \because \frac{-9}{4} + 2 = \frac{-9 + 8}{4} = \frac{-1}{4} \right] \\ &= \left( x - \frac{3}{2} \right)^2 - \left( \frac{1}{2} \right)^2 \quad \dots(ii) \end{aligned}$$

Putting this value in (i),  $I = \int \frac{1}{\sqrt{\left( x - \frac{3}{2} \right)^2 - \left( \frac{1}{2} \right)^2}} dx$

$$= \log \left| x - \frac{3}{2} + \sqrt{\left( x - \frac{3}{2} \right)^2 - \left( \frac{1}{2} \right)^2} \right| + c$$

$$\left[ \because \int \frac{1}{\sqrt{x^2 - a^2}} dx = \log \left| x + \sqrt{x^2 - a^2} \right| \right]$$

$$= \log \left| x - \frac{3}{2} + \sqrt{x^2 - 3x + 2} \right| + c. \quad [\text{By (ii)}]$$

14.  $\frac{1}{\sqrt{8 + 3x - x^2}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{8 + 3x - x^2}} dx \quad \dots(i)$

Here quadratic expression is  $8 + 3x - x^2 = -x^2 + 3x + 8$ .

Making coefficient of  $x^2$  unity,  $= -(x^2 - 3x - 8)$ .

To complete squares, adding and subtracting

$$\begin{aligned} & \left( \frac{1}{2} \text{ coefficient of } x \right)^2 = \left( \frac{3}{2} \right)^2 \\ 8 + 3x - x^2 &= - \left( x^2 - 3x + \left( \frac{3}{2} \right)^2 - \left( \frac{3}{2} \right)^2 - 8 \right) \\ &= - \left[ \left( x - \frac{3}{2} \right)^2 - \frac{9}{4} - 8 \right] = - \left[ \left( x - \frac{3}{2} \right)^2 - \frac{41}{4} \right] = \frac{41}{4} - \left( x - \frac{3}{2} \right)^2 \\ & \text{(See **Note** given in the solution of Q.N. 12)} \\ &= \left( \frac{\sqrt{41}}{2} \right)^2 - \left( x - \frac{3}{2} \right)^2 \quad \dots(ii) \end{aligned}$$

$$\begin{aligned}
 \text{Putting this value in (i), } I &= \int \frac{1}{\sqrt{\left(\frac{\sqrt{41}}{2}\right)^2 - \left(x - \frac{3}{2}\right)^2}} dx \\
 &= \sin^{-1} \frac{x - \frac{3}{2}}{\frac{\sqrt{41}}{2}} + c \quad \left[ \because \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} \right] \\
 &= \sin^{-1} \left( \frac{2x - 3}{\sqrt{41}} \right) + c.
 \end{aligned}$$

15.  $\frac{1}{\sqrt{(x-a)(x-b)}}$

**Sol.** Let  $I = \int \frac{1}{\sqrt{(x-a)(x-b)}} dx = \int \frac{1}{\sqrt{x^2 - bx - ax + ab}} dx$

$$= \int \frac{1}{\sqrt{x^2 - x(a+b) + ab}} \dots(i)$$

Here Quadratic expression  $= x^2 - x(a+b) + ab$

Adding and subtracting  $\left(\frac{1}{2} \text{ coefficient of } x\right)^2 = \left(\frac{a+b}{2}\right)^2$

$$\begin{aligned}
 &= x^2 - x(a+b) + \left(\frac{a+b}{2}\right)^2 - \left(\frac{a+b}{2}\right)^2 + ab \\
 &= \left(x - \left(\frac{a+b}{2}\right)\right)^2 - \left(\frac{(a+b)^2}{4} - ab\right) \\
 &= \left(x - \left(\frac{a+b}{2}\right)\right)^2 - \left(\frac{(a+b)^2 - 4ab}{4}\right) = \left(x - \left(\frac{a+b}{2}\right)\right)^2 - \frac{(a-b)^2}{4} \\
 &= \left(x - \left(\frac{a+b}{2}\right)\right)^2 - \left(\frac{a-b}{2}\right)^2 \dots(ii)
 \end{aligned}$$

$$\begin{aligned}
 (\because (a+b)^2 - 4ab &= a^2 + b^2 + 2ab - 4ab \\
 &= a^2 + b^2 - 2ab = (a-b)^2)
 \end{aligned}$$

Putting this value in (i),

$$\begin{aligned}
 I &= \int \frac{1}{\sqrt{\left(x - \left(\frac{a+b}{2}\right)\right)^2 - \left(\frac{a-b}{2}\right)^2}} dx \\
 &= \log \left| x - \left(\frac{a+b}{2}\right) + \sqrt{\left(x - \left(\frac{a+b}{2}\right)\right)^2 - \left(\frac{a-b}{2}\right)^2} \right| + c \\
 &\quad \left[ \because \int \frac{1}{\sqrt{x^2 - a^2}} dx = \log |x + \sqrt{x^2 - a^2}| \right]
 \end{aligned}$$

$$= \log \left| x - \left( \frac{a+b}{2} \right) + \sqrt{x^2 - x(a+b) + ab} \right| + c \quad [\text{By (ii)}]$$

**Note.** Method to evaluate  $\int \frac{\text{Linear}}{\text{Quadratic}} dx$  or  $\int \frac{\text{Linear}}{\sqrt{\text{Quadratic}}} dx$

or  $\int \text{Linear} \sqrt{\text{Quadratic}} dx$ .

Write linear = A  $\frac{d}{dx}$  (Quadratic) + B.

Find values of A and B by comparing coefficients of  $x$  and constant terms on both sides.

16.  $\frac{4x+1}{\sqrt{2x^2+x-3}}$

**Sol.** Let  $I = \int \frac{4x+1}{\sqrt{2x^2+x-3}} dx \quad \dots(i)$

Here  $\frac{d}{dx}$  (Quadratic  $2x^2 + x - 3$ ) is  $(4x + 1)$ , the numerator.

So put  $2x^2 + x - 3 = t$ .

$$\therefore (4x + 1) = \frac{dt}{dx} \Rightarrow (4x + 1) dx = dt$$

$$\begin{aligned} \therefore \text{From (i), } I &= \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{\frac{1}{2}} + c \\ &= 2\sqrt{t} + c = 2\sqrt{2x^2+x-3} + c. \end{aligned}$$

17.  $\frac{x+2}{\sqrt{x^2-1}}$

**Sol.** Let  $I = \int \frac{x+2}{\sqrt{x^2-1}} dx = \int \left( \frac{x}{\sqrt{x^2-1}} + \frac{2}{\sqrt{x^2-1}} \right) dx$

$$= \int \frac{x}{\sqrt{x^2-1}} dx + 2 \int \frac{1}{\sqrt{x^2-1^2}} dx$$

$$= \int \frac{x}{\sqrt{x^2-1}} dx + 2 \log |x + \sqrt{x^2-1^2}| + c \quad \dots(i)$$

$$\text{Let } I_1 = \int \frac{x}{\sqrt{x^2-1}} dx = \frac{1}{2} \int \frac{2x}{\sqrt{x^2-1}} dx$$

$$\text{Put } x^2 - 1 = t. \text{ Therefore } 2x = \frac{dt}{dx} \text{ or } 2x dx = dt$$

$$\therefore I_1 = \frac{1}{2} \int \frac{dt}{\sqrt{t}} = \frac{1}{2} \int t^{-1/2} dt = \frac{1}{2} \frac{t^{1/2}}{\frac{1}{2}} = \sqrt{t} = \sqrt{x^2-1}$$

Putting this value of  $(I_1) = \int \frac{x}{\sqrt{x^2-1}} dx = \sqrt{x^2-1}$  in (i)

$$I = \sqrt{x^2-1} + 2 \log |x + \sqrt{x^2-1}| + c.$$

18.  $\frac{5x-2}{1+2x+3x^2}$

**Sol.** Let  $I = \int \frac{5x-2}{1+2x+3x^2} dx$  ... (i)  $\left| \int \frac{\text{Linear}}{\text{Quadratic}} dx \right.$

**Let Linear = A  $\frac{d}{dx}$  (Quadratic) + B**

i.e.,  $5x-2 = A \frac{d}{dx} (1+2x+3x^2) + B$

or  $5x-2 = A(2+6x) + B$

i.e.,  $5x-2 = 2A+6Ax+B$

... (ii)

Comparing coefficients of  $x$ ,  $6A = 5 \Rightarrow A = \frac{5}{6}$

Comparing constants,  $2A+B = -2$

Putting  $A = \frac{5}{6}$ ,  $\frac{10}{6} + B = -2$

$\Rightarrow B = -2 - \frac{10}{6} = \frac{-22}{6}$  or  $B = \frac{-11}{3}$

Putting values of A and B in (ii),  $5x-2 = \frac{5}{6}(2+6x) - \frac{11}{3}$

Putting this value of  $5x-2$  in (i),

$$I = \int \frac{\frac{5}{6}(2+6x) - \frac{11}{3}}{1+2x+3x^2} dx$$

$\Rightarrow I = \frac{5}{6} \int \frac{2+6x}{1+2x+3x^2} dx - \frac{11}{3} \int \frac{1}{1+2x+3x^2} dx$

$$= \frac{5}{6} I_1 - \frac{11}{3} I_2$$

... (iii)

Here  $I_1 = \int \frac{2+6x}{1+2x+3x^2} dx$

Put Denominator  $1+2x+3x^2 = t$ .

$\therefore 2+6x = \frac{dt}{dx} \Rightarrow (2+6x) dx = dt$

$\therefore I_1 = \int \frac{dt}{t} = \int \frac{1}{t} dt = \log |t| = \log |1+2x+3x^2|$  ... (iv)

Again  $I_2 = \int \frac{1}{1+2x+3x^2} dx = \int \frac{1}{3x^2+2x+1} dx \left| \int \frac{1}{\text{Quadratic}} dx \right.$

Now Quadratic Expression =  $3x^2+2x+1$ .

Making coefficient of  $x^2$  unity =  $3 \left( x^2 + \frac{2}{3}x + \frac{1}{3} \right)$

$$\begin{aligned}\text{Completing squares} &= 3 \left[ x^2 + \frac{2}{3}x + \left(\frac{1}{3}\right)^2 + \frac{1}{3} - \frac{1}{9} \right] \\ &= 3 \left[ \left(x + \frac{1}{3}\right)^2 + \frac{2}{9} \right] \quad \left| \because \frac{1}{3} - \frac{1}{9} = \frac{3-1}{9} = \frac{2}{9} \right.\end{aligned}$$

$$\begin{aligned}\Rightarrow I_2 &= \int \frac{1}{3 \left[ \left(x + \frac{1}{3}\right)^2 + \frac{2}{9} \right]} dx = \frac{1}{3} \int \frac{1}{\left(x + \frac{1}{3}\right)^2 + \left(\frac{\sqrt{2}}{3}\right)^2} dx \\ &= \frac{1}{3} \cdot \frac{1}{\left(\frac{\sqrt{2}}{3}\right)} \tan^{-1} \frac{x + \frac{1}{3}}{\frac{\sqrt{2}}{3}} \quad \left[ \because \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right]\end{aligned}$$

$$\Rightarrow I_2 = \frac{1}{3} \cdot \frac{3}{\sqrt{2}} \tan^{-1} \frac{3x+1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \tan^{-1} \left( \frac{3x+1}{\sqrt{2}} \right) \quad \dots(v)$$

Putting values of  $I_1$  and  $I_2$  from (iv) and (v) in (iii), we have

$$I = \frac{5}{6} \log |1 + 2x + 3x^2| - \frac{11}{3} \cdot \frac{1}{\sqrt{2}} \tan^{-1} \left( \frac{3x+1}{\sqrt{2}} \right) + c.$$

**Integrate the functions in Exercises 19 to 23:**

19.  $\frac{6x+7}{\sqrt{(x-5)(x-4)}}$

**Sol.** Let  $I = \int \frac{6x+7}{\sqrt{(x-5)(x-4)}} dx = \int \frac{6x+7}{\sqrt{x^2-4x-5x+20}} dx$

$$\text{i.e., } I = \int \frac{6x+7}{\sqrt{x^2-9x+20}} dx \quad \dots(i) \quad \left| \int \frac{\text{Linear}}{\sqrt{\text{Quadratic}}} dx \right.$$

**Let Linear = A  $\frac{d}{dx}$  (Quadratic) + B**

$$\text{i.e., } 6x+7 = A(2x-9) + B = 2Ax - 9A + B \quad \dots(ii)$$

Comparing coefficients of  $x$ ,  $2A = 6 \Rightarrow A = 3$

Comparing constants,  $-9A + B = 7$ .

Putting  $A = 3$ ,  $-27 + B = 7 \Rightarrow B = 34$

Putting values of  $A$  and  $B$  in (ii),

$$6x+7 = 3(2x-9) + 34$$

Putting this value of  $6x+7$  in (i),

$$\begin{aligned}I &= \int \frac{3(2x-9) + 34}{\sqrt{x^2-9x+20}} dx \\ &= 3 \int \frac{2x-9}{\sqrt{x^2-9x+20}} dx + 34 \int \frac{1}{\sqrt{x^2-9x+20}} dx \\ &= 3 I_1 + 34 I_2 \quad \dots(iii) \\ I_1 &= \int \frac{2x-9}{\sqrt{x^2-9x+20}} dx\end{aligned}$$

$$\text{Put } x^2 - 9x + 20 = t. \quad \therefore 2x - 9 = \frac{dt}{dx}$$

$$\Rightarrow (2x - 9) dx = dt$$

$$\begin{aligned} \therefore I_1 &= \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{1/2} = 2\sqrt{t} \\ &= 2\sqrt{x^2 - 9x + 20} \end{aligned} \quad \dots(iv)$$

$$\begin{aligned} I_2 &= \int \frac{1}{\sqrt{x^2 - 9x + 20}} dx = \int \frac{1}{\sqrt{x^2 - 9x + \left(\frac{9}{2}\right)^2 + 20 - \frac{81}{4}}} dx \\ &= \int \frac{1}{\sqrt{\left(x - \frac{9}{2}\right)^2 - \frac{1}{4}}} dx = \int \frac{1}{\sqrt{\left(x - \frac{9}{2}\right)^2 - \left(\frac{1}{2}\right)^2}} dx \\ &= \log \left| x - \frac{9}{2} + \sqrt{\left(x - \frac{9}{2}\right)^2 - \left(\frac{1}{2}\right)^2} \right| \\ &\quad \left[ \because \int \frac{1}{\sqrt{x^2 - a^2}} dx = \log |x + \sqrt{x^2 - a^2}| \right] \\ I_2 &= \log \left| x - \frac{9}{2} + \sqrt{x^2 - 9x + 20} \right| \end{aligned} \quad \dots(v)$$

$$\left( \because \left(x - \frac{9}{2}\right)^2 - \left(\frac{1}{2}\right)^2 = x^2 + \frac{81}{4} - 9x - \frac{1}{4} = x^2 - 9x + 20 \right)$$

Putting values of  $I_1$  and  $I_2$  from (iv) and (v) in (iii),

$$I = 6\sqrt{x^2 - 9x + 20} + 34 \log \left| x - \frac{9}{2} + \sqrt{x^2 - 9x + 20} \right| + c.$$

20.  $\frac{x+2}{\sqrt{4x-x^2}}$

**Sol.** Let  $I = \int \frac{x+2}{\sqrt{4x-x^2}} dx \quad \dots(i) \quad \left| \int \frac{\text{Linear}}{\sqrt{\text{Quadratic}}} dx \right.$

Let Linear = A  $\frac{d}{dx}$  (Quadratic) + B

i.e.,  $x + 2 = A(4 - 2x) + B$   
 $= 4A - 2Ax + B$  ...(ii)

Comparing coefficients of  $x$ :  $-2A = 1 \Rightarrow A = \frac{-1}{2}$

Comparing constants:  $4A + B = 2$

Putting  $A = \frac{-1}{2}$ ,  $-2 + B = 2 \Rightarrow B = 4$

Putting values of A and B in (ii),  $x + 2 = \frac{-1}{2} (4 - 2x) + 4$

Putting this value of  $x + 2$  in (i),

$$I = \int \frac{-\frac{1}{2}(4-2x) + 4}{\sqrt{4x-x^2}} dx = \frac{-1}{2} \int \frac{4-2x}{\sqrt{4x-x^2}} dx + 4 \int \frac{1}{\sqrt{4x-x^2}} dx$$

$$= \frac{-1}{2} I_1 + 4 I_2 \quad \dots(iii) \quad I_1 = \int \frac{4-2x}{\sqrt{4x-x^2}} dx$$

Put  $4x - x^2 = t \quad \therefore \quad 4 - 2x = \frac{dt}{dx} \Rightarrow (4 - 2x) dx = dt$

$$\therefore I_1 = \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{1/2} = 2\sqrt{t} = 2\sqrt{4x-x^2} \quad \dots(iv)$$

$$I_2 = \int \frac{1}{\sqrt{4x-x^2}} dx$$

Quadratic Expression is  $4x - x^2 = -x^2 + 4x$   
 $= -(x^2 - 4x) = -(x^2 - 4x + 4 - 4) = -((x-2)^2 - 2^2) = 2^2 - (x-2)^2$

$$\therefore I_2 = \int \frac{1}{\sqrt{2^2 - (x-2)^2}} dx = \sin^{-1} \frac{x-2}{2} \quad \dots(v)$$

$$\left( \because \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} \right)$$

Putting values of  $I_1$  and  $I_2$  from (iv) and (v) in (iii),

$$I = -\sqrt{4x-x^2} + 4 \sin^{-1} \frac{x-2}{2} + c.$$

21.  $\frac{x+2}{\sqrt{x^2+2x+3}}$

Sol. Let  $I = \int \frac{x+2}{\sqrt{x^2+2x+3}} dx \quad \dots(i)$

Let **Linear** = A  $\frac{d}{dx}$  (**Quadratic**) + B

i.e.,  $x+2 = A(2x+2) + B$   
 $= 2Ax + 2A + B \quad \dots(ii)$

Comparing coefficients of  $x$ ,  $2A = 1 \Rightarrow A = \frac{1}{2}$

Comparing constants,  $2A + B = 2$

Putting  $A = \frac{1}{2}$ ,  $1 + B = 2 \Rightarrow B = 1$

Putting values of A and B in (ii),  $x+2 = \frac{1}{2}(2x+2) + 1$

Putting this value of  $(x+2)$  in (i),

$$I = \int \frac{\frac{1}{2}(2x+2) + 1}{\sqrt{x^2+2x+3}} dx = \frac{1}{2} \int \frac{2x+2}{\sqrt{x^2+2x+3}} dx + \int \frac{dx}{\sqrt{x^2+2x+3}}$$

$$\Rightarrow I = \frac{1}{2} I_1 + I_2 \quad \dots(iii) \quad I_1 = \int \frac{2x+2}{\sqrt{x^2+2x+3}} dx$$

$$\text{Put } x^2 + 2x + 3 = t \quad \therefore (2x+2) = \frac{dt}{dx} \Rightarrow (2x+2) dx = dt$$

$$I_1 = \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{\frac{1}{2}} = 2\sqrt{t} = 2\sqrt{x^2+2x+3} \quad \dots(iv)$$

$$\begin{aligned} I_2 &= \int \frac{1}{\sqrt{x^2+2x+3}} dx = \int \frac{1}{\sqrt{x^2+2x+1+2}} dx \\ &= \int \frac{1}{\sqrt{(x+1)^2 + (\sqrt{2})^2}} dx = \log |x+1 + \sqrt{(x+1)^2 + (\sqrt{2})^2}| \\ &\quad \left[ \because \int \frac{1}{\sqrt{x^2+a^2}} dx = \log |x + \sqrt{x^2+a^2}| \right] \\ &= \log |x+1 + \sqrt{x^2+2x+3}| \quad \dots(v) \end{aligned}$$

Putting values from (iv) and (v) in (iii),

$$I = \sqrt{x^2+2x+3} + \log |x+1 + \sqrt{x^2+2x+3}| + c.$$

**22.**  $\frac{x+3}{x^2-2x-5}$

**Sol.** Let  $I = \int \frac{x+3}{x^2-2x-5} dx \quad \dots(i)$

$$\begin{aligned} \text{Let } x+3 &= A \frac{d}{dx} (x^2-2x-5) + B \\ \text{or } x+3 &= A(2x-2) + B \quad \dots(ii) \\ &= 2Ax - 2A + B \end{aligned}$$

Comparing coefficients of  $x$  on both sides,  $2A = 1 \Rightarrow A = \frac{1}{2}$

Comparing constants,  $-2A + B = 3$

Putting  $A = \frac{1}{2}$ ,  $-1 + B = 3 \Rightarrow B = 4$

Putting values of  $A$  and  $B$  in (ii),  $x+3 = \frac{1}{2} (2x-2) + 4$

Putting this value in (i),

$$\begin{aligned} I &= \int \frac{\frac{1}{2}(2x-2) + 4}{x^2-2x-5} dx = \frac{1}{2} \int \frac{2x-2}{x^2-2x-5} dx + 4 \int \frac{1}{x^2-2x-5} dx \\ &= \frac{1}{2} I_1 + 4 I_2 \quad \dots(iii) \end{aligned}$$

$$I_1 = \int \frac{2x-2}{x^2-2x-5} dx$$

Put  $x^2 - 2x - 5 = t$ . Therefore  $(2x-2) = \frac{dt}{dx} \Rightarrow (2x-2) dx = dt$

$$\therefore I_1 = \int \frac{dt}{t} = \log |t| = \log |x^2 - 2x - 5| \quad \dots(iv)$$

$$\begin{aligned} \text{Again } I_2 &= \int \frac{1}{x^2 - 2x - 5} dx \\ &= \int \frac{1}{x^2 - 2x + 1 - 1 - 5} dx = \int \frac{1}{(x-1)^2 - 6} dx \\ &= \int \frac{1}{(x-1)^2 - 6} dx = \frac{1}{2\sqrt{6}} \log \left| \frac{x-1-\sqrt{6}}{x-1+\sqrt{6}} \right| \quad \dots(v) \\ &\quad \left[ \because \int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| \right] \end{aligned}$$

Putting values of  $I_1$  and  $I_2$  from (iv) and (v) in (iii),

$$I = \frac{1}{2} \log |x^2 - 2x - 5| + \frac{2}{\sqrt{6}} \log \left| \frac{x-1-\sqrt{6}}{x-1+\sqrt{6}} \right| + c.$$

23.  $\frac{5x+3}{\sqrt{x^2+4x+10}}$

**Sol.** Let  $I = \int \frac{5x+3}{\sqrt{x^2+4x+10}} dx \quad \dots(i)$

**Let Linear = A  $\frac{d}{dx}$  (Quadratic) + B**

$$\text{i.e., } 5x+3 = A(2x+4) + B = 2Ax + 4A + B \quad \dots(ii)$$

Comparing coefficients of  $x$  on both sides,  $2A = 5 \Rightarrow A = \frac{5}{2}$

Comparing constants,  $4A + B = 3$

Putting  $A = \frac{5}{2}$ ,  $10 + B = 3 \Rightarrow B = -7$

Putting values of  $A$  and  $B$  in (ii),  $5x+3 = \frac{5}{2}(2x+4) - 7$

$$\begin{aligned} \text{Putting this value in (i), } I &= \int \frac{\frac{5}{2}(2x+4) - 7}{\sqrt{x^2+4x+10}} dx \\ &= \frac{5}{2} \int \frac{2x+4}{\sqrt{x^2+4x+10}} dx - 7 \int \frac{1}{\sqrt{x^2+4x+10}} dx \end{aligned}$$

or  $I = \frac{5}{2} I_1 - 7 I_2 \quad \dots(iii)$

$$I_1 = \int \frac{2x+4}{\sqrt{x^2+4x+10}} dx$$

Put  $x^2 + 4x + 10 = t$ . Therefore  $2x+4 = \frac{dt}{dx} \Rightarrow (2x+4) dx = dt$

$$\begin{aligned}
 \therefore I_1 &= \int \frac{dt}{\sqrt{t}} = \int t^{-1/2} dt = \frac{t^{1/2}}{\frac{1}{2}} = 2\sqrt{t} \\
 &= 2\sqrt{x^2 + 4x + 10} \quad \dots(iv) \\
 I_2 &= \int \frac{1}{\sqrt{x^2 + 4x + 10}} dx = \int \frac{1}{\sqrt{x^2 + 4x + 4 + 6}} dx \\
 &= \int \frac{1}{\sqrt{(x+2)^2 + (\sqrt{6})^2}} dx = \log |x + 2 + \sqrt{(x+2)^2 + (\sqrt{6})^2}| \\
 &\quad \left[ \because \int \frac{1}{\sqrt{x^2 + a^2}} dx = \log |x + \sqrt{x^2 + a^2}| \right] \\
 &= \log |x + 2 + \sqrt{x^2 + 4x + 10}| \quad \dots(v)
 \end{aligned}$$

Putting values of  $I_1$  and  $I_2$  from (iv) and (v) in (iii),

$$I = 5\sqrt{x^2 + 4x + 10} - 7 \log |x + 2 + \sqrt{x^2 + 4x + 10}| + c.$$

**Choose the correct answer in Exercises 24 and 25.**

24.  $\int \frac{dx}{x^2 + 2x + 2}$  equals
- (A)  $x \tan^{-1}(x + 1) + C$  (B)  $\tan^{-1}(x + 1) + C$   
 (C)  $(x + 1) \tan^{-1} x + C$  (D)  $\tan^{-1} x + C$ .

**Sol.**  $\int \frac{dx}{x^2 + 2x + 2} = \int \frac{1}{x^2 + 2x + 1 + 1} dx = \int \frac{1}{(x + 1)^2 + 1^2} dx$

$$\begin{aligned}
 &= \frac{1}{1} \tan^{-1} \frac{(x + 1)}{1} + C \left[ \because \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right] \\
 &= \tan^{-1}(x + 1) + C
 \end{aligned}$$

$\therefore$  Option (B) is the correct answer.

25.  $\int \frac{dx}{\sqrt{9x - 4x^2}}$  equals
- (A)  $\frac{1}{9} \sin^{-1} \left( \frac{9x - 8}{8} \right) + C$  (B)  $\frac{1}{2} \sin^{-1} \left( \frac{8x - 9}{9} \right) + C$   
 (C)  $\frac{1}{3} \sin^{-1} \left( \frac{9x - 8}{8} \right) + C$  (D)  $\frac{1}{2} \sin^{-1} \left( \frac{9x - 8}{8} \right) + C$ .

**Sol.** Let  $I = \int \frac{dx}{\sqrt{9x - 4x^2}} = \int \frac{dx}{\sqrt{-4x^2 + 9x}} \quad \dots(i)$

Here Quadratic expression is  $-4x^2 + 9x = -4 \left( x^2 - \frac{9}{4}x \right)$

$$\begin{aligned}
 &= -4 \left[ x^2 - \frac{9}{4}x + \left( \frac{9}{8} \right)^2 - \left( \frac{9}{8} \right)^2 \right] = -4 \left[ \left( x - \frac{9}{8} \right)^2 - \left( \frac{9}{8} \right)^2 \right] \\
 &= 4 \left[ - \left( x - \frac{9}{8} \right)^2 + \left( \frac{9}{8} \right)^2 \right] = 4 \left[ \left( \frac{9}{8} \right)^2 - \left( x - \frac{9}{8} \right)^2 \right]
 \end{aligned}$$

Putting this value in (i),

$$I = \int \frac{1}{\sqrt{4 \left[ \left( \frac{9}{8} \right)^2 - \left( x - \frac{9}{8} \right)^2 \right]}} dx = \frac{1}{2} \int \frac{1}{\sqrt{\left[ \left( \frac{9}{8} \right)^2 - \left( x - \frac{9}{8} \right)^2 \right]}} dx$$

$$= \frac{1}{2} \sin^{-1} \frac{x - \frac{9}{8}}{\frac{9}{8}} + C$$

$$= \frac{1}{2} \sin^{-1} \left( \frac{8x - 9}{9} \right) + C \quad \left[ \because \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} \right]$$

$\therefore$  Option (B) is the correct answer.



## Exercise 7.5

**Integrate the (rational) functions in Exercises 1 to 6:**

1.  $\frac{x}{(x+1)(x+2)}$

**Sol.** To integrate the (rational) function  $\frac{x}{(x+1)(x+2)}$ .

Let integrand  $\frac{x}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2}$  ... (i)

(Partial Fractions)

Multiplying by L.C.M. =  $(x+1)(x+2)$ ,

$$x = A(x+2) + B(x+1) = Ax + 2A + Bx + B$$

Comparing coefficients of  $x$  on both sides,  $A + B = 1$  ... (ii)

Comparing constants,  $2A + B = 0$  ... (iii)

Let us solve Eqns. (ii) and (iii) for  $A$  and  $B$ .

Eqn. (iii) – Eqn. (ii) gives,  $A = -1$

Putting  $A = -1$  in (ii),  $-1 + B = 1 \Rightarrow B = 2$

Putting values of  $A$  and  $B$  in (i),  $\frac{x}{(x+1)(x+2)} = \frac{-1}{x+1} + \frac{2}{x+2}$

$$\begin{aligned} \therefore \int \frac{x}{(x+1)(x+2)} dx &= - \int \frac{1}{x+1} dx + 2 \int \frac{1}{x+2} dx \\ &= - \log |x+1| + 2 \log |x+2| + c \\ &= \log |x+2|^2 - \log |x+1| + c = \log \frac{(x+2)^2}{|x+1|} + c. \end{aligned}$$

( $\because |t|^2 = t^2$ )

2.  $\frac{1}{x^2-9}$

**Sol.** To integrate the (rational) function  $\frac{1}{x^2-9}$

$$\int \frac{1}{x^2-9} dx = \int \frac{1}{x^2-3^2} dx$$

$$\begin{aligned}
 &= \frac{1}{2 \times 3} \log \left| \frac{x-3}{x+3} \right| + c \left[ \because \int \frac{1}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| \right] \\
 &= \frac{1}{6} \log \left| \frac{x-3}{x+3} \right| + c.
 \end{aligned}$$

**OR**

$$\text{Integrand } \frac{1}{x^2 - 9} = \frac{1}{(x-3)(x+3)} = \frac{A}{x-3} + \frac{B}{x+3}$$

Now proceed as in the solution of Q.No.1.

3.  $\frac{3x-1}{(x-1)(x-2)(x-3)}$

**Sol.** To integrate the (rational) function  $\frac{3x-1}{(x-1)(x-2)(x-3)}$

$$\text{Let integrand } \frac{3x-1}{(x-1)(x-2)(x-3)} = \frac{A}{x-1} + \frac{B}{x-2} + \frac{C}{x-3} \quad \dots(i)$$

Multiplying by L.C.M. =  $(x-1)(x-2)(x-3)$ , we have

$$\begin{aligned}
 3x-1 &= A(x-2)(x-3) + B(x-1)(x-3) + C(x-1)(x-2) \\
 &= A(x^2 - 5x + 6) + B(x^2 - 4x + 3) + C(x^2 - 3x + 2) \\
 &= Ax^2 - 5Ax + 6A + Bx^2 - 4Bx + 3B + Cx^2 - 3Cx + 2C
 \end{aligned}$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides, we have

$$\text{Coefficients of } x^2: A + B + C = 0 \quad \dots(ii)$$

$$\text{Coefficient of } x: -5A - 4B - 3C = 3 \text{ or } 5A + 4B + 3C = -3 \quad \dots(iii)$$

$$\text{Constants: } 6A + 3B + 2C = -1 \quad \dots(iv)$$

Let us solve (ii), (iii) and (iv) for A, B, C.

Let us first form two Eqns. in two unknowns say A and B.

Eqn. (iii) - 3 Eqn. (i) gives (to eliminate C),

$$5A + 4B + 3C - 3A - 3B - 3C = -3$$

$$\text{or } 2A + B = -3 \quad \dots(v)$$

Eqn. (iv) - 2 Eqn. (i) gives (to eliminate C),

$$6A + 3B + 2C - 2A - 2B - 2C = -1$$

$$\text{or } 4A + B = -1 \quad \dots(vi)$$

Eqn. (vi) - Eqn. (v) gives (to eliminate B),

$$2A = -1 + 3 = 2 \Rightarrow A = \frac{2}{2} = 1.$$

$$\text{Putting } A = 1 \text{ in (v), } 2 + B = -3 \Rightarrow B = -5$$

$$\text{Putting } A = 1 \text{ and } B = -5 \text{ in (ii), } 1 - 5 + C = 0$$

$$\text{or } C - 4 = 0 \text{ or } C = 4$$

Putting values of A, B, C in (i),

$$\frac{3x-1}{(x-1)(x-2)(x-3)} = \frac{1}{x-1} - \frac{5}{x-2} + \frac{4}{x-3}$$

$$\begin{aligned}
 \therefore \int \frac{3x-1}{(x-1)(x-2)(x-3)} \\
 &= \int \frac{1}{x-1} dx - 5 \int \frac{1}{x-2} dx + 4 \int \frac{1}{x-3} dx \\
 &= \log |x-1| - 5 \log |x-2| + 4 \log |x-3| + c.
 \end{aligned}$$

4.  $\frac{x}{(x-1)(x-2)(x-3)}$

**Sol.** To integrate the (rational) function  $\frac{x}{(x-1)(x-2)(x-3)}$ .

$$\text{Let integrand } \frac{x}{(x-1)(x-2)(x-3)} = \frac{A}{x-1} + \frac{B}{x-2} + \frac{C}{x-3} \quad \dots(i)$$

(Partial fractions)

Multiplying by L.C.M. =  $(x-1)(x-2)(x-3)$ ,

$$\begin{aligned}
 x &= A(x-2)(x-3) + B(x-1)(x-3) + C(x-1)(x-2) \\
 &= A(x^2 - 5x + 6) + B(x^2 - 4x + 3) + C(x^2 - 3x + 2) \\
 &= Ax^2 - 5Ax + 6A + Bx^2 - 4Bx + 3B + Cx^2 - 3Cx + 2C
 \end{aligned}$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides, we have

$$x^2: \quad A + B + C = 0 \quad \dots(ii)$$

$$x: \quad -5A - 4B - 3C = 1 \quad \text{or} \quad 5A + 4B + 3C = -1 \quad \dots(iii)$$

$$\text{Constants: } 6A + 3B + 2C = 0 \quad \dots(iv)$$

Let us solve Eqns. (ii), (iii) and (iv) for A, B, C.

Let us first form two Eqns. in two unknowns say A and B.

Eqn. (iii) - 3 × Eqn. (ii) gives | To eliminate C

$$5A + 4B + 3C - 3A - 3B - 3C = -1 \quad \text{or} \quad 2A + B = -1 \quad \dots(v)$$

Eqn. (iv) - 2 × Eqn. (ii) gives | To eliminate C

$$4A + B = 0 \quad \dots(vi)$$

Eqn. (vi) - Eqn. (v) gives (To eliminate B)

$$2A = 1 \quad \therefore \quad A = \frac{1}{2}$$

$$\text{Putting } A = \frac{1}{2} \text{ in (v), } 1 + B = -1 \Rightarrow B = -2$$

$$\text{Putting } A = \frac{1}{2} \text{ and } B = -2 \text{ in (ii),}$$

$$\frac{1}{2} - 2 + C = 0 \Rightarrow C = \frac{-1}{2} + 2 = \frac{-1+4}{2} = \frac{3}{2}$$

Putting these values of A, B, C in (i), we have

$$\begin{aligned}
 \frac{x}{(x-1)(x-2)(x-3)} &= \frac{\frac{1}{2}}{x-1} - \frac{2}{x-2} + \frac{\frac{3}{2}}{x-3} \\
 \therefore \int \frac{x}{(x-1)(x-2)(x-3)} dx \\
 &= \frac{1}{2} \int \frac{1}{x-1} dx - 2 \int \frac{1}{x-2} dx + \frac{3}{2} \int \frac{1}{x-3} dx \\
 &= \frac{1}{2} \log |x-1| - 2 \log |x-2| + \frac{3}{2} \log |x-3| + c.
 \end{aligned}$$

5.  $\frac{2x}{x^2 + 3x + 2}$

**Sol.** To integrate the (rational) function  $\frac{2x}{x^2 + 3x + 2}$ .

$$\text{Now } x^2 + 3x + 2 = x^2 + 2x + x + 2 = x(x + 2) + 1(x + 2) \\ = (x + 1)(x + 2)$$

$$\therefore \text{Integrand } \frac{2x}{x^2 + 3x + 2} = \frac{2x}{(x + 1)(x + 2)} \\ = \frac{A}{x + 1} + \frac{B}{x + 2} \quad \dots(i)$$

(Partial Fractions)

Multiplying both sides by L.C.M. =  $(x + 1)(x + 2)$ ,

$$2x = A(x + 2) + B(x + 1) = Ax + 2A + Bx + B$$

Comparing coefficients of  $x$  and constant terms on both sides, we have

$$\text{Coefficients of } x: A + B = 2 \quad \dots(ii)$$

$$\text{Constant terms: } 2A + B = 0 \quad \dots(iii)$$

Let us solve (ii) and (iii) for  $A$  and  $B$ .

$$(iii) - (ii) \text{ gives } A = -2.$$

$$\text{Putting } A = -2 \text{ in (ii), } -2 + B = 2. \quad \therefore B = 4$$

$$\text{Putting values of } A \text{ and } B \text{ in (i), } \frac{2x}{x^2 + 3x + 2} = \frac{-2}{x + 1} + \frac{4}{x + 2}$$

$$\therefore \int \frac{2x}{x^2 + 3x + 2} dx = -2 \int \frac{1}{x + 1} dx + 4 \int \frac{1}{x + 2} dx \\ = -2 \log |x + 1| + 4 \log |x + 2| + c \\ = 4 \log |x + 2| - 2 \log |x + 1| + c$$

**Remark:** Alternative method to evaluate  $\int \frac{2x}{x^2 + 3x + 2} dx$

is  $\int \frac{\text{Linear}}{\text{Quadratic}} dx$  as explained in solutions in Exercise 7.4

(Exercise 18 and Exercise 22.)

6.  $\frac{1 - x^2}{x(1 - 2x)}$

**Sol.** To integrate (rational) function  $\frac{1 - x^2}{x(1 - 2x)} = \frac{1 - x^2}{x - 2x^2} = \frac{-x^2 + 1}{-2x^2 + x}$

[Here Degree of numerator = Degree of Denominator = 2

$\therefore$  We must divide numerator by denominator to make the degree of numerator smaller than degree of denominator so that we can form partial fractions.]

$$\begin{array}{r}
 -2x^2 + x \quad ) \quad -x^2 + 1 \quad \left( \frac{1}{2} \right. \\
 \underline{-x^2 + \frac{x}{2}} \\
 \phantom{-2x^2 + x} + \phantom{-x^2 + \frac{x}{2}} - \\
 \phantom{-2x^2 + x} - \frac{x}{2} + 1
 \end{array}$$

$$\therefore \frac{1-x^2}{x(1-2x)} = \text{Quotient} + \frac{\text{Remainder}}{\text{Divisor}} = \frac{1}{2} + \frac{\left(-\frac{x}{2}+1\right)}{x(1-2x)}$$

$$\begin{aligned}
 \therefore \int \frac{1-x^2}{x(1-2x)} dx &= \int \left( \frac{1}{2} + \frac{\left(-\frac{x}{2}+1\right)}{x(1-2x)} \right) dx \\
 &= \frac{1}{2} \int 1 dx + \int \frac{-\frac{x}{2}+1}{x(1-2x)} dx \quad \dots(i)
 \end{aligned}$$

$$\text{Let integrand } \frac{-\frac{x}{2}+1}{x(1-2x)} = \frac{A}{x} + \frac{B}{1-2x} \quad \dots(ii)$$

Multiplying by L.C.M. =  $x(1-2x)$ ,

$$-\frac{x}{2} + 1 = A(1-2x) + Bx = A - 2Ax + Bx$$

$$\text{Comparing coefficients of } x, \quad -2A + B = \frac{-1}{2} \quad \dots(iii)$$

$$\text{Comparing constants, } A = 1 \quad \dots(iv)$$

Putting  $A = 1$  from (iv) in (iii),

$$-2 + B = \frac{-1}{2} \Rightarrow B = \frac{-1}{2} + 2 = \frac{-1+4}{2} \quad \text{or} \quad B = \frac{3}{2}$$

Putting values of A and B in (ii),

$$\frac{-\frac{x}{2}+1}{x(1-2x)} = \frac{1}{x} + \frac{\frac{3}{2}}{1-2x}$$

$$\begin{aligned}
 \therefore \int \frac{-\frac{x}{2}+1}{x(1-2x)} dx &= \int \frac{1}{x} dx + \frac{3}{2} \int \frac{1}{1-2x} dx \\
 &= \log |x| + \frac{3}{2} \log \frac{|1-2x|}{-2 \rightarrow \text{Coefficient of } x} + c \\
 &= \log |x| - \frac{3}{4} \log |1-2x| + c
 \end{aligned}$$

Putting this value in (i),

$$\int \frac{1-x^2}{x(1-2x)} dx = \frac{1}{2} x + \log |x| - \frac{3}{4} \log |1-2x| + c.$$

**Integrate the following functions in Exercises 7 to 12:**

7.  $\frac{x}{(x^2 + 1)(x - 1)}$

**Sol.** To integrate the (rational) function  $\frac{x}{(x^2 + 1)(x - 1)}$ .

Let integrand  $\frac{x}{(x^2 + 1)(x - 1)} = \frac{Ax + B}{x^2 + 1} + \frac{C}{x - 1}$  ... (i)

(Partial Fractions)

Multiplying by L.C.M. =  $(x^2 + 1)(x - 1)$  on both sides,

$$x = (Ax + B)(x - 1) + C(x^2 + 1)$$

$$\Rightarrow x = Ax^2 - Ax + Bx - B + Cx^2 + C,$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides, we have

$$x^2 \quad A + C = 0 \quad \dots (ii)$$

$$x \quad -A + B = 1 \quad \dots (iii)$$

$$\text{Constants} \quad -B + C = 0 \quad \dots (iv)$$

Let us solve Eqns. (ii), (iii) and (iv) for A, B, C

Adding (ii) and (iii) to eliminate A,  $B + C = 1$  ... (v)

Adding (iv) and (v),  $2C = 1 \Rightarrow C = \frac{1}{2}$

From (iv),  $-B = -C \Rightarrow B = C = \frac{1}{2}$

From (ii),  $A = -C = -\frac{1}{2}$

Putting these values of A, B, C in (i),

$$\begin{aligned} \frac{x}{(x^2 + 1)(x - 1)} &= \frac{-\frac{1}{2}x + \frac{1}{2}}{x^2 + 1} + \frac{\frac{1}{2}}{x - 1} \\ &= \frac{-1}{2} \cdot \frac{x}{x^2 + 1} + \frac{1}{2} \cdot \frac{1}{x^2 + 1} + \frac{1}{2} \cdot \frac{1}{x - 1} \\ &= \frac{-1}{4} \cdot \frac{2x}{x^2 + 1} + \frac{1}{2} \cdot \frac{1}{x^2 + 1} + \frac{1}{2} \cdot \frac{1}{x - 1} \end{aligned}$$

$$\therefore \int \frac{x}{(x^2 + 1)(x - 1)} dx$$

$$= \frac{-1}{4} \int \frac{2x}{x^2 + 1} dx + \frac{1}{2} \int \frac{1}{x^2 + 1} dx + \frac{1}{2} \int \frac{1}{x - 1} dx$$

$$\Rightarrow \int \frac{x}{(x^2 + 1)(x - 1)} dx = \frac{-1}{4} \log |x^2 + 1| + \frac{1}{2} \tan^{-1} x$$

$$+ \frac{1}{2} \log |x - 1| + c \quad \left( \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right)$$

$$= \frac{-1}{4} \log (x^2 + 1) + \frac{1}{2} \tan^{-1} x + \frac{1}{2} \log |x - 1| + c$$

$[\because x^2 + 1 > 0 \Rightarrow |x^2 + 1| = x^2 + 1]$

$$= \frac{1}{2} \log |x - 1| - \frac{1}{4} \log (x^2 + 1) + \frac{1}{2} \tan^{-1} x + c.$$

8.  $\frac{x}{(x-1)^2(x+2)}$

**Sol.** To integrate the (rational) function  $\frac{x}{(x-1)^2(x+2)}$ .

Let integrand  $\frac{x}{(x-1)^2(x+2)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+2} \quad \dots(i)$

(Partial fractions)

Multiplying both sides of (i) by L.C.M. =  $(x-1)^2(x+2)$

$$x = A(x-1)(x+2) + B(x+2) + C(x-1)^2$$

or  $x = A(x^2 + 2x - x - 2) + B(x+2) + C(x^2 + 1 - 2x)$

or  $x = Ax^2 + Ax - 2A + Bx + 2B + Cx^2 + C - 2Cx$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides

$x^2$   $A + C = 0 \quad \dots(ii)$

$x$   $A + B - 2C = 1 \quad \dots(iii)$

**Constants**  $-2A + 2B + C = 0 \quad \dots(iv)$

Let us solve (ii), (iii) and (iv) for A, B, C

From (ii),  $A = -C$

Putting  $A = -C$  in (iv),  $2C + 2B + C = 0$

$$\Rightarrow 2B = -3C \Rightarrow B = \frac{-3C}{2}$$

Putting values of A and B in (iii),

$$-C - \frac{-3C}{2} - 2C = 1 \Rightarrow -2C - 3C - 4C = 2$$

$$\Rightarrow -9C = 2 \Rightarrow C = \frac{-2}{9}$$

Putting  $C = \frac{-2}{9}$ ,  $B = \frac{-3C}{2} = \frac{-3}{2} \left( \frac{-2}{9} \right) = \frac{1}{3} \therefore A = -C = \frac{2}{9}$

Putting these values of A, B, C in (i),

$$\frac{x}{(x-1)^2(x+2)} = \frac{\frac{2}{9}}{x-1} + \frac{\frac{1}{3}}{(x-1)^2} - \frac{\frac{2}{9}}{x+2}$$

$$\begin{aligned} \therefore \int \frac{x}{(x-1)^2(x+2)} dx &= \frac{2}{9} \int \frac{1}{x-1} dx + \frac{1}{3} \int (x-1)^{-2} dx - \frac{2}{9} \int \frac{1}{x+2} dx \\ &= \frac{2}{9} \log |x-1| + \frac{1}{3} \frac{(x-1)^{-1}}{(-1)(1)} - \frac{2}{9} \log |x+2| + c \\ &= \frac{2}{9} (\log |x-1| - \log |x+2|) - \frac{1}{3(x-1)} + c \\ &= \frac{2}{9} \log \left| \frac{x-1}{x+2} \right| - \frac{1}{3(x-1)} + c. \end{aligned}$$

9.  $\frac{3x+5}{x^3-x^2-x+1}$

**Sol.** To integrate the (rational) function  $\frac{3x+5}{x^3-x^2-x+1}$ .

$$\begin{aligned}\text{Now denominator} &= x^3 - x^2 - x + 1 \\ &= x^2(x-1) - 1(x-1) = (x-1)(x^2-1) \\ &= (x-1)(x-1)(x+1) = (x-1)^2(x+1)\end{aligned}$$

$$\therefore \text{Integrand } \frac{3x+5}{x^3-x^2-x+1} = \frac{3x+5}{(x-1)^2(x+1)}$$

$$= \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1} \quad \dots(i) \quad (\text{Partial fractions})$$

$$\begin{aligned}\text{Multiplying by L.C.M.} &= (x-1)^2(x+1), \\ 3x+5 &= A(x-1)(x+1) + B(x-1) + C(x-1)^2 \\ &= A(x^2-1) + B(x-1) + C(x^2+1-2x) \\ &= Ax^2 - A + Bx + B + Cx^2 + C - 2Cx\end{aligned}$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides,

$$x^2 \quad A + C = 0 \quad \dots(ii)$$

$$x \quad B - 2C = 3 \quad \dots(iii)$$

$$\text{Constants} \quad -A + B + C = 5 \quad \dots(iv)$$

Let us solve Eqns. (ii), (iii) and (iv) for A, B, C.

From (ii),  $A = -C$  and from (iii),  $B = 2C + 3$

Putting these values of A and B in (iv),

$$C + 2C + 3 + C = 5 \quad \Rightarrow \quad 4C = 2 \quad \Rightarrow \quad C = \frac{2}{4} = \frac{1}{2}$$

$$\therefore \quad A = -C = -\frac{1}{2}$$

$$\text{and} \quad B = 2C + 3 = 2\left(\frac{1}{2}\right) + 3 = 1 + 3 = 4.$$

Putting these values of A, B, C in (i)

$$\begin{aligned}\frac{3x+5}{x^3-x^2-x+1} &= \frac{-\frac{1}{2}}{x-1} + \frac{4}{(x-1)^2} + \frac{\frac{1}{2}}{x+1} \\ \therefore \int \frac{3x+5}{x^3-x^2-x+1} dx &= \frac{-1}{2} \int \frac{1}{x-1} dx + 4 \int (x-1)^{-2} dx + \frac{1}{2} \int \frac{1}{x+1} dx \\ &= \frac{-1}{2} \log |x-1| + 4 \frac{(x-1)^{-1}}{(-1)(1)} + \frac{1}{2} \log |x+1| + c \\ &\quad \downarrow \\ &\quad \text{Coeff. of } x \\ &= \frac{1}{2} (\log |x+1| - \log |x-1|) - \frac{4}{x-1} + c \\ &= \frac{1}{2} \log \left| \frac{x+1}{x-1} \right| - \frac{4}{x-1} + c.\end{aligned}$$

$$10. \quad \frac{2x-3}{(x^2-1)(2x+3)}$$

**Sol.** To integrate the rational function  $\frac{2x-3}{(x^2-1)(2x+3)}$ .

$$\begin{aligned}\text{Let integrand } \frac{2x-3}{(x^2-1)(2x+3)} &= \frac{2x-3}{(x-1)(x+1)(2x+3)} \\ &= \frac{A}{x-1} + \frac{B}{x+1} + \frac{C}{2x+3} \quad \dots(i)\end{aligned}$$

Multiplying both sides by L.C.M. =  $(x-1)(x+1)(2x+3)$ ,  
 $2x-3 = A(x+1)(2x+3) + B(x-1)(2x+3) + C(x-1)(x+1)$   
 or  $2x-3 = A(2x^2 + 3x + 2x + 3) + B(2x^2 + 3x - 2x - 3) + C(x^2 - 1)$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides,  
 $x^2 \quad 2A + 2B + C = 0 \quad \dots(ii)$

$x \quad 5A + B = 2 \quad \dots(iii)$

**Constants**  $3A - 3B - C = -3 \quad \dots(iv)$

Let us solve Eqns. (ii), (iii) and (iv) for A, B, C.

Eqn. (ii) + Eqn. (iv) gives (to eliminate C)  
 $5A - B = -3 \quad \dots(v)$

Adding Eqns. (iii) and (v),  $10A = -1 \Rightarrow A = \frac{-1}{10}$

Putting  $A = \frac{-1}{10}$  in (iii),  $\frac{-5}{10} + B = 2 \Rightarrow B = 2 + \frac{1}{2} = \frac{5}{2}$

Putting values of A and B in (ii),

$$\frac{-1}{5} + 5 + C = 0 \quad \therefore C = \frac{1}{5} - 5 = \frac{1-25}{5} = \frac{-24}{5}$$

Putting values of A, B, C in (i),

$$\begin{aligned}\frac{2x-3}{(x^2-1)(2x+3)} &= \frac{\frac{-1}{10}}{x-1} + \frac{\frac{5}{2}}{x+1} - \frac{\frac{24}{5}}{2x+3} \\ \therefore \int \frac{2x-3}{(x^2-1)(2x+3)} dx &= \frac{-1}{10} \int \frac{1}{x-1} dx + \frac{5}{2} \int \frac{1}{x+1} dx - \frac{24}{5} \int \frac{1}{2x+3} dx \\ &= \frac{-1}{10} \frac{\log |x-1|}{1 \rightarrow \text{Coeff. of } x} + \frac{5}{2} \frac{\log |x+1|}{1} - \frac{24}{5} \frac{\log |2x+3|}{2 \rightarrow \text{Coeff. of } x} + c \\ &= \frac{-1}{10} \log |x-1| + \frac{5}{2} \log |x+1| - \frac{12}{5} \log |2x+3| + c \\ &= \frac{5}{2} \log |x+1| - \frac{1}{10} \log |x-1| - \frac{12}{5} \log |2x+3| + c.\end{aligned}$$

11.  $\frac{5x}{(x+1)(x^2-4)}$

**Sol.** To integrate the rational function  $\frac{5x}{(x+1)(x^2-4)}$ .

$$\begin{aligned}\text{Let integrand } \frac{5x}{(x+1)(x^2-4)} &= \frac{5x}{(x+1)(x+2)(x-2)} \\ &= \frac{A}{x+1} + \frac{B}{x+2} + \frac{C}{x-2} \quad \dots(i) \text{ (Partial fractions)}\end{aligned}$$

Multiplying both sides of (i) by L.C.M.

$$\begin{aligned} &= (x+1)(x+2)(x-2), \\ 5x &= A(x+2)(x-2) + B(x+1)(x-2) + C(x+1)(x+2) \\ &= A(x^2-4) + B(x^2-x-2) + C(x^2+3x+2) \\ &= Ax^2-4A+Bx^2-Bx-2B+Cx^2+3Cx+2C. \end{aligned}$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides,

$$x^2 \quad A + B + C = 0 \quad \dots(ii)$$

$$x \quad -B + 3C = 5 \quad \dots(iii)$$

$$\text{Constants} \quad -4A - 2B + 2C = 0$$

$$\text{Dividing by } -2, \quad 2A + B - C = 0 \quad \dots(iv)$$

Let us solve (ii), (iii) and (iv) for A, B, C

Eqn. (ii)  $\times 2$  - Eqn. (iv) gives (To eliminate A) because Eqn. (iii) does not involve A.

$$2A + 2B + 2C - (2A + B - C) = 0,$$

$$\text{i.e.,} \quad 2A + 2B + 2C - 2A - B + C = 0$$

$$\Rightarrow \quad B + 3C = 0 \quad \dots(v)$$

Adding Eqns. (iii) and (v),

$$6C = 5 \quad \Rightarrow \quad C = \frac{5}{6}$$

$$\text{Putting } C = \frac{5}{6} \text{ in (iii), } -B + \frac{15}{6} = 5 \quad \Rightarrow \quad -B = 5 - \frac{15}{6}$$

$$\Rightarrow \quad -B = \frac{30-15}{6} = \frac{15}{6} = \frac{5}{2} \quad \Rightarrow \quad B = -\frac{5}{2}$$

$$\text{Putting } B = -\frac{5}{2} \text{ and } C = \frac{5}{6} \text{ in (ii), } A - \frac{5}{2} + \frac{5}{6} = 0$$

$$\Rightarrow \quad A = \frac{5}{2} - \frac{5}{6} = \frac{15-5}{6} = \frac{10}{6} = \frac{5}{3}$$

Putting values of A, B, C in (i),

$$\frac{5x}{(x+1)(x^2-4)} = \frac{5}{3} \cdot \frac{1}{x+1} - \frac{5}{2} \cdot \frac{1}{x+2} + \frac{5}{6} \cdot \frac{1}{x-2}$$

$$\therefore \int \frac{5x}{(x+1)(x^2-4)} dx = \frac{5}{3} \int \frac{1}{x+1} dx - \frac{5}{2} \int \frac{1}{x+2} dx + \frac{5}{6} \int \frac{1}{x-2} dx$$

$$= \frac{5}{3} \log |x+1| - \frac{5}{2} \log |x+2| + \frac{5}{6} \log |x-2| + c.$$

$$12. \quad \frac{x^3 + x + 1}{x^2 - 1}$$

**Sol.** Here degree of numerator is greater than degree of denominator. Therefore, dividing the numerator by the denominator,

$$\begin{array}{r} x^2 - 1 \overline{) x^3 + x + 1} \phantom{0} \\ \underline{x^3 - x} \phantom{0} \\ 2x + 1 \phantom{0} \end{array}$$

$$\therefore \frac{x^3 + x + 1}{x^2 - 1} = x + \frac{2x + 1}{x^2 - 1} \quad \dots(i)$$

$$\left[ \text{Rational function} = \text{Quotient} + \frac{\text{Remainder}}{\text{Divisor}} \right]$$

$$\text{Let } \frac{2x+1}{x^2-1} = \frac{2x+1}{(x+1)(x-1)} = \frac{A}{x+1} + \frac{B}{x-1} \quad \dots(ii)$$

Multiplying by L.C.M. =  $(x+1)(x-1)$ , we have

$$2x+1 = A(x-1) + B(x+1)$$

or  $2x+1 = Ax - A + Bx + B$

By equating the coefficients of  $x$  and constant terms, we get

$$A + B = 2 \quad \dots(iii)$$

$$\text{and } -A + B = 1 \quad \dots(iv)$$

$$(iii) + (iv) \text{ gives } 2B = 3 \Rightarrow B = \frac{3}{2}$$

$$\text{Putting } B = \frac{3}{2} \text{ in (iii), we get } A + \frac{3}{2} = 2 \text{ or } A = \frac{1}{2}$$

Putting values of  $A$  and  $B$  in eqn. (ii), we have

$$\frac{2x+1}{x^2-1} = \frac{\frac{1}{2}}{x+1} + \frac{\frac{3}{2}}{x-1}$$

Putting this value of  $\frac{2x+1}{x^2-1}$  in (i),

$$\frac{x^3+x+1}{x^2-1} = x + \frac{\frac{1}{2}}{x+1} + \frac{\frac{3}{2}}{x-1}$$

$$\begin{aligned} \therefore \int \frac{x^3+x+1}{x^2-1} dx &= \int x dx + \frac{1}{2} \int \frac{1}{x+1} dx + \frac{3}{2} \int \frac{1}{x-1} dx \\ &= \frac{x^2}{2} + \frac{1}{2} \log |x+1| + \frac{3}{2} \log |x-1| + c. \end{aligned}$$

**Integrate the following functions in Exercises 13 to 17:**

$$13. \frac{2}{(1-x)(1+x^2)}$$

**Sol.** To find integral of the Rational function  $\frac{2}{(1-x)(1+x^2)}$ .

$$\text{Let integrand } \frac{2}{(1-x)(1+x^2)} = \frac{A}{1-x} + \frac{Bx+C}{1+x^2} \quad \dots(i)$$

(Partial Fractions)

Multiplying by L.C.M. =  $(1-x)(1+x^2)$

$$2 = A(1+x^2) + (Bx+C)(1-x)$$

or

$$2 = A + Ax^2 + Bx - Bx^2 + C - Cx$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms, we have

$$x^2 \quad A - B = 0 \quad \dots(ii)$$

$$x \quad B - C = 0 \quad \dots(iii)$$

$$\text{Constant terms } A + C = 2 \quad \dots(iv)$$

Let us solve (ii), (iii), (iv) for  $A$ ,  $B$ ,  $C$

From (ii),  $A = B$  and from (iii),  $B = C$

$\therefore A = B = C$   
 Putting  $A = C$  in (iv),  $C + C = 2$  or  $2C = 2$  or  $C = 1$   
 $\therefore A = C = 1$   $\therefore B = A = 1$   
 Putting these values of A, B, C in (i),

$$\begin{aligned}\frac{2}{(1-x)(1+x^2)} &= \frac{1}{1-x} + \frac{x+1}{1+x^2} = \frac{1}{1-x} + \frac{x}{1+x^2} + \frac{1}{1+x^2} \\ &= \frac{1}{1-x} + \frac{1}{2} \frac{2x}{1+x^2} + \frac{1}{1+x^2}\end{aligned}$$

$$\begin{aligned}\therefore \int \frac{2}{(1-x)(1+x^2)} dx &= \int \frac{1}{1-x} dx + \frac{1}{2} \int \frac{2x}{1+x^2} dx + \int \frac{1}{1+x^2} dx \\ &= \frac{\log |1-x|}{-1 \rightarrow \text{Coefficient of } x} + \frac{1}{2} \log |1+x^2| + \tan^{-1} x + c \\ &\quad \left[ \because \int \frac{2x}{1+x^2} dx = \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right] \\ &= -\log |1-x| + \frac{1}{2} \log (1+x^2) + \tan^{-1} x + c \\ &\quad (\because 1+x^2 > 0, \text{ therefore } |1+x^2| = 1+x^2) \\ \text{Note. } \log |1-x| &= \log |-(x-1)| \\ &= \log |x-1| \text{ because } |-t| = |t|.\end{aligned}$$

14.  $\frac{3x-1}{(x+2)^2}$

**Sol.** To find integral of rational function  $\frac{3x-1}{(x+2)^2}$ .

Let  $I = \int \frac{3x-1}{(x+2)^2} dx$  ... (i)

Form  $\int \frac{\text{Polynomial function}}{(\text{Linear})^k} dx$  where  $k$  is a positive integer,

put  $\text{Linear} = t$ .

Here put  $x+2 = t \Rightarrow x = t-2$

$\therefore \frac{dx}{dt} = 1 \Rightarrow dx = dt$

Putting these values in (i),

$$\begin{aligned}I &= \int \frac{3(t-2)-1}{t^2} dt = \int \frac{3t-6-1}{t^2} dt = \int \frac{3t-7}{t^2} dt \\ &= \int \left( \frac{3t}{t^2} - \frac{7}{t^2} \right) dt = \int \left( \frac{3}{t} - \frac{7}{t^2} \right) dt \\ &= 3 \int \frac{1}{t} dt - 7 \int t^{-2} dt = 3 \log |t| - 7 \frac{t^{-1}}{-1} + c \\ &= 3 \log |t| + \frac{7}{t} + c\end{aligned}$$

Putting  $t = x + 2$ ,  $= 3 \log |x + 2| + \frac{7}{x + 2} + c$ .

**Remark.** Alternative solution is Let  $\frac{3x-1}{(x+2)^2} = \frac{A}{x+2} + \frac{B}{(x+2)^2}$ .

15.  $\frac{1}{x^4 - 1}$

**Sol.** To find integral of  $\frac{1}{x^4 - 1}$ .

Let integrand  $\frac{1}{x^4 - 1} = \frac{1}{(x^2 - 1)(x^2 + 1)}$ .

Put  $x^2 = y$  **only** to form partial fractions.

$$= \frac{1}{(y-1)(y+1)} = \frac{A}{y-1} + \frac{B}{y+1} \quad \dots(i)$$

Multiplying by L.C.M.  $= (y-1)(y+1)$

$$1 = A(y+1) + B(y-1) \text{ or } 1 = Ay + A + By - B$$

Comparing coeffs. of  $y$  and constant terms, we have

$$\text{Coefficients of } y: \quad A + B = 0 \quad \dots(ii)$$

$$\text{Constant terms} \quad A - B = 1 \quad \dots(iii)$$

$$\text{Adding (ii) and (iii), } 2A = 1 \quad \Rightarrow \quad A = \frac{1}{2}$$

$$\text{Putting } A = \frac{1}{2} \text{ in (ii), } \frac{1}{2} + B = 0 \quad \Rightarrow \quad B = -\frac{1}{2}$$

Putting values of  $A, B$  and  $y$  in (i),

$$\frac{1}{x^4 - 1} = \frac{\frac{1}{2}}{x^2 - 1} - \frac{\frac{1}{2}}{x^2 + 1}$$

$$\begin{aligned} \therefore \int \frac{1}{x^4 - 1} dx &= \frac{1}{2} \int \frac{1}{x^2 - 1^2} dx - \frac{1}{2} \int \frac{1}{x^2 + 1} dx \\ &= \frac{1}{2} \cdot \frac{1}{2 \cdot 1} \log \left| \frac{x-1}{x+1} \right| - \frac{1}{2} \tan^{-1} x + c \end{aligned}$$

$$\left[ \because \int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| \right]$$

**Note.** Must put  $y = x^2$  in (i) along with values of  $A$  and  $B$  before writing values of integrals.

**Remark.** Alternative solution is:

$$\begin{aligned} \frac{1}{x^4 - 1} &= \frac{1}{(x^2 - 1)(x^2 + 1)} = \frac{1}{(x-1)(x+1)(x^2 + 1)} \\ &= \frac{A}{x-1} + \frac{B}{x+1} + \frac{Cx + D}{x^2 + 1} \end{aligned}$$

But the above given solution is better.

$$16. \frac{1}{x(x^n + 1)}$$

**Sol.** Let  $I = \int \frac{1}{x(x^n + 1)} dx$

Multiplying both numerator and denominator of integrand by  $nx^{n-1}$ .

$$\left[ \because \frac{d}{dx} (x^n + 1) = nx^{n-1} \right]$$

$$I = \int \frac{nx^{n-1}}{n x^{n-1} x(x^n + 1)} dx = \frac{1}{n} \int \frac{n x^{n-1}}{x^n (x^n + 1)} dx \quad \dots(i)$$

( $\because n - 1 + 1 = n$ )

**Put  $x^n = t$ .** Therefore  $n x^{n-1} = \frac{dt}{dx}$ .  $\therefore n x^{n-1} dx = dt$ .

$\therefore$  From (i),  $I = \frac{1}{n} \int \frac{dt}{t(t+1)} = \frac{1}{n} \int \frac{1}{t(t+1)} dt$

Adding and subtracting  $t$  in the numerator of integrand,

$$= \frac{1}{n} \int \frac{t+1-t}{t(t+1)} dt = \frac{1}{n} \int \left( \frac{t+1}{t(t+1)} - \frac{t}{t(t+1)} \right) dt \left[ \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right]$$

$$= \frac{1}{n} \left[ \int \frac{1}{t} dt - \int \frac{1}{t+1} dt \right] = \frac{1}{n} [\log |t| - \log |t+1| + c]$$

$$= \frac{1}{n} \log \left| \frac{t}{t+1} \right| + c$$

Putting  $t = x^n$ ,  $= \frac{1}{n} \log \left| \frac{x^n}{x^n + 1} \right| + c$

**Remark:** Alternative solution for  $\int \frac{1}{t(t+1)} dt$  is:

Let  $\frac{1}{t(t+1)} = \frac{A}{t} + \frac{B}{t+1}$

But the above given solution is better.

$$17. \frac{\cos x}{(1 - \sin x)(2 - \sin x)}$$

**Sol.** Let  $I = \int \frac{\cos x}{(1 - \sin x)(2 - \sin x)} dx \quad \dots(i)$

**Put  $\sin x = t$ .** Therefore  $\cos x = \frac{dt}{dx} \Rightarrow \cos x dx = dt$ ,

$\therefore$  From (i),  $\int \frac{1}{(1-t)(2-t)} dt = \int \frac{(2-t) - (1-t)}{(1-t)(2-t)} dt$

[ $\because$  Difference of two factors in the denominator namely  $1-t$  and  $2-t$  is  $(2-t) - (1-t) = 2-t-1+t = 1$ ]

$$= \int \left( \frac{2-t}{(1-t)(2-t)} - \frac{(1-t)}{(1-t)(2-t)} \right) dt \left[ \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right]$$

$$\begin{aligned}
 &= \int \left( \frac{1}{1-t} - \frac{1}{2-t} \right) dt = \int \frac{1}{1-t} dt - \int \frac{1}{2-t} dt \\
 &= \frac{\log |1-t|}{-1 \rightarrow \text{Coefficient of } t} - \frac{\log |2-t|}{-1} + c \\
 &= -\log |1-t| + \log |2-t| + c \\
 &= \log |2-t| - \log |1-t| + c = \log \left| \frac{2-t}{1-t} \right| + c
 \end{aligned}$$

Putting  $t = \sin x$ ,  $= \log \left| \frac{2 - \sin x}{1 - \sin x} \right| + c$

**Remark:** Alternative solution for  $\int \frac{1}{(1-t)(2-t)} dt$  is

Let  $\frac{1}{(1-t)(2-t)} = \frac{A}{1-t} + \frac{B}{2-t}$

**Integrate the following functions for Exercises 18 to 21:**

18.  $\frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)}$

**Sol.** To integrate the rational function  $\frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)}$  ... (i)

**Put  $x^2 = y$  in the integrand to get**

$$= \frac{(y+1)(y+2)}{(y+3)(y+4)} = \frac{y^2+3y+2}{y^2+7y+12} \quad \dots (ii)$$

Here degree of numerator = degree of denominator (= 2)

So have to perform long division to make the degree of numerator smaller than degree of denominator so that the concept of forming partial fractions becomes valid.

$$\begin{array}{r}
 y^2 + 7y + 12 \overline{) y^2 + 3y + 2} \quad (1) \\
 \underline{y^2 + 7y + 12} \phantom{0} \\
 -4y - 10
 \end{array}$$

$\therefore$  From (i) and (ii),

$$\frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)} = \frac{(y+1)(y+2)}{(y+3)(y+4)} = 1 + \frac{(-4y-10)}{(y+3)(y+4)} \quad \dots (iii)$$

Let us form partial fractions of  $\frac{(-4y-10)}{(y+3)(y+4)}$ .

Let  $\frac{-4y-10}{(y+3)(y+4)} = \frac{A}{y+3} + \frac{B}{y+4} \quad \dots (iv)$

Multiplying by L.C.M. =  $(y+3)(y+4)$

$$-4y - 10 = A(y+4) + B(y+3) = Ay + 4A + By + 3B$$

Comparing coefficients of  $y$ ,  $A + B = -4 \quad \dots (v)$

Comparing constants,  $4A + 3B = -10 \quad \dots (vi)$

Let us solve Eqns. (v) and (vi) for A and B.

Eqn. (v)  $\times 4$  gives,  $4A + 4B = -16 \quad \dots (vii)$

Eqn. (vi) – Eqn. (vii) gives,  $-B = 6$  or  $B = -6$ .

Putting  $B = -6$  in (v),  $A - 6 = -4 \Rightarrow A = -4 + 6 = 2$

Putting these values of A and B in (iv),

$$\frac{-4y - 10}{(y+3)(y+4)} = \frac{2}{y+3} - \frac{6}{y+4}$$

Putting this value in (iii),

$$\frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)} = 1 + \frac{2}{y+3} - \frac{6}{y+4}$$

**In R.H.S., Putting  $y = x^2$  (before integration)**

$$= 1 + \frac{2}{x^2+3} - \frac{6}{x^2+4}$$

$$\begin{aligned} \therefore \int \frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)} dx \\ &= \int 1 dx + 2 \int \frac{1}{x^2+(\sqrt{3})^2} dx - 6 \int \frac{1}{x^2+2^2} dx \\ &= x + 2 \cdot \frac{1}{\sqrt{3}} \tan^{-1} \frac{x}{\sqrt{3}} - 6 \cdot \frac{1}{2} \tan^{-1} \frac{x}{2} + c \\ &= x + \frac{2}{\sqrt{3}} \tan^{-1} \frac{x}{\sqrt{3}} - 3 \tan^{-1} \frac{x}{2} + c. \end{aligned}$$

19.  $\frac{2x}{(x^2+1)(x^2+3)}$

**Sol.** Let  $I = \int \frac{2x}{(x^2+1)(x^2+3)} dx$

**Put  $x^2 = t$ .** Differentiating both sides  $2x dx = dt$

$$\therefore I = \int \frac{dt}{(t+1)(t+3)}$$

Dividing and multiplying by 2,

$$(\because (t+3) - (t+1) = t+3 - t - 1 = 2)$$

$$\begin{aligned} &= \frac{1}{2} \int \frac{2}{(t+1)(t+3)} dt = \frac{1}{2} \int \frac{(t+3) - (t+1)}{(t+1)(t+3)} dt \\ &= \frac{1}{2} \int \left( \frac{1}{t+1} - \frac{1}{t+3} \right) dt = \frac{1}{2} [\log |t+1| - \log |t+3|] + c \\ &= \frac{1}{2} \log \left| \frac{t+1}{t+3} \right| + c = \frac{1}{2} \log \left| \frac{x^2+1}{x^2+3} \right| + c = \frac{1}{2} \log \left( \frac{x^2+1}{x^2+3} \right) + c. \end{aligned}$$

20.  $\frac{1}{x(x^4-1)}$

**Sol.** Let  $I = \int \frac{1}{x(x^4-1)} dx$

Multiplying both numerator and denominator of integrand by  $4x^3$ .

$$\left( \because \frac{d}{dx} (x^4-1) = 4x^3 \right)$$

$$I = \int \frac{4x^3}{4x^4(x^4-1)} dx = \frac{1}{4} \int \frac{4x^3}{x^4(x^4-1)} dx \quad \dots(i)$$

**Put  $x^4 = t$ .** Therefore  $4x^3 = \frac{dt}{dx} \Rightarrow 4x^3 dx = dt$ .

$$\therefore \text{ From (i), } I = \frac{1}{4} \int \frac{dt}{t(t-1)} = \frac{1}{4} \int \frac{t-(t-1)}{t(t-1)} dt$$

$$[\because t-(t-1) = t-t+1 = 1]$$

$$= \frac{1}{4} \int \left( \frac{t}{t(t-1)} - \frac{(t-1)}{t(t-1)} \right) dt = \frac{1}{4} \int \left( \frac{1}{t-1} - \frac{1}{t} \right) dt$$

$$= \frac{1}{4} \left[ \int \frac{1}{t-1} dt - \int \frac{1}{t} dt \right] = \frac{1}{4} [\log |t-1| - \log |t|] + c$$

$$= \frac{1}{4} \log \left| \frac{t-1}{t} \right| + c$$

$$\text{Putting } t = x^4, = \frac{1}{4} \log \left| \frac{x^4-1}{x^4} \right| + c.$$

**Remark:** Alternative solution is:

$$\frac{1}{x(x^4-1)} = \frac{1}{x(x^2-1)(x^2+1)} = \frac{1}{x(x-1)(x+1)(x^2+1)}$$

$$= \frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+1} + \frac{Dx+E}{x^2+1}$$

But the solution given above is much better.

21.  $\frac{1}{(e^x-1)}$

**Sol.** Let  $I = \int \frac{1}{e^x-1} dx \quad \dots(i)$

**Put  $e^x = t$ .** Therefore  $e^x = \frac{dt}{dx} \Rightarrow e^x dx = dt \Rightarrow dx = \frac{dt}{e^x}$

**(Rule to evaluate  $\int f(e^x) dx$ , put  $e^x = t$ )**

$$\therefore \text{ From (i), } I = \int \frac{1}{t-1} \frac{dt}{e^x} = \int \frac{1}{t-1} \frac{dt}{t} = \int \frac{1}{t(t-1)} dt$$

$$= \int \frac{t-(t-1)}{t(t-1)} dt = \int \left( \frac{t}{t(t-1)} - \frac{(t-1)}{t(t-1)} \right) dt = \int \frac{1}{t-1} dt - \int \frac{1}{t} dt$$

$$= \log |t-1| - \log |t| + c = \log \left| \frac{t-1}{t} \right| + c.$$

$$\text{Putting } t = e^x, = \log \left| \frac{e^x-1}{e^x} \right| + c.$$

Choose the correct answer in each of the Exercises 22 and 23:

22.  $\int \frac{x \, dx}{(x-1)(x-2)}$  equals

(A)  $\log \left| \frac{(x-1)^2}{x-2} \right| + C$

(B)  $\log \left| \frac{(x-2)^2}{x-1} \right| + C$

(C)  $\log \left| \left( \frac{x-1}{x-2} \right)^2 \right| + C$

(D)  $\log | (x-1)(x-2) | + C.$

**Sol.** Let integrand  $\frac{x}{(x-1)(x-2)} = \frac{A}{x-1} + \frac{B}{x-2}$  ... (i)

(Partial fractions)

Multiplying by L.C.M. =  $(x-1)(x-2)$ ,  
 $x = A(x-2) + B(x-1)$   
 $= Ax - 2A + Bx - B$

Comparing coefficients of  $x$  and constant terms on both sides,

Coefficients of  $x$ :  $A + B = 1$  ... (ii)

Constant terms:  $-2A - B = 0$  ... (iii)

Let us solve (ii) and (iii) for  $A$  and  $B$

Adding (ii) and (iii),  $-A = 1$  or  $A = -1$

Putting  $A = -1$  in (ii)  $-1 + B = 1$  or  $B = 2$

Putting values of  $A$  and  $B$  in (i),

$$\frac{x}{(x-1)(x-2)} = \frac{-1}{x-1} + \frac{2}{x-2}$$

$$\begin{aligned} \therefore \int \frac{x}{(x-1)(x-2)} \, dx &= - \int \frac{1}{x-1} \, dx + 2 \int \frac{1}{x-2} \, dx \\ &= -\log |x-1| + 2 \log |x-2| + c \\ &= \log |(x-2)^2| - \log |x-1| + c \\ &\quad (\because n \log m = \log m^n) \\ &= \log \left| \frac{(x-2)^2}{x-1} \right| + c \end{aligned}$$

$\therefore$  Option (B) is the correct answer.

23.  $\int \frac{dx}{x(x^2+1)}$  equals

(A)  $\log |x| - \frac{1}{2} \log (x^2 + 1) + C$

(B)  $\log |x| + \frac{1}{2} \log (x^2 + 1) + C$

(C)  $-\log |x| + \frac{1}{2} \log (x^2 + 1) + C$

(D)  $\frac{1}{2} \log |x| + \log (x^2 + 1) + C.$

**Sol.** Let  $I = \int \frac{1}{x(x^2+1)} \, dx$

Multiplying both numerator and denominator of integrand by  $2x$ .

$$\left( \because \frac{d}{dx} (x^2 + 1) = 2x \right)$$

$$\Rightarrow I = \int \frac{2x}{2x^2(x^2 + 1)} dx \quad \dots(i)$$

**Put  $x^2 = t$ .**  $\therefore 2x = \frac{dt}{dx} \Rightarrow 2x dx = dt$

$$\therefore \text{ From (i), } I = \int \frac{dt}{2t(t+1)} = \frac{1}{2} \int \frac{1}{t(t+1)} dt$$

Adding and subtracting  $t$  in the numerator of integrand,

$$\begin{aligned} &= \frac{1}{2} \int \frac{(t+1) - t}{t(t+1)} dt = \frac{1}{2} \int \left( \frac{1}{t} - \frac{1}{t+1} \right) dt \\ &= \frac{1}{2} (\log |t| - \log |t+1|) + c \end{aligned}$$

Putting  $t = x^2$ ,  $I = \frac{1}{2} (\log |x^2| - \log |x^2 + 1|) + c$

$$\begin{aligned} &= \frac{1}{2} (2 \log |x| - \log (x^2 + 1)) + c \\ &\quad (\because x^2 + 1 \geq 1 > 0 \text{ and hence } |x^2 + 1| = x^2 + 1) \\ &= \log |x| - \frac{1}{2} \log (x^2 + 1) + c \end{aligned}$$

$\therefore$  Option (A) is the correct answer.

## Exercise 7.6

Integrate the functions in Exercises 1 to 8:

1.  $x \sin x$

Sol.  $\int x \sin x \, dx$   
I II

$$\begin{aligned}\text{Applying Product Rule I } \int \text{II} \, dx &= \int \left( \frac{d}{dx} (\text{I}) \int \text{II} \, dx \right) dx \\ &= x \int \sin x \, dx - \int \left( \frac{d}{dx} (x) \int \sin x \, dx \right) dx \\ &= x (-\cos x) - \int 1 (-\cos x) \, dx = -x \cos x - \int -\cos x \, dx \\ &= -x \cos x + \int \cos x \, dx = -x \cos x + \sin x + c\end{aligned}$$

**Note.**  $\int \sin x \, dx = -\cos x$ .

2.  $x \sin 3x$

Sol.  $\int x \sin 3x \, dx$   
I II

$$\begin{aligned}\text{Applying Product Rule I } \int \text{II} \, dx &= \int \left( \frac{d}{dx} (\text{I}) \int \text{II} \, dx \right) dx \\ &= x \int \sin 3x \, dx - \int \left( \frac{d}{dx} (x) \int \sin 3x \, dx \right) dx \\ &= x \left( \frac{-\cos 3x}{3} \right) - \int \left[ 1 \left( \frac{-\cos 3x}{3} \right) \right] dx + c \\ &= \frac{-1}{3} x \cos 3x + \frac{1}{3} \int \cos 3x \, dx + c\end{aligned}$$

$$= \frac{-1}{3} x \cos 3x + \frac{1}{3} \frac{\sin 3x}{3} + c = \frac{-1}{3} x \cos 3x + \frac{1}{9} \sin 3x + c.$$

3.  $x^2 e^x$

**Sol.**  $\int \underset{\text{I}}{x^2} \underset{\text{II}}{e^x} dx$

Applying Product Rule I  $\int \text{II } dx - \int \left( \frac{d}{dx} (\text{I}) \int \text{II } dx \right) dx$

$$= x^2 \int e^x dx - \int \left[ \left( \frac{d}{dx} x^2 \right) \int e^x dx \right] dx = x^2 e^x - \int 2x e^x dx$$

$$= x^2 e^x - 2 \int \underset{\text{I}}{x} \underset{\text{II}}{e^x} dx$$

Again Applying Product Rule

$$= x^2 e^x - 2 \left[ x \int e^x dx - \int \left[ \frac{d}{dx} (x) \int e^x dx \right] dx \right]$$

$$= x^2 e^x - 2 \left( x e^x - \int 1 \cdot e^x dx \right) = x^2 e^x - 2 \left( x e^x - \int e^x dx \right)$$

$$= x^2 e^x - 2x e^x + 2 \int e^x dx + c = x^2 e^x - 2x e^x + 2e^x + c$$

$$= e^x (x^2 - 2x + 2) + c.$$

4.  $x \log x$

**Sol.**  $\int \underset{\text{I}}{x} \log \underset{\text{II}}{x} dx = \int (\log x) \cdot \underset{\text{II}}{x} dx$

Applying Product Rule I  $\int \text{II } dx - \int \left[ \frac{d}{dx} (\text{I}) \int \text{II } dx \right] dx$

$$= (\log x) \int x dx - \int \left[ \frac{d}{dx} (\log x) \int x dx \right] dx$$

$$= (\log x) \frac{x^2}{2} - \int \frac{1}{x} \frac{x^2}{2} dx = \frac{1}{2} x^2 \log x - \frac{1}{2} \int x dx$$

$$\left( \because \frac{x^2}{x} = \frac{x \cdot x}{x} = x \right)$$

$$= \frac{1}{2} x^2 \log x - \frac{1}{2} \frac{x^2}{2} + c = \frac{x^2}{2} \log x - \frac{x^2}{4} + c.$$

5.  $x \log 2x$

**Sol.**  $\int \underset{\text{I}}{x} \log \underset{\text{II}}{2x} dx = \int (\log 2x) \cdot \underset{\text{II}}{x} dx$

Applying Product Rule I  $\int \text{II } dx - \int \left( \frac{d}{dx} (\text{I}) \int \text{II } dx \right) dx$

$$= (\log 2x) \int x dx - \int \left( \frac{d}{dx} (\log 2x) \int x dx \right) dx$$

$$= (\log 2x) \frac{x^2}{2} - \int \frac{1}{2x} \cdot 2 \cdot \frac{x^2}{2} dx$$

$$\begin{aligned}
 &= \frac{1}{2} x^2 \log 2x - \frac{1}{2} \int x \, dx \quad \left[ \because \frac{x^2}{x} = \frac{x \cdot x}{x} = x \right] \\
 &= \frac{1}{2} x^2 \log 2x - \frac{1}{2} \cdot \frac{x^2}{2} + c = \frac{x^2}{2} \log 2x - \frac{x^2}{4} + c.
 \end{aligned}$$

### 6. $x^2 \log x$

**Sol.**  $\int x^2 \log x \, dx = \int \underset{\text{I}}{(\log x)} \underset{\text{II}}{x^2 \, dx}$

Applying Product Rule:  $\int \text{I} \, \text{II} \, dx = \int \left( \frac{d}{dx} (\text{I}) \int \text{II} \, dx \right) dx$

$$= \log x \int x^2 \, dx - \int \left( \frac{d}{dx} (\log x) \int x^2 \, dx \right) dx$$

$$\begin{aligned}
 &= (\log x) \frac{x^3}{3} - \int \frac{1}{x} \cdot \frac{x^3}{3} \, dx = \frac{x^3}{3} \log x - \frac{1}{3} \int x^2 \, dx \quad \left[ \because \frac{x^3}{x} = x^2 \right] \\
 &= \frac{x^3}{3} \log x - \frac{1}{3} \cdot \frac{x^3}{3} + c = \frac{x^3}{3} \log x - \frac{x^3}{9} + c.
 \end{aligned}$$

### 7. $x \sin^{-1} x$

**Sol.** Let  $\text{I} = \int x \sin^{-1} x \, dx$ .

**Put  $x = \sin \theta$ .** Differentiating both sides  $dx = \cos \theta \, d\theta$

$$\begin{aligned}
 \therefore \text{I} &= \int \sin \theta \cdot \theta \cdot \cos \theta \, d\theta = \frac{1}{2} \int \theta \cdot 2 \sin \theta \cos \theta \, d\theta \\
 &= \frac{1}{2} \int \theta \sin 2\theta \, d\theta
 \end{aligned}$$

Integrating by parts

$$\begin{aligned}
 &= \frac{1}{2} \left[ \theta \left( -\frac{\cos 2\theta}{2} \right) - \int 1 \cdot \left( -\frac{\cos 2\theta}{2} \right) d\theta \right] \\
 &= \frac{1}{4} \left[ -\theta \cos 2\theta + \int \cos 2\theta \, d\theta \right] = \frac{1}{4} \left[ -\theta \cos 2\theta + \frac{\sin 2\theta}{2} \right] + c \\
 &= \frac{1}{4} [-\theta (1 - 2 \sin^2 \theta) + \sin \theta \cos \theta] + c \\
 &\quad (\because \sin 2\theta = 2 \sin \theta \cos \theta) \\
 &= \frac{1}{4} [-\sin^{-1} x \cdot (1 - 2x^2) + x \sqrt{1 - x^2}] + c
 \end{aligned}$$

$$\left[ \because \cos \theta = \sqrt{1 - \sin^2 \theta} = \sqrt{1 - x^2} \right]$$

$$= \frac{1}{4} (2x^2 - 1) \sin^{-1} x + \frac{x\sqrt{1-x^2}}{4} + c.$$

### 8. $x \tan^{-1} x$

**Sol.** Let  $\text{I} = \int x \tan^{-1} x \, dx = \int (\tan^{-1} x) \cdot x \, dx$

$$= (\tan^{-1} x) \cdot \frac{x^2}{2} - \int \frac{1}{1+x^2} \cdot \frac{x^2}{2} \, dx$$

$$\begin{aligned}
&= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} dx \\
&= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} \int \left( 1 - \frac{1}{1+x^2} \right) dx \\
&\quad \left[ \because \frac{x^2}{1+x^2} = \frac{1+x^2-1}{1+x^2} = 1 - \frac{1}{1+x^2} \right] \\
&= \frac{x^2}{2} \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x) + c \\
&= \frac{1}{2} [x^2 \tan^{-1} x - x + \tan^{-1} x] + c = \frac{1}{2} [(x^2 + 1) \tan^{-1} x - x] + c.
\end{aligned}$$

**Integrate the functions in Exercises 9 to 15:**

**9.  $x \cos^{-1} x$**

**Sol.** Let  $I = \int x \cos^{-1} x \, dx$  ... (i)

**Put  $\cos^{-1} x = \theta$ .** Therefore  $x = \cos \theta$ .

$$\therefore \frac{dx}{d\theta} = -\sin \theta \Rightarrow dx = -\sin \theta \, d\theta$$

$$\begin{aligned}
\therefore \text{From (i), } I &= \int (\cos \theta) \theta (-\sin \theta \, d\theta) = \frac{-1}{2} \int \theta (2 \sin \theta \cos \theta) \, d\theta \\
&= \frac{-1}{2} \int \theta \sin 2\theta \, d\theta \\
&\quad \text{I} \quad \text{II}
\end{aligned}$$

Applying Product Rule:  $I \int II \, d\theta - \int \left[ \frac{d}{d\theta} (I) \int II \, d\theta \right] d\theta$

$$\begin{aligned}
&= \frac{-1}{2} \left[ \theta \left( \frac{-\cos 2\theta}{2} \right) - \int 1 \left( \frac{-\cos 2\theta}{2} \right) d\theta \right] \\
&= \frac{-1}{2} \left[ \frac{-1}{2} \theta \cos 2\theta + \frac{1}{2} \int \cos 2\theta \, d\theta \right] = \frac{1}{4} \theta \cos 2\theta - \frac{1}{4} \left( \frac{\sin 2\theta}{2} \right) + c \\
&= \frac{1}{4} \theta \cos 2\theta - \frac{1}{8} (2 \sin \theta \cos \theta) + c \\
&= \frac{1}{4} \theta (2 \cos^2 \theta - 1) - \frac{1}{4} \sqrt{1 - \cos^2 \theta} \cdot \cos \theta + c \\
&\text{Putting } \cos \theta = x \text{ and } \theta = \cos^{-1} x; \\
&= \frac{1}{4} (\cos^{-1} x) (2x^2 - 1) - \frac{1}{4} \sqrt{1 - x^2} \cdot x + c \\
&= (2x^2 - 1) \frac{\cos^{-1} x}{4} - \frac{x}{4} \sqrt{1 - x^2} + c.
\end{aligned}$$

**10.  $(\sin^{-1} x)^2$**

**Sol.** **Put  $x = \sin \theta$ .** Differentiating both sides,  $dx = \cos \theta \, d\theta$

$$\begin{aligned}
\therefore \int (\sin^{-1} x)^2 dx &= \int \theta^2 \cos \theta \, d\theta = \theta^2 \sin \theta - \int 2\theta \sin \theta \, d\theta \\
&\quad \text{I} \quad \text{II} \\
&= \theta^2 \sin \theta - 2 \int \theta \sin \theta \, d\theta \\
&\quad \text{I} \quad \text{II}
\end{aligned}$$

$$\begin{aligned}
 &= \theta^2 \sin \theta - 2 \left[ \theta (-\cos \theta) - \int 1 \cdot (-\cos \theta) d\theta \right] \\
 &= \theta^2 \sin \theta + 2\theta \cos \theta - 2 \int \cos \theta d\theta = \theta^2 \sin \theta + 2\theta \cos \theta - 2 \sin \theta + c \\
 &= x (\sin^{-1} x)^2 + 2\sqrt{1-x^2} \sin^{-1} x - 2x + c. \\
 &\left( \because \cos \theta = \sqrt{1-\sin^2 \theta} = \sqrt{1-x^2} \right)
 \end{aligned}$$

11.  $\frac{x \cos^{-1} x}{\sqrt{1-x^2}}$

**Sol.** Let  $I = \int \frac{x \cos^{-1} x}{\sqrt{1-x^2}} dx$  ... (i)

**Put**  $\cos^{-1} x = \theta$ .  $\Rightarrow x = \cos \theta$

Therefore  $\frac{dx}{d\theta} = -\sin \theta \Rightarrow dx = -\sin \theta d\theta$

$$\begin{aligned}
 \therefore \text{ From (i), } I &= \int \frac{(\cos \theta) \theta}{\sqrt{1-\cos^2 \theta}} (-\sin \theta d\theta) \\
 &= - \int \frac{\theta \cos \theta \sin \theta}{\sin \theta} d\theta \quad (\because \sqrt{1-\cos^2 \theta} = \sqrt{\sin^2 \theta} = \sin \theta) \\
 &= - \int \theta \cos \theta d\theta \\
 &\quad \text{I} \quad \text{II}
 \end{aligned}$$

Applying Product Rule:  $I \int II d\theta = \int \left[ \frac{d}{d\theta} (I) \int II d\theta \right] d\theta$

$$\begin{aligned}
 &= - \left[ \theta \cdot \sin \theta - \int 1 \cdot \sin \theta d\theta \right] = - \theta \sin \theta + \int \sin \theta d\theta \\
 &= - \theta \sin \theta - \cos \theta + c = - \theta \sqrt{1-\cos^2 \theta} - \cos \theta + c
 \end{aligned}$$

Putting  $\theta = \cos^{-1} x$  and  $\cos \theta = x$ ,

$$= - (\cos^{-1} x) \sqrt{1-x^2} - x + c = - [\sqrt{1-x^2} \cos^{-1} x + x] + c.$$

12.  $x \sec^2 x$

**Sol.**  $\int x \sec^2 x dx$   
I II

Applying Product Rule:  $I \int II dx = \int \left[ \frac{d}{dx} (I) \int II dx \right] dx$

$$\begin{aligned}
 &= x \int \sec^2 x dx - \int \left[ \frac{d}{dx} (x) \int \sec^2 x dx \right] dx \\
 &= x \tan x - \int 1 \cdot \tan x dx = x \tan x - \int \tan x dx \\
 &= x \tan x - (-\log |\cos x|) + c = x \tan x + \log |\cos x| + c.
 \end{aligned}$$

13.  $\tan^{-1} x$

**Sol.** Let  $I = \int \tan^{-1} x dx = \int (\tan^{-1} x) \cdot 1 dx$

$$\begin{aligned}
 &= \tan^{-1} x \cdot x - \int \frac{1}{1+x^2} \cdot x dx = x \tan^{-1} x - \frac{1}{2} \int \frac{2x}{1+x^2} dx \\
 &= x \tan^{-1} x - \frac{1}{2} \log |(1+x^2)| + c. \left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right]
 \end{aligned}$$

$$= x \tan^{-1} x - \frac{1}{2} \log (1 + x^2) + c$$

$[\because 1 + x^2 \geq 1 > 0 \text{ and hence } |1 + x^2| = 1 + x^2]$

**14.  $x (\log x)^2$**

**Sol.**  $\int x (\log x)^2 dx = \int \underset{\text{I}}{(\log x)^2} \cdot \underset{\text{II}}{x dx}$

Applying Product Rule:  $\int \underset{\text{I}}{(\log x)^2} \underset{\text{II}}{x dx} - \int \left[ \frac{d}{dx} (\text{I}) \int \text{II} dx \right] dx$

$$= (\log x)^2 \int x dx - \int \left[ \frac{d}{dx} (\log x)^2 \int x dx \right] dx$$

$$= (\log x)^2 \frac{x^2}{2} - \int \frac{2(\log x)}{x} \frac{x^2}{2} dx$$

$$\left[ \because \frac{d}{dx} (\log x)^2 = 2(\log x)^1 \frac{d}{dx} (\log x) = 2 \log x \cdot \frac{1}{x} = \frac{2 \log x}{x} \right]$$

$$= \underset{\text{I}}{\frac{x^2}{2}} (\log x)^2 - \int \underset{\text{I}}{(\log x)} \underset{\text{II}}{x dx} \quad \left[ \because \frac{x^2}{x} = \frac{x \cdot x}{x} = x \right]$$

Again applying Product Rule:  $\int \underset{\text{I}}{(\log x)} \underset{\text{II}}{x dx} - \int \left[ \frac{d}{dx} (\text{I}) \int \text{II} dx \right] dx$

$$= \frac{x^2}{2} (\log x)^2 - \left[ (\log x) \frac{x^2}{2} - \int \left( \frac{1}{x} \frac{x^2}{2} \right) dx \right] + c$$

$$= \frac{x^2}{2} (\log x)^2 - \frac{x^2}{2} \log x + \frac{1}{2} \int x dx + c$$

$$= \frac{x^2}{2} (\log x)^2 - \frac{x^2}{2} \log x + \frac{x^2}{4} + c.$$

**15.  $(x^2 + 1) \log x$**

**Sol.**  $\int (x^2 + 1) \log x dx = \int \underset{\text{I}}{(\log x)} \underset{\text{II}}{(x^2 + 1) dx}$

Applying Product Rule:  $\int \underset{\text{I}}{(\log x)} \underset{\text{II}}{(x^2 + 1) dx} - \int \left[ \frac{d}{dx} (\text{I}) \int \text{II} dx \right] dx$

$$= \log x \left( \frac{x^3}{3} + x \right) - \int \frac{1}{x} \left( \frac{x^3}{3} + x \right) dx$$

$$= \left( \frac{x^3}{3} + x \right) \log x - \int \left( \frac{x^2}{3} + 1 \right) dx$$

$$= \left( \frac{x^3}{3} + x \right) \log x - \frac{1}{3} \int x^2 dx - \int 1 dx$$

$$= \left( \frac{x^3}{3} + x \right) \log x - \frac{1}{3} \frac{x^3}{3} - x + c = \left( \frac{x^3}{3} + x \right) \log x - \frac{x^3}{9} - x + c.$$

**Integrate the functions in Exercises 16 to 22:****16.  $e^x (\sin x + \cos x)$** **Sol.** Here  $I = \int e^x (\sin x + \cos x) dx$ It is of the form  $\int e^x [f(x) + f'(x)] dx$ Let us take  $f(x) = \sin x$  so that  $f'(x) = \cos x$ 

$$I = e^x f(x) + c = e^x \sin x + c.$$

$$\left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) + c \right]$$

**17.  $\frac{x e^x}{(1+x)^2}$** **Sol.** Here  $I = \int \frac{x e^x}{(x+1)^2} dx = \int \frac{(x+1)-1}{(x+1)^2} e^x dx$ 

$$= \int e^x \left[ \frac{x+1}{(x+1)^2} - \frac{1}{(x+1)^2} \right] dx = \int e^x \left[ \frac{1}{x+1} + \frac{-1}{(x+1)^2} \right] dx$$

It is of the form  $\int e^x [f(x) + f'(x)] dx$ Let us take  $f(x) = \frac{1}{x+1}$  so that  $f'(x) = \frac{d}{dx} [(x+1)^{-1}]$ 

$$= -(x+1)^{-2} = \frac{-1}{(x+1)^2}$$

$$\therefore I = e^x f(x) + c = \frac{e^x}{x+1} + c. \left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) + c \right]$$

**18.  $e^x \left( \frac{1+\sin x}{1+\cos x} \right)$** **Sol.** Here  $I = \int e^x \cdot \frac{1+\sin x}{1+\cos x} dx = \int e^x \cdot \frac{1+2\sin \frac{x}{2} \cos \frac{x}{2}}{2 \cos^2 \frac{x}{2}} dx$ 

$$= \int e^x \cdot \left[ \frac{1}{2 \cos^2 \frac{x}{2}} + \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \cos^2 \frac{x}{2}} \right] dx = \int e^x \left( \frac{1}{2} \sec^2 \frac{x}{2} + \tan \frac{x}{2} \right) dx$$

$$= \int e^x \left( \tan \frac{x}{2} + \frac{1}{2} \sec^2 \frac{x}{2} \right) dx$$

It is of the form  $\int e^x [f(x) + f'(x)] dx$ Let us take  $f(x) = \tan \frac{x}{2}$  so that  $f'(x) = \frac{1}{2} \sec^2 \frac{x}{2}$ 

$$\therefore I = e^x f(x) + c = e^x \tan \frac{x}{2} + c.$$

$$\left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) + c \right]$$

$$19. e^x \left( \frac{1}{x} - \frac{1}{x^2} \right)$$

$$\text{Sol. Let } I = \int e^x \left( \frac{1}{x} - \frac{1}{x^2} \right) dx$$

It is of the form  $\int e^x (f(x) + f'(x)) dx$

$$\text{Here } f(x) = \frac{1}{x} = x^{-1} \quad \text{and} \quad \text{so } f'(x) = (-1) x^{-2} = \frac{-1}{x^2}$$

$$\therefore I = e^x f(x) + c \quad [\because \int e^x (f(x) + f'(x)) dx = e^x f(x) + c]$$

$$= e^x \frac{1}{x} + c = \frac{e^x}{x} + c.$$

$$20. \frac{(x-3)e^x}{(x-1)^3}$$

$$\text{Sol. Here } I = \int \frac{(x-3)e^x}{(x-1)^3} dx = \int \frac{(x-1)-2}{(x-1)^3} e^x dx$$

$$= \int e^x \left[ \frac{x-1}{(x-1)^3} - \frac{2}{(x-1)^3} \right] dx = \int e^x \left[ \frac{1}{(x-1)^2} + \frac{-2}{(x-1)^3} \right] dx$$

It is of the form  $\int e^x [f(x) + f'(x)] dx$

$$\text{Let us take } f(x) = \frac{1}{(x-1)^2} \quad \text{so that } f'(x) = \frac{d}{dx} [(x-1)^{-2}]$$

$$= -2(x-1)^{-3} = \frac{-2}{(x-1)^3}$$

$$\therefore I = e^x f(x) + c = \frac{e^x}{(x-1)^2} + c.$$

$$[\because \int e^x (f(x) + f'(x)) dx = e^x f(x)]$$

**Note. Rule to evaluate**  $\int e^{ax} \sin bx dx$  or  $\int e^{ax} \cos bx dx$

$$\text{Let } I = \int e^{ax} \sin bx dx \quad \text{or} \quad \int e^{ax} \cos bx dx$$

$$\text{I} \quad \text{II} \quad \text{I} \quad \text{II}$$

Integrate twice by product Rule and transpose term containing I from R.H.S. to L.H.S.

$$21. e^{2x} \sin x$$

$$\text{Sol. Let } I = \int e^{2x} \sin x dx \quad \dots(i)$$

$$\text{Applying Product Rule: } I \int \text{II} dx - \int \left[ \frac{d}{dx} (\text{I}) \int \text{II} dx \right] dx$$

$$\Rightarrow I = e^{2x} (-\cos x) - \int e^{2x} \cdot 2 \cdot (-\cos x) dx$$

$$\left[ \because \frac{d}{dx} e^{2x} = e^{2x} \frac{d}{dx} (2x) = 2e^{2x} \right]$$

$$\Rightarrow I = -e^{2x} \cos x + 2 \int e^{2x} \cos x \, dx$$

Again Applying Product Rule:

$$I = -e^{2x} \cos x + 2 \left[ e^{2x} \sin x - \int 2e^{2x} \sin x \, dx \right]$$

$$\Rightarrow I = -e^{2x} \cos x + 2e^{2x} \sin x - 4 \int e^{2x} \sin x \, dx$$

$$\Rightarrow I = e^{2x} (-\cos x + 2 \sin x) - 4I \quad [\text{By (i)}]$$

$$\text{Transposing } -4I \text{ to L.H.S.; } 5I = e^{2x} (2 \sin x - \cos x)$$

$$\therefore I \left( = \int e^{2x} \sin x \, dx \right) = \frac{e^{2x}}{5} (2 \sin x - \cos x) + c$$

**Remark:** The above question can also be done as:

Applying Product Rule: taking  $\sin x$  as first function and  $e^{2x}$  as second function.

22.  $\sin^{-1} \left( \frac{2x}{1+x^2} \right)$

**Sol.** Put  $x = \tan \theta$ . Differentiating both sides  $dx = \sec^2 \theta \, d\theta$ .

$$\therefore \int \sin^{-1} \left( \frac{2x}{1+x^2} \right) dx = \int \sin^{-1} \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right) \cdot \sec^2 \theta \, d\theta$$

$$= \int \sin^{-1} (\sin 2\theta) \cdot \sec^2 \theta \, d\theta = \int 2\theta \sec^2 \theta \, d\theta$$

$$= 2 \int \theta \sec^2 \theta \, d\theta$$

Applying product rule

$$= 2 [\theta \cdot \tan \theta - \int 1 \cdot \tan \theta \, d\theta] = 2 [\theta \tan \theta - \int \tan \theta \, d\theta]$$

$$= 2 [\theta \tan \theta - \log \sec \theta] + c$$

$$= 2 [\tan^{-1} x \cdot x - \log \sqrt{1+x^2}] + c$$

$$[\because \sec \theta = \sqrt{1 + \tan^2 \theta} = \sqrt{1 + x^2}]$$

$$= 2 \left[ x \tan^{-1} x - \frac{1}{2} \log (1 + x^2) \right] + c$$

$$= 2x \tan^{-1} x - \log (1 + x^2) + c.$$

**Choose the correct answer in Exercises 23 and 24.**

23.  $\int x^2 e^{x^3} \, dx$  equals

(A)  $\frac{1}{3} e^{x^3} + C$

(B)  $\frac{1}{3} e^{x^2} + C$

(C)  $\frac{1}{2} e^{x^3} + C$

(D)  $\frac{1}{2} e^{x^2} + C$

**Sol.** Let  $I = \int x^2 e^{x^3} \, dx = \frac{1}{3} \int e^{(x^3)} (3x^2) \, dx \quad \left[ \because \frac{d}{dx} x^3 = 3x^2 \right] \dots(i)$

Put  $x^3 = t$ . Therefore  $3x^2 = \frac{dt}{dx}$ . Therefore  $3x^2 \, dx = dt$

$$\therefore \text{ From (i), } I = \frac{1}{3} \int e^t dt = \frac{1}{3} e^t + C$$

$$\text{Putting } t = x^3, \quad = \frac{1}{3} e^{x^3} + C$$

$\therefore$  Option (B) is the correct answer.

**24.  $\int e^x \sec x (1 + \tan x) dx$  equals**

(A)  $e^x \cos x + C$

(B)  $e^x \sec x + C$

(C)  $e^x \sin x + C$

(D)  $e^x \tan x + C$

**Sol.** Let  $I = \int e^x \sec x (1 + \tan x) dx = \int e^x (\sec x + \sec x \tan x) dx$


It is of the form  $\int e^x (f(x) + f'(x)) dx$

Here  $f(x) = \sec x$  and so  $f'(x) = \sec x \tan x$

$$\therefore I = e^x f(x) + C \quad \left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) + C \right]$$

$$= e^x \sec x + C$$

$\therefore$  Option (B) is the correct answer.



## Exercise 7.7

**I. Rule to evaluate  $\int \sqrt{\text{Pure Quadratic}} \, dx$ , i.e.,  $\int \sqrt{ax^2 + b} \, dx$ .**

**Apply directly one of these formulae according to form of integrand:**

$$1. \int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}.$$

$$2. \int \sqrt{x^2 + a^2} \, dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log \left| x + \sqrt{x^2 + a^2} \right|.$$

$$3. \int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log \left| x + \sqrt{x^2 - a^2} \right|.$$

**II. Rule to evaluate  $\int \sqrt{\text{Quadratic}} \, dx$ , i.e.,  $\int \sqrt{ax^2 + bx + c} \, dx$**

**Step I.** Make coefficient of  $x^2$  unity by taking  $|a|$  common.

Now complete the squares by adding and subtracting

$$\left( \frac{1}{2} \text{Coefficient of } x \right)^2.$$

Now applying one of the above three formulae (according to the form of the integrand) will give value of required integral.

**Integrate the functions in Exercises 1 to 9:**

1.  $\sqrt{4 - x^2}$

**Sol.** 
$$\begin{aligned} \int \sqrt{4 - x^2} \, dx &= \int \sqrt{2^2 - x^2} \, dx \\ &= \frac{x}{2} \sqrt{2^2 - x^2} + \frac{2^2}{2} \sin^{-1} \frac{x}{2} + c \end{aligned}$$

$$\begin{aligned} \left[ \because \int \sqrt{a^2 - x^2} \, dx &= \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right] \\ &= \frac{x}{2} \sqrt{4 - x^2} + 2 \sin^{-1} \frac{x}{2} + c \end{aligned}$$

2.  $\sqrt{1 - 4x^2}$

**Sol.**  $\int \sqrt{1 - 4x^2} \, dx = \int \sqrt{1^2 - (2x)^2} \, dx$

$$= \frac{\frac{(2x)}{2} \sqrt{1^2 - (2x)^2} + \frac{1^2}{2} \sin^{-1} \left( \frac{2x}{1} \right)}{2 \rightarrow \text{Coefficient of } x \text{ in } 2x} + c$$

$$\left[ \because \int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]$$

$$= \frac{1}{2} \left[ x \sqrt{1 - 4x^2} + \frac{1}{2} \sin^{-1} \frac{2x}{1} \right] + c = \frac{x}{2} \sqrt{1 - 4x^2} + \frac{1}{4} \sin^{-1} 2x + c.$$

3.  $\sqrt{x^2 + 4x + 6}$

**Sol.**  $\int \sqrt{x^2 + 4x + 6} \, dx$

Coefficient of  $x^2$  is unity. So let us complete squares by adding and subtracting  $\left( \frac{1}{2} \text{Coefficient of } x \right)^2 = 2^2$

$$= \int \sqrt{x^2 + 4x + 4 + 6 - 4} \, dx = \int \sqrt{(x + 2)^2 + 2} \, dx$$

$$= \int \sqrt{(x + 2)^2 + (\sqrt{2})^2} \, dx = \left( \frac{x + 2}{2} \right) \sqrt{(x + 2)^2 + (\sqrt{2})^2}$$

$$+ \frac{(\sqrt{2})^2}{2} \log \left| x + 2 + \sqrt{(x + 2)^2 + (\sqrt{2})^2} \right| + c$$

$$\left[ \because \int \sqrt{x^2 + a^2} \, dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 + a^2}| \right]$$

$$= \frac{(x + 2)}{2} \sqrt{x^2 + 4x + 6}$$

$$+ \frac{2}{2} \log |x + 2 + \sqrt{x^2 + 4x + 6}| + c$$

$$= \frac{(x + 2)}{2} \sqrt{x^2 + 4x + 6} + \log |x + 2 + \sqrt{x^2 + 4x + 6}| + c.$$

4.  $\sqrt{x^2 + 4x + 1}$

**Sol.**  $\int \sqrt{x^2 + 4x + 1} \, dx = \int \sqrt{x^2 + 4x + 2^2 + 1 - 4} \, dx$

$$\left( \text{We have added and subtracted } \left( \frac{1}{2} \text{coefficient of } x \right)^2 = 2^2 \right)$$

$$\begin{aligned}
 &= \int \sqrt{(x+2)^2 - 3} \, dx = \int \sqrt{(x+2)^2 - (\sqrt{3})^2} \, dx \\
 &= \left( \frac{x+2}{2} \right) \sqrt{(x+2)^2 - (\sqrt{3})^2} \\
 &\quad - \frac{(\sqrt{3})^2}{2} \log \left| x+2 + \sqrt{(x+2)^2 - (\sqrt{3})^2} \right| + c \\
 &\quad \left[ \because \int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| \right] \\
 &= \left( \frac{x+2}{2} \right) \sqrt{x^2 + 4x + 1} - \frac{3}{2} \log \left| x+2 + \sqrt{x^2 + 4x + 1} \right| + c \\
 &\quad [\because (x+2)^2 - (\sqrt{3})^2 = x^2 + 4x + 4 - 3 = x^2 + 4x + 1]
 \end{aligned}$$

5.  $\int \sqrt{1 - 4x - x^2} \, dx$

**Sol.**  $\int \sqrt{1 - 4x - x^2} \, dx = \int \sqrt{-x^2 - 4x + 1} \, dx$

Making coefficient of  $x^2$  unity

$$= \int \sqrt{-(x^2 + 4x - 1)} \, dx$$

**(Note.** You can't take this  $(-)$  sign out of this bracket because square root of  $-1$  is imaginary)

$$= \int \sqrt{-(x^2 + 4x + 2^2 - 4 - 1)} \, dx = \int \sqrt{-(x+2)^2 - 5} \, dx$$

$$= \int \sqrt{5 - (x+2)^2} \, dx = \int \sqrt{(\sqrt{5})^2 - (x+2)^2} \, dx$$

$$= \frac{x+2}{2} \sqrt{(\sqrt{5})^2 - (x+2)^2} + \frac{(\sqrt{5})^2}{2} \sin^{-1} \left( \frac{x+2}{\sqrt{5}} \right) + c$$

$$\left[ \because \int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]$$

$$= \frac{x+2}{2} \sqrt{1 - 4x - x^2} + \frac{5}{2} \sin^{-1} \left( \frac{x+2}{\sqrt{5}} \right) + c$$

$$[\because (\sqrt{5})^2 - (x+2)^2 = 5 - (x^2 + 4x + 4x)]$$

$$= 5 - x^2 - 4 - 4x = 1 - 4x - x^2]$$

6.  $\int \sqrt{x^2 + 4x - 5} \, dx$

**Sol.**  $\int \sqrt{x^2 + 4x - 5} \, dx = \int \sqrt{x^2 + 4x + 2^2 - 4 - 5} \, dx$

$$= \int \sqrt{(x+2)^2 - 9} \, dx = \int \sqrt{(x+2)^2 - 3^2} \, dx$$

$$= \left( \frac{x+2}{2} \right) \sqrt{(x+2)^2 - 3^2} - \frac{3^2}{2} \log \left| x+2 + \sqrt{(x+2)^2 - 3^2} \right| + c$$

$$\left[ \because \int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| \right]$$

$$= \left( \frac{x+2}{2} \right) \sqrt{x^2+4x-5} - \frac{9}{2} \log \left| x+2+\sqrt{x^2+4x-5} \right| + c$$

$$[\because (x+2)^2 - 3^2 = x^2 + 4x + 4 - 9 = x^2 + 4x - 5]$$

7.  $\sqrt{1+3x-x^2}$

**Sol.**  $\int \sqrt{1+3x-x^2} \, dx = \int \sqrt{-x^2+3x+1} \, dx$

$$= \int \sqrt{-(x^2-3x-1)} \, dx$$

$$= \int \sqrt{-\left[x^2-3x+\left(\frac{3}{2}\right)^2 - \frac{9}{4} - 1\right]} \, dx = \int \sqrt{-\left[\left(x-\frac{3}{2}\right)^2 - \frac{13}{4}\right]} \, dx$$

$$= \int \sqrt{\frac{13}{4} - \left(x-\frac{3}{2}\right)^2} \, dx = \int \sqrt{\left(\frac{\sqrt{13}}{2}\right)^2 - \left(x-\frac{3}{2}\right)^2} \, dx$$

$$= \left(\frac{x-\frac{3}{2}}{2}\right) \sqrt{\left(\frac{\sqrt{13}}{2}\right)^2 - \left(x-\frac{3}{2}\right)^2} + \frac{\left(\frac{\sqrt{13}}{2}\right)^2}{2} \sin^{-1} \left(\frac{x-\frac{3}{2}}{\frac{\sqrt{13}}{2}}\right) + c$$

$$\left[ \because \int \sqrt{a^2-x^2} \, dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]$$

$$= \left(\frac{2x-3}{4}\right) \sqrt{1+3x-x^2} + \frac{13}{8} \sin^{-1} \left(\frac{2x-3}{\sqrt{13}}\right) + c$$

$$\left[ \because \left(\frac{\sqrt{13}}{2}\right)^2 - \left(x-\frac{3}{2}\right)^2 = \frac{13}{4} - \left(x^2 + \frac{9}{4} - 3x\right) \right]$$

$$= \frac{13}{4} - x^2 - \frac{9}{4} + 3x = 1 + 3x - x^2$$

8.  $\sqrt{x^2+3x}$

**Sol.**  $\int \sqrt{x^2+3x} \, dx = \int \sqrt{x^2+3x+\left(\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2} \, dx = \int \sqrt{\left(x+\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2} \, dx$

$$= \frac{x+\frac{3}{2}}{2} \sqrt{\left(x+\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2} - \frac{\left(\frac{3}{2}\right)^2}{2} \log \left| x+\frac{3}{2} + \sqrt{\left(x+\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2} \right| + c$$

$$\left[ \because \int \sqrt{x^2-a^2} \, dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \log |x+\sqrt{x^2-a^2}| \right]$$

$$= \frac{2x+3}{4} \sqrt{x^2+3x} - \frac{9}{8} \log \left| x + \frac{3}{2} + \sqrt{x^2+3x} \right| + c$$

$$\left[ \because \left( x + \frac{3}{2} \right)^2 - \left( \frac{3}{2} \right)^2 = x^2 + 3x + \frac{9}{4} - \frac{9}{4} = x^2 + 3x \right]$$

9.  $\sqrt{1+\frac{x^2}{9}}$

**Sol.**  $\int \sqrt{1+\frac{x^2}{9}} dx = \int \sqrt{\frac{9+x^2}{9}} dx = \int \frac{\sqrt{x^2+3^2}}{3} dx = \frac{1}{3} \int \sqrt{x^2+3^2} dx$

$$= \frac{1}{3} \left[ \frac{x}{2} \sqrt{x^2+3^2} + \frac{3^2}{2} \log \left| x + \sqrt{x^2+3^2} \right| \right] + c$$

$$\left[ \because \int \sqrt{x^2+a^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2+a^2}| \right]$$

$$= \frac{x}{6} \sqrt{x^2+9} + \frac{3}{2} \log |x + \sqrt{x^2+9}| + c.$$

**Choose the correct answer in Exercises 10 to 11:**

10.  $\int \sqrt{1+x^2} dx$  is equal to

- (A)  $\frac{x}{2} \sqrt{1+x^2} + \frac{1}{2} \log \left| \left( x + \sqrt{1+x^2} \right) \right| + C$
- (B)  $\frac{2}{3} (1+x^2)^{3/2} + C$  (C)  $\frac{2}{3} x (1+x^2)^{3/2} + C$
- (D)  $\frac{x^2}{2} \sqrt{1+x^2} + \frac{1}{2} x^2 \log \left| x + \sqrt{1+x^2} \right| + C.$

**Sol.**  $\int \sqrt{1+x^2} dx = \int \sqrt{x^2+1^2} dx$

$$= \frac{x}{2} \sqrt{x^2+1^2} + \frac{1^2}{2} \log |x + \sqrt{x^2+1^2}| + C$$

$$\left[ \because \int \sqrt{x^2+a^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2+a^2}| \right]$$

$$= \frac{x}{2} \sqrt{x^2+1} + \frac{1}{2} \log |x + \sqrt{x^2+1}| + C.$$

11.  $\int \sqrt{x^2-8x+7} dx$  is equal to

- (A)  $\frac{1}{2} (x-4) \sqrt{x^2-8x+7} + 9 \log \left| x-4 + \sqrt{x^2-8x+7} \right| + C$
- (B)  $\frac{1}{2} (x+4) \sqrt{x^2-8x+7} + 9 \log \left| x+4 + \sqrt{x^2-8x+7} \right| + C$
- (C)  $\frac{1}{2} (x-4) \sqrt{x^2-8x+7} - 3\sqrt{2} \log \left| x-4 + \sqrt{x^2-8x+7} \right| + C$
- (D)  $\frac{1}{2} (x-4) \sqrt{x^2-8x+7} - \frac{9}{2} \log \left| x-4 + \sqrt{x^2-8x+7} \right| + C.$

**Sol.**  $\int \sqrt{x^2 - 8x + 7} \, dx = \int \sqrt{x^2 - 8x + 4^2 - 16 + 7} \, dx$

$$= \int \sqrt{(x-4)^2 - 9} \, dx = \int \sqrt{(x-4)^2 - 3^2} \, dx$$

$$= \left( \frac{x-4}{2} \right) \sqrt{(x-4)^2 - 3^2} - \frac{3^2}{2} \log |x-4 + \sqrt{(x-4)^2 - 3^2}| + C$$

$$\left[ \because \int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| \right]$$

$$= \left( \frac{x-4}{2} \right) \sqrt{x^2 - 8x + 7} - \frac{9}{2} \log |x-4 + \sqrt{x^2 - 8x + 7}| + C.$$

[ $\because (x-4)^2 - 3^2 = x^2 - 8x + 16 - 9 = x^2 - 8x + 7$ ]



## Exercise 7.8

**Definition of definite integral as the limit of a sum:**

$$\int_a^b f(x) \, dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)]$$

where  $nh = b - a$

**Note.** The series within brackets represents the sum of  $n$  terms.

**Evaluate the following definite integrals as limit of sums:**

1.  $\int_a^b x \, dx$

**Sol. Step I.** Comparing  $\int_a^b x \, dx$  with  $\int_a^b f(x) \, dx$  we have

$$a = a, \, b = b \text{ and } f(x) = x \quad \dots(i)$$

$$\therefore nh = b - a = b - a$$

**Step II.** Putting  $x = a, a+h, a+2h, \dots, a+(n-1)h$  in (i), we have  $f(a) = a, f(a+h) = a+h,$

$$f(a+2h) = a+2h, \dots, f(a+(n-1)h) = a+(n-1)h$$

**Step III.** Putting these values in

$$\int_a^b f(x) \, dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)]$$

where  $nh = b - a$ , we have

$$\int_a^b x \, dx = \lim_{h \rightarrow 0} h [a + (a+h) + (a+2h) + \dots + (a+(n-1)h)]$$

where  $nh = b - a$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[na + h(1 + 2 + 3 + \dots + (n-1))]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ anh + hh \frac{n(n-1)}{2} \right] \left[ \because 1 + 2 + 3 + \dots + (n-1) = \frac{n(n-1)}{2} \right]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ anh + \frac{nh(nh-h)}{2} \right].$$

**Step IV.** Putting  $nh = b - a$ ,

$$= \lim_{h \rightarrow 0} \left[ a(b-a) + \frac{(b-a)(b-a-h)}{2} \right].$$

**Step V.** Taking Limits as  $h \rightarrow 0$  (i.e., putting  $h = 0$  here)

$$\begin{aligned} &= a(b-a) + \frac{(b-a)(b-a)}{2} \\ &= (b-a) \left[ a + \frac{b-a}{2} \right] = (b-a) \left[ \frac{2a+b-a}{2} \right] \\ &= \frac{(b-a)(b+a)}{2} = \frac{b^2 - a^2}{2}. \end{aligned}$$

2.  $\int_0^5 (x+1) dx$

**Sol. Step I.** Comparing  $\int_0^5 (x+1) dx$  with  $\int_a^b f(x) dx$ , we have

$$a = 0, b = 5 \text{ and } f(x) = x + 1 \quad \dots(i)$$

$$\therefore nh = b - a = 5 - 0 = 5.$$

**Step II.** Putting  $x = a, a + h, a + 2h, \dots, a + (n-1)h$  in (i), we have

$$f(a) = f(0) = 0 + 1 = 1, f(a+h) = f(h) = h + 1,$$

$$f(a+2h) = f(2h) = 2h + 1, \dots,$$

$$f(a+(n-1)h) = f((n-1)h) = (n-1)h + 1.$$

**Step III.** Putting these values in

$$\int_a^b f(x) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)], \text{ we have}$$

$$\int_0^5 (x+1) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [1 + (h+1) + (2h+1) + \dots + [(n-1)h+1]]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [n + h(1 + 2 + \dots + (n-1))] = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ nh + hh \frac{n(n-1)}{2} \right]$$

$$\left[ \because 1 + 2 + 3 + \dots + (n-1) = \frac{n(n-1)}{2} \right]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ nh + \frac{(nh)(nh-h)}{2} \right].$$

**Step IV.** Putting  $nh = 5$ ,  $= \lim_{h \rightarrow 0} \left[ 5 + \frac{5(5-h)}{2} \right]$ .

**Step V.** Taking limits as  $h \rightarrow 0$  (i.e., putting  $h = 0$  here)

$$= 5 + \frac{5(5-0)}{2} = 5 + \frac{25}{2} = \frac{10+25}{2} = \frac{35}{2}.$$

3.  $\int_2^3 x^2 dx$

**Sol. Step I.** Comparing  $\int_2^3 x^2 dx$  with  $\int_a^b f(x)$ , we have

$$a = 2, b = 3 \text{ and } f(x) = x^2 \quad \dots(i)$$

$$\therefore nh = b - a = 3 - 2 = 1.$$

**Step II.** Putting  $x = a, a + h, a + 2h, \dots, a + (n-1)h$  in (i), we have

$$f(a) = f(2) = 2^2 = 4$$

$$f(a + h) = f(2 + h) = (2 + h)^2 = 4 + 4h + h^2$$

$$f(a + 2h) = f(2 + 2h) = (2 + 2h)^2 = 4 + 8h + 2^2h^2$$

|

$$f(a + (n-1)h) = f(2 + (n-1)h) = (2 + (n-1)h)^2 \\ = 4 + 4(n-1)h + (n-1)^2h^2.$$

**Step III.** Putting these values in

$$\int_a^b f(x) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[f(a) + f(a+h) + f(a+2h) \\ + \dots + f(a+(n-1)h)]$$

where  $nh = 1$ , we have

$$\begin{aligned} \int_2^3 x^2 dx &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[4 + (4 + 4h + h^2) + (4 + 8h + 2^2h^2) \\ &\quad + \dots + (4 + 4(n-1)h + (n-1)^2h^2)] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[4n + 4h(1 + 2 + \dots + (n-1)) + h^2(1^2 + 2^2) \\ &\quad + \dots + (n-1)^2] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ 4nh + 4hh \frac{n(n-1)}{2} + hhh \frac{n(n-1)(2n-1)}{6} \right] \\ &\quad \left[ \because 1 + 2 + \dots + (n-1) = \frac{n(n-1)}{2} \text{ and } 1^2 + 2^2 + \dots \right. \\ &\quad \left. + (n-1)^2 = \frac{n(n-1)(2n-1)}{6} \right] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ 4nh + 4nh \frac{(nh-h)}{2} + \frac{nh(nh-h)(2nh-h)}{6} \right]. \end{aligned}$$

**Step IV.** Putting  $nh = 1$ ;

$$= \lim_{h \rightarrow 0} \left[ 4 + 2(1-h) + 1 \frac{(1-h)(2-h)}{6} \right].$$

**Step V.** Taking limits as  $h \rightarrow 0$  (i.e., putting  $h = 0$  here)

$$= 4 + 2(1 - 0) + \frac{1(2)}{6} = 6 + \frac{1}{3} = \frac{19}{3}.$$

4.  $\int_1^4 (x^2 - x) dx$

**Sol. Step I.** Comparing  $\int_1^4 (x^2 - x) dx$  with  $\int_a^b f(x) dx$ ,

we have

$$a = 1, b = 4, f(x) = x^2 - x \quad \dots(i)$$

$$\therefore nh = b - a = 4 - 1 = 3.$$

**Step II.** Putting  $x = a, a + h, a + 2h, \dots, a + (n - 1)h$  in (i),

$$f(a) = f(1) = 1^2 - 1 = 1 - 1 = 0$$

$$\begin{aligned} f(a + h) &= f(1 + h) = (1 + h)^2 - (1 + h) \\ &= 1 + h^2 + 2h - 1 - h = h + h^2 \end{aligned}$$

$$\begin{aligned} f(a + 2h) &= f(1 + 2h) = (1 + 2h)^2 - (1 + 2h) \\ &= 1 + 4h^2 + 4h - 1 - 2h \end{aligned}$$

$$\therefore \quad \quad \quad = 2h + 4h^2$$

$$\begin{aligned} f(a + (n - 1)h) &= (1 + (n - 1)h)^2 - (1 + (n - 1)h) \\ &= 1 + (n - 1)^2 h^2 + 2(n - 1)h - 1 - (n - 1)h \\ &= (n - 1)h + (n - 1)^2 h^2. \end{aligned}$$

**Step III.** Putting these values in

$$\int_a^b f(x) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[f(a) + f(a + h) + f(a + 2h) + \dots + f(a + (n - 1)h)]$$

we have

$$\begin{aligned} \int_1^4 (x^2 - x) dx &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[0 + h + h^2 + 2h + 4h^2 + \dots + (n - 1)h + (n - 1)^2 h^2] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h[h(1 + 2 + \dots + (n - 1)) + h^2(1^2 + 2^2 + \dots + (n - 1)^2)] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ h \cdot h \cdot \frac{n(n - 1)}{2} + h \cdot h \cdot h \cdot \frac{n(n - 1)(2n - 1)}{6} \right] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ nh \frac{(nh - h)}{2} + \frac{(nh)(nh - h)(2nh - h)}{6} \right]. \end{aligned}$$

**Step IV.** Putting  $nh = 3$

$$= \lim_{h \rightarrow 0} \left[ \frac{3(3 - h)}{2} + \frac{3(3 - h)(6 - h)}{6} \right].$$

**Step V.** Taking limits as  $h \rightarrow 0$  (Putting  $h = 0$  here)

$$= \frac{3(3 - 0)}{2} + \frac{3(3 - 0)(6 - 0)}{6} = \frac{9}{2} + 9 = \frac{27}{2}.$$

5.  $\int_{-1}^1 e^x dx$

**Sol. Step I.** Comparing  $\int_{-1}^1 e^x dx$  with  $\int_a^b f(x) dx$ , we have

$$a = -1, b = 1 \text{ and } f(x) = e^x \quad \dots(i)$$

$$\therefore nh = b - a = 1 - (-1) = 2.$$

**Step II.** Putting  $x = a, a + h, a + 2h, \dots, a + (n-1)h$  in (i), we have

$$f(a) = f(-1) = e^{-1}$$

$$f(a+h) = f(-1+h) = e^{-1+h} = e^{-1} \cdot e^h$$

$$f(a+2h) = f(-1+2h) = e^{-1+2h} = e^{-1} \cdot e^{2h}$$

⋮

⋮

$$f(a+(n-1)h) = f(-1+(n-1)h) = e^{-1+(n-1)h} = e^{-1} e^{(n-1)h}.$$

**Step III.** Putting these values in

$$\int_a^b f(x) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

we have

$$\begin{aligned} \int_{-1}^1 e^x dx &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [e^{-1} + e^{-1} e^h + e^{-1} e^{2h} + \dots + e^{-1} e^{(n-1)h}] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h e^{-1} \left[ \frac{(e^h)^n - 1}{e^h - 1} \right] \quad [\because \text{The series within brackets} \\ &\quad \text{is a G.P. series with First term } A = e^{-1} \text{ and common ratio } R = e^h, \\ &\quad \text{Number of terms is } n \text{ and } S_n \text{ of G.P.} = A \frac{(R^n - 1)}{R - 1}] \\ &= \int_{-1}^1 e^x dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h e^{-1} \frac{(e^{nh} - 1)}{e^h - 1}. \end{aligned}$$

**Step IV.** Putting  $nh = 2$ ,  $= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h e^{-1} \frac{(e^2 - 1)}{e^h - 1}$

$$\begin{aligned} &= e^{-1} (e^2 - 1) \lim_{h \rightarrow 0} \frac{h}{e^h - 1} = e^{-1} (e^2 - 1) \times 1 \left[ \because \lim_{x \rightarrow 0} \frac{x}{e^x - 1} = 1 \right] \\ &= e^{-1+2} - e^{-1} = e^1 - e^{-1} = e - e^{-1}. \end{aligned}$$

6.  $\int_0^4 (x + e^{2x}) dx$

**Sol. Step I.** Comparing  $\int_0^4 (x + e^{2x}) dx$  with  $\int_a^b f(x) dx$ , we have

$$a = 0, b = 4 \text{ and } f(x) = x + e^{2x} \quad \dots(i)$$

$$\therefore nh = b - a = 4 - 0 = 4.$$

**Step II.** Putting  $x = a, a + h, a + 2h, \dots, a + (n-1)h$  in (i), we have

$$f(a) = f(0) = 0 + e^0 = 1$$

$$\begin{aligned}
 f(a+h) &= f(h) = h + e^{2h} \\
 f(a+2h) &= f(2h) = 2h + e^{4h} \\
 &\vdots \\
 f(a+(n-1)h) &= f((n-1)h) = (n-1)h + e^{2(n-1)h}.
 \end{aligned}$$

**Step III.** Putting these values in

$$\int_a^b f(x) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

we have

$$\int_0^4 (x + e^{2x}) dx = \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [1 + (h + e^{2h}) + (2h + e^{4h}) + \dots + ((n-1)h + e^{2(n-1)h})]$$

(G.P. series : A = 1, R =  $e^{2h}$ ,  $n = n$ )

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h [(h + 2h + \dots + (n-1)h) + (1 + e^{2h} + e^{4h} + \dots + e^{2(n-1)h})]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h \left[ h(1 + 2 + \dots + (n-1)) + A \left( \frac{R^n - 1}{R - 1} \right) \right]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h \left[ h \frac{n(n-1)}{2} + \frac{1((e^{2h})^n - 1)}{e^{2h} - 1} \right]$$

$$= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} \left[ \frac{nh(nh - h)}{2} + \frac{h(e^{2nh} - 1)}{e^{2h} - 1} \right].$$

$$\text{Step IV. Putting } nh = 4, = \lim_{h \rightarrow 0} \left[ \frac{4(4-h)}{2} + \frac{h(e^8 - 1)}{e^{2h} - 1} \right].$$

**Step V.** Taking limits as  $h \rightarrow 0$

$$= \frac{4(4-0)}{2} + (e^8 - 1) \lim_{h \rightarrow 0} \frac{h}{e^{2h} - 1} = 8 + (e^8 - 1) \frac{1}{2} \lim_{h \rightarrow 0} \frac{2h}{e^{2h} - 1}$$

$$= 8 + \frac{(e^8 - 1)}{2} \left[ \because \lim_{h \rightarrow 0} \frac{e^{2h}}{e^{2h} - 1} \left( \Rightarrow \lim_{x \rightarrow 0} \frac{x}{e^x - 1} \right) = 1 \right]$$



### Exercise 7.9

Evaluate the definite integrals in Exercises 1 to 11:

**Result.** If  $\int f(x) \, dx = \phi(x)$ , then  $\int_a^b f(x) \, dx = \phi(b) - \phi(a) \quad \dots(i)$

(This is known as **Second Fundamental Theorem**).

1.  $\int_{-1}^1 (x+1) \, dx$

**Sol.**  $\int_{-1}^1 (x+1) \, dx = \left( \frac{x^2}{2} + x \right)_{-1}^1 = \phi(b) - \phi(a)$

(By Second Fundamental Theorem given in Eqn. (i) page 496)

$$= \left( \frac{1^2}{2} + 1 \right) - \left( \frac{(-1)^2}{2} - 1 \right) = \frac{1}{2} + 1 - \left( \frac{1}{2} - 1 \right)$$

$$= \frac{1}{2} + 1 - \frac{1}{2} + 1 = 2.$$

**Remark.** [Constant  $c$  will never occur in the value of a definite integral because  $c$  in the value of  $\phi(b)$  gets cancelled with  $c$  in  $\phi(a)$  when we subtract them to get  $\phi(b) - \phi(a)$ ].

2.  $\int_2^3 \frac{1}{x} \, dx$

**Sol.**  $\int_2^3 \frac{1}{x} \, dx = (\log |x|)_2^3 = \phi(b) - \phi(a) = \log |3| - \log |2|$

$$= \log 3 - \log 2 = \log \frac{3}{2}. \quad [\because |x| = x \text{ if } x \geq 0]$$

3.  $\int_1^2 (4x^3 - 5x^2 + 6x + 9) \, dx$

**Sol.**  $\int_1^2 (4x^3 - 5x^2 + 6x + 9) \, dx = \left( 4 \frac{x^4}{4} - 5 \frac{x^3}{3} + 6 \frac{x^2}{2} + 9x \right)_1^2$

$$= \left( x^4 - \frac{5}{3}x^3 + 3x^2 + 9x \right)_1^2$$

$$= \left[ 2^4 - \frac{5}{3}(2)^3 + 3(2)^2 + 9(2) \right] - \left[ 1 - \frac{5}{3} + 3 + 9 \right]$$

$$= \left( 16 - \frac{40}{3} + 12 + 18 \right) - \left( 13 - \frac{5}{3} \right)$$

$$= 46 - \frac{40}{3} - \left( 13 - \frac{5}{3} \right) = 46 - \frac{40}{3} - 13 + \frac{5}{3}$$

$$= 33 - \frac{40}{3} + \frac{5}{3} = \frac{99 - 40 + 5}{3} = \frac{104 - 40}{3} = \frac{64}{3}.$$

4.  $\int_0^{\frac{\pi}{4}} \sin 2x \, dx$

**Sol.**  $\int_0^{\frac{\pi}{4}} \sin 2x \, dx = \left( \frac{-\cos 2x}{2} \right)_0^{\frac{\pi}{4}} = \frac{-\cos \frac{\pi}{2}}{2} - \left( \frac{-\cos 0}{2} \right)$

$$= \frac{-0}{2} - \left( \frac{-1}{2} \right) = 0 + \frac{1}{2} = \frac{1}{2}.$$

5.  $\int_0^{\frac{\pi}{2}} \cos 2x \, dx$

**Sol.**  $\int_0^{\frac{\pi}{2}} \cos 2x \, dx = \left( \frac{\sin 2x}{2} \right)_0^{\frac{\pi}{2}} = \frac{\sin \pi}{2} - \frac{\sin 0}{2}$   
 $= \frac{0}{2} - \frac{0}{2} = 0$   
 $[\because \sin \pi = \sin 180^\circ = \sin (180^\circ - 0^\circ) = \sin 0 = 0]$

6.  $\int_4^5 e^x \, dx$

**Sol.**  $\int_4^5 e^x \, dx = (e^x)_4^5 = e^5 - e^4 = e^4 (e - 1).$

7.  $\int_0^{\frac{\pi}{4}} \tan x \, dx$

**Sol.**  $\int_0^{\frac{\pi}{4}} \tan x \, dx = (\log |\sec x|)_0^{\frac{\pi}{4}}$   
 $= \log \left| \sec \frac{\pi}{4} \right| - \log |\sec 0| = \log |\sqrt{2}| - \log |1|$   
 $= \log \sqrt{2} - \log 1 = \log 2^{1/2} - 0 = \frac{1}{2} \log 2.$

8.  $\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \operatorname{cosec} x \, dx$

**Sol.**  $\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \operatorname{cosec} x \, dx = (\log |\operatorname{cosec} x - \cot x|)_{\frac{\pi}{6}}^{\frac{\pi}{4}}$   
 $= \log \left| \operatorname{cosec} \frac{\pi}{4} - \cot \frac{\pi}{4} \right| - \log \left| \operatorname{cosec} \frac{\pi}{6} - \cot \frac{\pi}{6} \right|$   
 $= \log |\sqrt{2} - 1| - \log |2 - \sqrt{3}|$   
 $= \log (\sqrt{2} - 1) - \log (2 - \sqrt{3}) \quad [\because |x| = x \text{ if } x \geq 0]$   
 $= \log \left( \frac{\sqrt{2} - 1}{2 - \sqrt{3}} \right).$

9.  $\int_0^1 \frac{dx}{\sqrt{1-x^2}}$

**Sol.**  $\int_0^1 \frac{dx}{\sqrt{1-x^2}} = (\sin^{-1} x)_0^1 \quad \left[ \because \int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1} \frac{x}{a} \right]$   
 $= \sin^{-1} 1 - \sin^{-1} 0 = \frac{\pi}{2} - 0 = \frac{\pi}{2}. \quad \left[ \because \sin \frac{\pi}{2} = 1 \text{ and } \sin 0 = 0 \right]$

10.  $\int_0^1 \frac{dx}{1+x^2}$

$$\begin{aligned}
 \text{Sol. } \int_0^1 \frac{dx}{1+x^2} &= \left( \tan^{-1} x \right)_0^1 & \left[ \because \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right] \\
 &= \tan^{-1} 1 - \tan^{-1} 0 = \frac{\pi}{4} - 0 = \frac{\pi}{4}. \\
 & \left[ \because \tan \frac{\pi}{4} = 1 \text{ and } \tan 0 = 0 \right]
 \end{aligned}$$

$$11. \int_2^3 \frac{dx}{x^2-1}$$

$$\begin{aligned}
 \text{Sol. } \int_2^3 \frac{1}{x^2-1} dx &= \int_2^3 \frac{1}{x^2-1^2} dx \\
 &= \left( \frac{1}{2(1)} \log \left| \frac{x-1}{x+1} \right| \right)_2^3 \left[ \because \int \frac{1}{x^2-a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| \right] \\
 &= \frac{1}{2} \log \left| \frac{3-1}{3+1} \right| - \frac{1}{2} \log \left| \frac{2-1}{2+1} \right| = \frac{1}{2} \log \left| \frac{1}{2} \right| - \frac{1}{2} \log \left| \frac{1}{3} \right| \\
 &= \frac{1}{2} \left( \log \frac{1}{2} - \log \frac{1}{3} \right) \quad [\because |x| = x \text{ if } x \geq 0] \\
 &= \frac{1}{2} \left[ \log \left( \frac{1}{\frac{1}{3}} \right) \right] = \frac{1}{2} \log \frac{3}{2}.
 \end{aligned}$$

**Evaluate the definite integrals in Exercises 12 to 20:**

$$12. \int_0^{\frac{\pi}{2}} \cos^2 x \, dx$$

$$\begin{aligned}
 \text{Sol. } \int_0^{\frac{\pi}{2}} \cos^2 x \, dx &= \int_0^{\frac{\pi}{2}} \frac{1+\cos 2x}{2} dx = \int_0^{\frac{\pi}{2}} \frac{1}{2} (1+\cos 2x) dx \\
 &= \frac{1}{2} \int_0^{\frac{\pi}{2}} (1+\cos 2x) dx = \frac{1}{2} \left( x + \frac{\sin 2x}{2} \right)_0^{\frac{\pi}{2}} \\
 &= \frac{1}{2} \left[ \frac{\pi}{2} + \frac{1}{2} \sin \pi - \left( 0 + \frac{1}{2} \sin 0 \right) \right] = \frac{1}{2} \left[ \frac{\pi}{2} + 0 - 0 \right] \\
 &= \frac{\pi}{4}. \quad [\because \sin \pi = \sin 180^\circ = \sin (180^\circ - 0^\circ) = \sin 0 = 0]
 \end{aligned}$$

$$13. \int_2^3 \frac{x \, dx}{x^2+1}$$

$$\begin{aligned}
 \text{Sol. } \int_2^3 \frac{x}{x^2+1} dx &= \frac{1}{2} \int_2^3 \frac{2x}{x^2+1} dx \\
 &= \frac{1}{2} \left( \log |x^2+1| \right)_2^3. & \left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right] \\
 & \text{(Here } f(x) = x^2+1 \text{ and } f'(x) = 2x)
 \end{aligned}$$

$$= \frac{1}{2} (\log |10| - \log |5|) = \frac{1}{2} (\log 10 - \log 5)$$

$$= \frac{1}{2} \log \frac{10}{5} = \frac{1}{2} \log 2.$$

14.  $\int_0^1 \frac{2x+3}{5x^2+1} dx$

**Sol.**  $\int_0^1 \frac{2x+3}{5x^2+1} dx = \int_0^1 \left( \frac{2x}{5x^2+1} + \frac{3}{5x^2+1} \right) dx$

$$= \int_0^1 \frac{2x}{5x^2+1} dx + 3 \int_0^1 \frac{dx}{5x^2+1}$$

$$= \frac{1}{5} \int_0^1 \frac{10x}{5x^2+1} dx + 3 \int_0^1 \frac{dx}{(\sqrt{5}x)^2+1^2}$$

$$= \frac{1}{5} (\log |5x^2+1|)_0^1 + 3 \cdot \frac{1}{1} \cdot \frac{\left( \tan^{-1} \left( \frac{\sqrt{5}x}{1} \right) \right)_0^1}{\sqrt{5} \rightarrow \text{Coefficient of } x}$$

$$\left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \text{ and } \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right]$$

$$= \frac{1}{5} (\log 6 - \log 1) + \frac{3}{\sqrt{5}} (\tan^{-1} \sqrt{5} - \tan^{-1} 0)$$

$$= \frac{1}{5} \log 6 + \frac{3}{\sqrt{5}} \tan^{-1} \sqrt{5}.$$

15.  $\int_0^1 x e^{x^2} dx$

**Sol.** To evaluate  $\int_0^1 x e^{x^2} dx$

Let us first evaluate  $\int x e^{x^2} dx$

$$= \frac{1}{2} \int e^{x^2} (2x dx) \quad \dots(i)$$

Put  $x^2 = t$ . Therefore  $2x = \frac{dt}{dx} \therefore 2x dx = dt$

$$\therefore \text{ From (i), } \int x e^{x^2} dx = \frac{1}{2} \int e^t dt = \frac{1}{2} e^t$$

**Putting  $t = x^2$ ,**  $= \frac{1}{2} e^{x^2} \quad \dots(ii)$

$$\therefore \text{ The given integral } \int_0^1 x e^{x^2} dx = \frac{1}{2} (e^{x^2})_0^1 \quad [\text{By (ii)}]$$

$$= \frac{1}{2} (e^1 - e^0) = \frac{1}{2} (e - 1).$$

**Note.** Please note that limits 0 and 1 specified in the given integral are limits for  $x$ .

Therefore after substituting  $x^2 = t$  and evaluating the integral, we must put back  $t = x^2$  and only then use  $\int_a^b f(x) dx = \phi(b) - \phi(a)$ .

**Remark.** In the next Exercise 7.10 we shall also learn to change the limits of integration from values of  $x$  to values of  $t$  and then we may use our discretion even here also.

16.  $\int_1^2 \frac{5x^2}{x^2 + 4x + 3} dx$

**Sol.**  $\int_1^2 \frac{5x^2}{x^2 + 4x + 3} dx = \int_1^2 \frac{5x^2}{(x+1)(x+3)} \dots(i)$   
 $[\because x^2 + 4x + 3 = x^2 + 3x + x + 3$   
 $= x(x+3) + 1(x+3) = (x+1)(x+3)]$

The integrand  $\frac{5x^2}{(x+1)(x+3)}$  is a rational function and degree of numerator = degree of denominator.

So let us apply long division.

$$\begin{array}{r} (x+1)(x+3) = x^2 + 4x + 3 \overline{) 5x^2} \quad (5 \\ \underline{5x^2 + 20x + 15} \\ -20x - 15 \end{array}$$

$$\therefore \frac{5x^2}{(x+1)(x+3)} = 5 + \frac{(-20x-15)}{(x+1)(x+3)}$$

Putting this value in (i),

$$\begin{aligned} \int_1^2 \frac{5x^2}{x^2 + 4x + 3} dx &= \int_1^2 \left( 5 + \frac{(-20x-15)}{(x+1)(x+3)} \right) dx \\ &= \int_1^2 5 dx + \int_1^2 \frac{-20x-15}{(x+1)(x+3)} dx = 5(x)_1^2 + I \\ &= 5(2-1) + I = 5 + I \end{aligned} \dots(ii)$$

where  $I = \int_1^2 \frac{-20x-15}{(x+1)(x+3)} dx$

Let integrand of  $I = \frac{-20x-15}{(x+1)(x+3)} = \frac{A}{x+1} + \frac{B}{x+3} \dots(iii)$   
(Partial Fractions)

Multiplying both sides by L.C.M. =  $(x+1)(x+3)$ ,  
 $-20x - 15 = A(x+3) + B(x+1)$   
 $= Ax + 3A + Bx + B$

Comparing coefficients of  $x$  and constant terms on both sides, we have

**Coefficients of  $x$ :**  $A + B = -20 \dots(iv)$

**Constant terms:**  $3A + B = -15 \dots(v)$

Subtracting (iv) and (v),  $-2A = -5$ . Therefore  $A = \frac{5}{2}$ .

Putting  $A = \frac{5}{2}$  in (iv),  $\frac{5}{2} + B = -20 \Rightarrow B = -20 - \frac{5}{2}$

$$\text{or } B = \frac{-40-5}{2} = \frac{-45}{2}$$

Putting these values of A and B in (iii),

$$\frac{-20x-15}{(x+1)(x+3)} = \frac{\frac{5}{2}}{x+1} - \frac{\frac{45}{2}}{x+3}$$

$$\begin{aligned} \therefore I &= \int_1^2 \frac{-20x-15}{(x+1)(x+3)} dx = \frac{5}{2} \int_1^2 \frac{1}{x+1} dx - \frac{45}{2} \int_1^2 \frac{1}{x+3} dx \\ &= \frac{5}{2} (\log |x+1|)_1^2 - \frac{45}{2} (\log |x+3|)_1^2 \\ &= \frac{5}{2} (\log |3| - \log |2|) - \frac{45}{2} (\log |5| - \log |4|) \\ &= \frac{5}{2} \log \frac{3}{2} - \frac{45}{2} \log \frac{5}{4} \quad [\because |x| = x \text{ if } x \geq 0] \\ &= \frac{5}{2} \left( \log \frac{3}{2} - 9 \log \frac{5}{4} \right) \end{aligned}$$

Putting this value of I in (ii),

$$\int_1^2 \frac{5x^2}{x^2+4x+3} dx = 5 + \frac{5}{2} \left( \log \frac{3}{2} - 9 \log \frac{5}{4} \right) = 5 - \frac{5}{2} \left( 9 \log \frac{5}{4} - \log \frac{3}{2} \right)$$

$$17. \int_0^{\frac{\pi}{4}} (2 \sec^2 x + x^3 + 2) dx$$

$$\begin{aligned} \text{Sol. } \int_0^{\frac{\pi}{4}} (2 \sec^2 x + x^3 + 2) dx &= 2 \int_0^{\frac{\pi}{4}} \sec^2 x dx + \int_0^{\frac{\pi}{4}} x^3 dx + 2 \int_0^{\frac{\pi}{4}} 1 dx \\ &= 2 \left( \tan x \right)_0^{\frac{\pi}{4}} + \left( \frac{x^4}{4} \right)_0^{\frac{\pi}{4}} + 2 \left( x \right)_0^{\frac{\pi}{4}} \\ &= 2 \left( \tan \frac{\pi}{4} - \tan 0 \right) + \frac{\left( \frac{\pi}{4} \right)^4}{4} - 0 + 2 \left( \frac{\pi}{4} - 0 \right) \\ &= 2(1-0) + \frac{\left( \frac{\pi^4}{256} \right)}{4} + \frac{2\pi}{4} = 2 + \frac{\pi^4}{1024} + \frac{\pi}{2} . \end{aligned}$$

$$18. \int_0^{\pi} \left( \sin^2 \frac{x}{2} - \cos^2 \frac{x}{2} \right) dx$$

$$\begin{aligned} \text{Sol. } \int_0^{\pi} \left( \sin^2 \frac{x}{2} - \cos^2 \frac{x}{2} \right) dx &= \int_0^{\pi} \left[ \left( \frac{1-\cos x}{2} \right) - \left( \frac{1+\cos x}{2} \right) \right] dx \\ &\left( \because \sin^2 \theta = \frac{1-\cos 2\theta}{2} \text{ and } \cos^2 \theta = \frac{1+\cos 2\theta}{2} \right) \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\pi} \left( \frac{1 - \cos x - 1 - \cos x}{2} \right) dx = \int_0^{\pi} \frac{-2 \cos x}{2} dx \\
 &= - \int_0^{\pi} \cos x \, dx = - (\sin x)_0^{\pi} = - (\sin \pi - \sin 0) = - (0 - 0) = 0. \\
 &\quad [\because \sin \pi = \sin 180^\circ = \sin (180^\circ - 0) = \sin 0 = 0]
 \end{aligned}$$

19.  $\int_0^2 \frac{6x+3}{x^2+4} dx$

**Sol.**  $\int_0^2 \frac{6x+3}{x^2+4} dx = \int_0^2 \frac{6x}{x^2+4} dx + 3 \int_0^2 \frac{1}{x^2+4} dx$

$$\begin{aligned}
 &= 3 \int_0^2 \frac{2x}{x^2+4} dx + 3 \frac{1}{2} \left( \tan^{-1} \frac{x}{2} \right)_0^2 \\
 &= 3 \left( \log |x^2+4| \right)_0^2 + \frac{3}{2} (\tan^{-1} 1 - \tan^{-1} 0) \\
 &\quad \left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \text{ and } \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right] \\
 &= 3 (\log 8 - \log 4) + \frac{3}{2} \left( \frac{\pi}{4} - 0 \right) \quad \left[ \because \tan \frac{\pi}{4} = 1 \right] \\
 &= 3 \log \frac{8}{4} + \frac{3\pi}{8} = 3 \log 2 + \frac{3\pi}{8}.
 \end{aligned}$$

20.  $\int_0^1 \left( x e^x + \sin \frac{\pi x}{4} \right) dx$

**Sol.**  $\int_0^1 \left( x e^x + \sin \frac{\pi x}{4} \right) dx = \int_0^1 \underset{\text{I}}{x} \underset{\text{II}}{e^x} dx + \int_0^1 \sin \frac{\pi x}{4} dx$

Applying Product Rule on first definite integral,

$$\begin{aligned}
 &\left( \text{I} \int \text{II} dx \right)_0^1 - \int_0^1 \left( \frac{d}{dx} (\text{I}) \int \text{II} dx \right) dx \\
 &= \left( x e^x \right)_0^1 - \int_0^1 1 \cdot e^x dx - \frac{\left( \cos \frac{\pi x}{4} \right)_0^1}{\frac{\pi}{4} \rightarrow \text{Coefficient of } x \text{ in } \frac{\pi x}{4}} \\
 &= e^1 - 0 - \int_0^1 e^x dx - \frac{4}{\pi} \left[ \cos \frac{\pi}{4} - \cos 0 \right] = e - \left( e^x \right)_0^1 - \frac{4}{\pi} \left( \frac{1}{\sqrt{2}} - 1 \right) \\
 &= e - (e - e^0) - \frac{4}{\pi\sqrt{2}} + \frac{4}{\pi} \\
 &= e - e + 1 - \frac{2 \cdot 2}{\pi\sqrt{2}} + \frac{4}{\pi} = 1 + \frac{4}{\pi} - \frac{2\sqrt{2}}{\pi}.
 \end{aligned}$$

Choose the correct answer in Exercises 21 and 22:

21.  $\int_1^{\sqrt{3}} \frac{dx}{1+x^2}$  equals

- (A)  $\frac{\pi}{3}$  (B)  $\frac{2\pi}{3}$  (C)  $\frac{\pi}{6}$  (D)  $\frac{\pi}{12}$

Sol.  $\int_1^{\sqrt{3}} \frac{dx}{1+x^2} = \left( \tan^{-1} x \right)_1^{\sqrt{3}} = \tan^{-1} \sqrt{3} - \tan^{-1} 1$   
 $= \frac{\pi}{3} - \frac{\pi}{4} \quad \left[ \because \tan \frac{\pi}{3} = \sqrt{3} \text{ and } \tan \frac{\pi}{4} = 1 \right]$   
 $= \frac{4\pi - 3\pi}{12} = \frac{\pi}{12}$

$\therefore$  Option (D) is the correct answer.

22.  $\int_0^{\frac{2}{3}} \frac{dx}{4+9x^2}$  equals

- (A)  $\frac{\pi}{6}$  (B)  $\frac{\pi}{12}$  (C)  $\frac{\pi}{24}$  (D)  $\frac{\pi}{4}$

Sol.  $\int_0^{\frac{2}{3}} \frac{dx}{4+9x^2} = \int_0^{\frac{2}{3}} \frac{dx}{(3x)^2 + 2^2} = \left[ \frac{1}{2} \frac{\tan^{-1} \frac{3x}{2}}{3 \rightarrow \text{Coefficient of } x \text{ in } 3x} \right]$   
 $= \frac{1}{6} \left[ \tan^{-1} \frac{3x}{2} \right]_0^{\frac{2}{3}} = \frac{1}{6} \left[ \tan^{-1} \left( \frac{3}{2} \times \frac{2}{3} \right) - \tan^{-1} 0 \right]$   
 $= \frac{1}{6} (\tan^{-1} 1 - \tan^{-1} 0) = \frac{1}{6} \left( \frac{\pi}{4} - 0 \right) = \frac{\pi}{24}$   
 $\left[ \because \tan \frac{\pi}{4} = 1 \text{ and } \tan 0 = 0 \right]$

$\therefore$  Option (C) is the correct answer.

### Exercise 7.10

Evaluate the integrals in Exercises 1 to 8 using substitution:

1.  $\int_0^1 \frac{x}{x^2+1} dx$

**Sol.** Let  $I = \int_0^1 \frac{x}{x^2+1} dx = \frac{1}{2} \int_0^1 \frac{2x}{x^2+1} dx \quad \dots(i)$

Put  $x^2 + 1 = t$ . Therefore  $2x = \frac{dt}{dx} \Rightarrow 2x dx = dt$ .

**To change the limits of integration from values of  $x$  to values of  $t$ .**

When  $x = 0$ ,  $t = 0 + 1 = 1$

When  $x = 1$ ,  $t = 1 + 1 = 2$

$\therefore$  From (i),  $I = \frac{1}{2} \int_1^2 \frac{dt}{t} = \frac{1}{2} (\log |t|)_1^2 = \frac{1}{2} (\log |2| - \log |1|)$

$$= \frac{1}{2} (\log 2 - \log 1) = \frac{1}{2} (\log 2 - 0) = \frac{1}{2} \log 2.$$

2.  $\int_0^{\frac{\pi}{2}} \sqrt{\sin \phi} \cos^5 \phi \, d\phi$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \sqrt{\sin \phi} \cos^5 \phi \, d\phi$  ... (i)

**Put  $\sin \phi = t$ .**

( $\because$  one factor of integrand is  $\cos^5 \phi$  where  $n = 5$  is odd.)

$$\therefore \cos \phi = \frac{dt}{d\phi} \quad \text{i.e.,} \quad \cos \phi \, d\phi = dt.$$

**To change the limits of integration from  $\phi$  to  $t$**

When  $\phi = 0$ ,  $t = \sin \phi = \sin 0 = 0$

When  $\phi = \frac{\pi}{2}$ ,  $t = \sin \phi = \sin \frac{\pi}{2} = 1$

$$\begin{aligned} \text{Now Integrand } \sqrt{\sin \phi} \cos^5 \phi &= \sqrt{\sin \phi} \cos^4 \phi \cos \phi \\ &= \sqrt{\sin \phi} (\cos^2 \phi)^2 \cos \phi = \sqrt{\sin \phi} (1 - \sin^2 \phi)^2 \cos \phi \end{aligned}$$

$$\begin{aligned} \therefore \text{ From (i), } I &= \int_0^{\frac{\pi}{2}} \sqrt{\sin \phi} (1 - \sin^2 \phi)^2 \cos \phi \, d\phi \\ &= \int_0^1 \sqrt{t} (1 - t^2)^2 \, dt = \int_0^1 t^{1/2} (1 + t^4 - 2t^2) \, dt \\ &= \int_0^1 \left( t^{1/2} + t^{9/2} - 2t^{5/2} \right) \, dt = \int_0^1 (t^{1/2} + t^{9/2} - 2t^{5/2}) \, dt \\ &= \int_0^1 t^{1/2} \, dt + \int_0^1 t^{9/2} \, dt - 2 \int_0^1 t^{5/2} \, dt \\ &= \frac{(t^{3/2})_0^1}{\frac{3}{2}} + \frac{(t^{11/2})_0^1}{\frac{11}{2}} - 2 \frac{(t^{7/2})_0^1}{\frac{7}{2}} \\ &= \frac{2}{3} (1 - 0) + \frac{2}{11} (1 - 0) - \frac{4}{7} (1 - 0) \\ &= \frac{2}{3} + \frac{2}{11} - \frac{4}{7} = \frac{2(77) + 2(21) - 4(33)}{3(11)(7)} \\ &= \frac{154 + 42 - 132}{231} = \frac{196 - 132}{231} = \frac{64}{231}. \end{aligned}$$

3.  $\int_0^1 \sin^{-1} \left( \frac{2x}{1+x^2} \right) dx$

**Sol.** Let  $I = \int_0^1 \sin^{-1} \left( \frac{2x}{1+x^2} \right) dx$  ... (i)

**Put  $x = \tan \theta$ .**  $\therefore \frac{dx}{d\theta} = \sec^2 \theta \Rightarrow dx = \sec^2 \theta \, d\theta$

**To change the limits of integration**

When  $x = 0$ ,  $\tan \theta = 0 = \tan 0 \Rightarrow \theta = 0$

When  $x = 1$ ,  $\tan \theta = 1 = \tan \frac{\pi}{4} \Rightarrow \theta = \frac{\pi}{4}$

$$\therefore \text{ From (i), } I = \int_0^{\frac{\pi}{4}} \left( \sin^{-1} \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right) \right) \sec^2 \theta \, d\theta$$

$$= \int_0^{\frac{\pi}{4}} (\sin^{-1} (\sin 2\theta)) \sec^2 \theta \, d\theta = \int_0^{\frac{\pi}{4}} 2\theta \sec^2 \theta \, d\theta$$

$$= 2 \int_0^{\frac{\pi}{4}} \theta \sec^2 \theta \, d\theta$$

I I I

Applying Product Rule of Integration

$$\begin{aligned} \left( \int_a^b I \cdot II \, dx = (I \int_a^b II) - \int_a^b \left( \frac{d}{dx} (I) \int_a^b II \, dx \right) dx \right) \\ = 2 \left[ (\theta \cdot \tan \theta)_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} 1 \cdot \tan \theta \, d\theta \right] \\ = 2 \left[ \frac{\pi}{4} \tan \frac{\pi}{4} - 0 - \int_0^{\frac{\pi}{4}} \tan \theta \, d\theta \right] = 2 \left[ \frac{\pi}{4} - (\log \sec \theta)_0^{\frac{\pi}{4}} \right] \\ = 2 \left[ \frac{\pi}{4} - \left( \log \sec \frac{\pi}{4} - \log \sec 0 \right) \right] = 2 \left[ \frac{\pi}{4} - (\log \sqrt{2} - \log 1) \right] \\ = \frac{\pi}{2} - 2 \log 2^{1/2} \quad (\because \log 1 = 0) \\ = \frac{\pi}{2} - 2 \cdot \frac{1}{2} \log 2 = \frac{\pi}{2} - \log 2. \end{aligned}$$

4.  $\int_0^2 x\sqrt{x+2} \, dx$

**Sol.** Let  $I = \int_0^2 x\sqrt{x+2} \, dx$  ... (i)

Put  $\sqrt{x+2} = t$ , i.e.,  $\sqrt{x+2} = t$ . Therefore  $x+2 = t^2$ .

$$\therefore \frac{dx}{dt} = 2t \Rightarrow dx = 2t \, dt$$

**To change the limits of Integration**

When  $x = 0$ ,  $t = \sqrt{x+2} = \sqrt{2}$

When  $x = 2$ ,  $t = \sqrt{x+2} = \sqrt{2+2} = \sqrt{4} = 2$ .

$$\therefore \text{ From (i), } I = \int_{\sqrt{2}}^2 (t^2 - 2) \cdot 2t \, dt$$

$$[\because x+2 = t^2 \Rightarrow x = t^2 - 2]$$

$$= 2 \int_{\sqrt{2}}^2 t^2(t^2 - 2) \, dt = 2 \int_{\sqrt{2}}^2 (t^4 - 2t^2) \, dt$$

$$= 2 \left[ \left( \frac{t^5}{5} \right)_{\sqrt{2}}^2 - 2 \left( \frac{t^3}{3} \right)_{\sqrt{2}}^2 \right] = 2 \left[ \frac{1}{5} (2^5 - (\sqrt{2})^5) - \frac{2}{3} (2^3 - (\sqrt{2})^3) \right]$$

$$\begin{aligned}
 &= 2 \left[ \frac{1}{5} (32 - 4\sqrt{2}) - \frac{2}{3} (8 - 2\sqrt{2}) \right] \left[ \because (\sqrt{2})^3 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 2\sqrt{2}, \right. \\
 &\quad \left. \text{and } (\sqrt{2})^5 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 4\sqrt{2} \right] \\
 &= 2 \left[ \frac{32}{5} - \frac{4\sqrt{2}}{5} - \frac{16}{3} + \frac{4\sqrt{2}}{3} \right] = 2 \left[ \frac{96 - 12\sqrt{2} - 80 + 20\sqrt{2}}{15} \right] \\
 &= \frac{2}{15} (16 + 8\sqrt{2}) = \frac{16}{15} (2 + \sqrt{2}) = \frac{16}{15} (\sqrt{2} \cdot \sqrt{2} + \sqrt{2}) \\
 &= \frac{16\sqrt{2}}{15} (\sqrt{2} + 1).
 \end{aligned}$$

5.  $\int_0^{\frac{\pi}{2}} \frac{\sin x}{1 + \cos^2 x} dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\sin x}{1 + \cos^2 x} dx = - \int_0^{\frac{\pi}{2}} \frac{-\sin x}{1 + \cos^2 x} dx \dots (i)$

**Put  $\cos x = t$ .** Therefore  $-\sin x = \frac{dt}{dx} \Rightarrow -\sin x dx = dt$ .

**To change the limits of Integration.**

When  $x = 0$ ,  $t = \cos 0 = 1$ , When  $x = \frac{\pi}{2}$ ,  $t = \cos \frac{\pi}{2} = 0$

$$\begin{aligned}
 \therefore \text{ From (i), } I &= - \int_1^0 \frac{dt}{1+t^2} = - \int_1^0 \frac{1}{t^2+1} dt \\
 &= - \left( \tan^{-1} t \right)_1^0 = - (\tan^{-1} 0 - \tan^{-1} 1) = - \left( 0 - \frac{\pi}{4} \right) \\
 &\left[ \because \tan 0 = 0 \Rightarrow \tan^{-1} 0 = 0 \text{ and } \tan \frac{\pi}{4} = 1 \Rightarrow \tan^{-1} 1 = \frac{\pi}{4} \right] = \frac{\pi}{4}.
 \end{aligned}$$

6.  $\int_0^2 \frac{dx}{x+4-x^2}$

**Sol.**  $\int_0^2 \frac{dx}{4+x-x^2} = \int_0^2 \frac{dx}{-x^2+x+4} = \int_0^2 \frac{dx}{-(x^2-x-4)}$

(Making coeff. of  $x^2$  numerically unity)

Completing squares by adding and subtracting

$$\begin{aligned}
 \left( \frac{1}{2} \text{coeff. of } x \right)^2 &= \left( \frac{1}{2} \right)^2 = \frac{1}{4} \therefore \int_0^2 \frac{dx}{- \left[ x^2 - x + \frac{1}{4} - \frac{1}{4} - 4 \right]} \\
 &= \int_0^2 \frac{dx}{- \left[ \left( x - \frac{1}{2} \right)^2 - \frac{17}{4} \right]} = \int_0^2 \frac{dx}{\frac{17}{4} - \left( x - \frac{1}{2} \right)^2} = \int_0^2 \frac{dx}{\left( \frac{\sqrt{17}}{2} \right)^2 - \left( x - \frac{1}{2} \right)^2}
 \end{aligned}$$

$$= \frac{1}{2 \times \frac{\sqrt{17}}{2}} \left[ \log \left| \frac{\frac{\sqrt{17}}{2} + \left(x - \frac{1}{2}\right)}{\frac{\sqrt{17}}{2} - \left(x - \frac{1}{2}\right)} \right| \right]_0^2 \left( \because \int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| \right)$$

$$= \frac{1}{\sqrt{17}} \left[ \log \left| \frac{\sqrt{17} + 2x - 1}{\sqrt{17} - 2x + 1} \right| \right]_0^2$$

$$= \frac{1}{\sqrt{17}} \left[ \log \left| \frac{\sqrt{17} + 3}{\sqrt{17} - 3} \right| - \log \left| \frac{\sqrt{17} - 1}{\sqrt{17} + 1} \right| \right]$$

$$= \frac{1}{\sqrt{17}} \log \left( \frac{\sqrt{17} + 3}{\sqrt{17} - 3} \times \frac{\sqrt{17} + 1}{\sqrt{17} - 1} \right) = \frac{1}{\sqrt{17}} \log \frac{20 + 4\sqrt{17}}{20 - 4\sqrt{17}}$$

$$(\because (\sqrt{17} + 3)(\sqrt{17} + 1) = 17 + \sqrt{17} + 3\sqrt{17} + 3 = 20 + 4\sqrt{17}.)$$

$$\text{Similarly } (\sqrt{17} - 3)(\sqrt{17} - 1) = 20 - 4\sqrt{17})$$

$$= \frac{1}{\sqrt{17}} \log \frac{4(5 + \sqrt{17})}{4(5 - \sqrt{17})} = \frac{1}{\sqrt{17}} \log \frac{5 + \sqrt{17}}{5 - \sqrt{17}}$$

$$= \frac{1}{\sqrt{17}} \log \left( \frac{5 + \sqrt{17}}{5 - \sqrt{17}} \times \frac{5 + \sqrt{17}}{5 + \sqrt{17}} \right) = \frac{1}{\sqrt{17}} \log \frac{(5 + \sqrt{17})^2}{25 - 17}$$

$$= \frac{1}{\sqrt{17}} \log \frac{42 + 10\sqrt{17}}{8} = \frac{1}{\sqrt{17}} \log \frac{21 + 5\sqrt{17}}{4}.$$

7.  $\int_{-1}^1 \frac{dx}{x^2 + 2x + 5}$

**Sol.** Let  $I = \int_{-1}^1 \frac{dx}{x^2 + 2x + 5} = \int_{-1}^1 \frac{dx}{x^2 + 2x + 1 + 4}$  (To complete squares)  
 $= \int_{-1}^1 \frac{1}{(x+1)^2 + 2^2} dx \quad \dots(i)$

**Put  $x + 1 = t$ .**  $\therefore \frac{dx}{dt} = 1 \Rightarrow dx = dt$

**To change the limits of Integration**

When  $x = -1$ ,  $t = -1 + 1 = 0$

When  $x = 1$ ,  $t = 1 + 1 = 2$

$$\therefore \text{ From (i), } I = \int_0^2 \frac{1}{t^2 + 2^2} dt = \frac{1}{2} \left( \tan^{-1} \frac{t}{2} \right)_0^2$$

$$\left[ \because \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \right]$$

$$= \frac{1}{2} \left[ \tan^{-1} \frac{2}{2} - \tan^{-1} \frac{0}{2} \right] = \frac{1}{2} (\tan^{-1} 1 - \tan^{-1} 0)$$

$$= \frac{1}{2} \left( \frac{\pi}{4} - 0 \right) = \frac{\pi}{8}. \quad \left[ \because \tan \frac{\pi}{4} = 1 \text{ and } \tan 0 = 0 \right]$$

$$8. \int_1^2 \left( \frac{1}{x} - \frac{1}{2x^2} \right) e^{2x} dx$$

$$\text{Sol. Let } I = \int_1^2 \left( \frac{1}{x} - \frac{1}{2x^2} \right) e^{2x} dx \quad \dots(i)$$

[Type  $\int (f(x) + g(x)) e^{ax} dx$ . Put  $ax = t$  and it will become  $\int (f(t) + g(t)) e^t dt = e^t f(t)$ ]

$$\text{Put } 2x = t \quad \therefore \quad 2 = \frac{dt}{dx} \Rightarrow 2dx = dt \Rightarrow dx = \frac{dt}{2}$$

**To change the limits of Integration**

When  $x = 1$ ,  $t = 2x = 2$ , When  $x = 2$ ,  $t = 2x = 4$

$$\therefore \text{ From (i), } I = \int_2^4 \left( \frac{1}{t} - \frac{1}{2\left(\frac{t}{2}\right)^2} \right) e^t \frac{dt}{2} \quad \left[ \because 2x = t \Rightarrow x = \frac{t}{2} \right]$$

$$\begin{aligned} \therefore I &= \int_2^4 \left( \frac{2}{t} - \frac{2}{t^2} \right) e^t \frac{dt}{2} = \int_2^4 \frac{1}{2} \cdot 2 \left( \frac{1}{t} - \frac{1}{t^2} \right) e^t dt \\ &= \int_2^4 \left( \frac{1}{t} - \frac{1}{t^2} \right) e^t dt = \int_2^4 (f(t) + f'(t)) e^t dt \\ &\quad \left( \text{Here } f(t) = \frac{1}{t} = t^{-1} \text{ and therefore } f'(t) = (-1)t^{-2} = \frac{-1}{t^2} \right) \\ &= \left( e^t f(t) \right)_2^4 = \left( \frac{e^t}{t} \right)_2^4 = \frac{e^4}{4} - \frac{e^2}{2} = \frac{e^4 - 2e^2}{4} = \frac{e^2(e^2 - 2)}{4}. \end{aligned}$$

**Choose the correct answer in Exercises 9 and 10.**

9. The value of the integral  $\int_{\frac{1}{3}}^1 \frac{(x - x^3)^{1/3}}{x^4} dx$  is

(A) 6                      (B) 0                      (C) 3                      (D) 4

$$\text{Sol. Let } I = \int_{\frac{1}{3}}^1 \frac{(x - x^3)^{1/3}}{x^4} dx$$

$$= \int_{\frac{1}{3}}^1 \frac{\left[ x^3 \left( \frac{x}{x^3} - 1 \right) \right]^{1/3}}{x^4} dx = \int_{\frac{1}{3}}^1 \frac{(x^3)^{1/3} \left( \frac{1}{x^2} - 1 \right)^{1/3}}{x^4} dx$$

$$= \int_{\frac{1}{3}}^1 \frac{x(x^{-2} - 1)^{1/3}}{x^4} dx = \int_{\frac{1}{3}}^1 (x^{-2} - 1)^{1/3} x^{-3} dx$$

$$I = \frac{-1}{2} \int_{\frac{1}{3}}^1 (x^{-2} - 1)^{1/3} (-2x^{-3}) dx \quad \dots(i)$$

$$\text{Put } x^{-2} - 1 = t$$

$$\text{Therefore } -2x^{-3} = \frac{dt}{dx} \Rightarrow -2x^{-3} dx = dt$$

**To change the limits of Integration**

$$\begin{aligned} \text{When } x = \frac{1}{3}, t = x^{-2} - 1 &= \left(\frac{1}{3}\right)^{-2} - 1 \\ &= (3^{-1})^{-2} - 1 = 3^2 - 1 = 9 - 1 = 8 \\ \text{When } x = 1, t = 1^{-2} - 1 &= 1 - 1 = 0 \end{aligned}$$

$$\begin{aligned} \therefore \text{ From (i), } I &= \frac{-1}{2} \int_8^0 t^{1/3} dt = \frac{-1}{2} \left[ \frac{t^{4/3}}{\frac{4}{3}} \right]_8^0 \\ &= \frac{-1}{2} \cdot \frac{3}{4} [0 - 8^{4/3}] = \frac{-3}{8} [- (2^3)^{4/3}] = \frac{-3}{8} (-2^4) = \frac{3}{8} \times 16 = 6 \\ \therefore \text{ Option (A) is the correct answer.} \end{aligned}$$

**10. If  $f(x) = \int_0^x t \sin t \, dt$ , then  $f'(x)$  is**

- (A)  $\cos x + x \sin x$   
(C)  $x \cos x$

- (B)  $x \sin x$   
(D)  $\sin x + x \cos x$

**Sol.**  $f(x) = \int_0^x t \sin t \, dt$

I II

Applying Product Rule of Integration

$$\begin{aligned} \left[ \int_a^b I \cdot II \, dx &= \left( I \int II \, dx \right)_a^b - \int_a^b \frac{d}{dx} (I) \int II \, dx \, dx \right] \\ \Rightarrow f(x) &= \left( t (-\cos t) \right)_0^x - \int_0^x 1 (-\cos t) \, dt \\ &= -x \cos x - 0 + \int_0^x \cos t \, dt = -x \cos x + (\sin t)_0^x \\ &= -x \cos x + \sin x - \sin 0 = -x \cos x + \sin x \\ \therefore f'(x) &= - (x (-\sin x) + (\cos x)1) + \cos x \\ &= x \sin x - \cos x + \cos x = x \sin x \\ \therefore \text{ Option (B) is the correct answer.} \end{aligned}$$

**OR**

$$f(x) = \int_0^x \sin t \, dt \qquad \therefore f'(x) = (t \sin t)_0^x$$

[ $\therefore$  Derivative operator and integral operator cancel with each other]

$$= x \sin x - 0 = x \sin x.$$



### Exercise 7.11

By using the properties of definite integrals, evaluate the integrals in Exercises 1 to 6:

1.  $\int_0^{\frac{\pi}{2}} \cos^2 x \, dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \cos^2 x \, dx$  ... (i)

$$\therefore I = \int_0^{\frac{\pi}{2}} \cos^2 \left( \frac{\pi}{2} - x \right) dx \quad \left[ \because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

or  $I = \int_0^{\frac{\pi}{2}} \sin^2 x \, dx$  ... (ii)

Adding Eqns. (i) and (ii),

$$2I = \int_0^{\frac{\pi}{2}} (\cos^2 x + \sin^2 x) \, dx = \int_0^{\frac{\pi}{2}} 1 \, dx = (x)_0^{\frac{\pi}{2}}$$

$$\Rightarrow 2I = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}.$$

2.  $\int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} \, dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} \, dx$  ... (i)

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin \left( \frac{\pi}{2} - x \right)}}{\sqrt{\sin \left( \frac{\pi}{2} - x \right)} + \sqrt{\cos \left( \frac{\pi}{2} - x \right)}} \, dx$$

$$\left[ \because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

or  $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} \, dx$  ... (ii)

Adding Eqns. (i) and (ii), we have

$$2I = \int_0^{\frac{\pi}{2}} \left( \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} + \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} \right) dx$$

$$= \int_0^{\frac{\pi}{2}} \left( \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} \right) dx = \int_0^{\frac{\pi}{2}} 1 \, dx$$

$$\Rightarrow 2I = (x)_0^{\frac{\pi}{2}} = \frac{\pi}{2} - 0 = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}.$$

3.  $\int_0^{\frac{\pi}{2}} \frac{\sin^{3/2} x \, dx}{\sin^{3/2} x + \cos^{3/2} x}$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\sin^{3/2} x}{\sin^{3/2} x + \cos^{3/2} x} \, dx$  ... (i)

Changing  $x$  to  $\frac{\pi}{2} - x$   $\left[ \because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$

$$\begin{aligned}
 I &= \int_0^{\frac{\pi}{2}} \frac{\sin^{3/2}\left(\frac{\pi}{2} - x\right)}{\sin^{3/2}\left(\frac{\pi}{2} - x\right) + \cos^{3/2}\left(\frac{\pi}{2} - x\right)} dx \\
 &= \int_0^{\frac{\pi}{2}} \frac{\cos^{3/2} x}{\cos^{3/2} x + \sin^{3/2} x} dx \quad \dots(ii)
 \end{aligned}$$

Adding Eqns. (i) and (ii),

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sin^{3/2} x + \cos^{3/2} x}{\sin^{3/2} x + \cos^{3/2} x} dx = \int_0^{\frac{\pi}{2}} 1 dx = [x]_0^{\frac{\pi}{2}} = \frac{\pi}{2} \therefore I = \frac{\pi}{4}.$$

4.  $\int_0^{\frac{\pi}{2}} \frac{\cos^5 x dx}{\sin^5 x + \cos^5 x}$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\cos^5 x}{\sin^5 x + \cos^5 x} dx \quad \dots(i)$

$$\begin{aligned}
 \therefore I &= \int_0^{\frac{\pi}{2}} \frac{\cos^5\left(\frac{\pi}{2} - x\right)}{\sin^5\left(\frac{\pi}{2} - x\right) + \cos^5\left(\frac{\pi}{2} - x\right)} dx \\
 &\left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]
 \end{aligned}$$

or  $I = \int_0^{\frac{\pi}{2}} \frac{\sin^5 x}{\cos^5 x + \sin^5 x} dx \quad \dots(ii)$

Adding Eqns. (i) and (ii), we have

$$\begin{aligned}
 2I &= \int_0^{\frac{\pi}{2}} \left( \frac{\cos^5 x}{\sin^5 x + \cos^5 x} + \frac{\sin^5 x}{\cos^5 x + \sin^5 x} \right) dx \\
 \Rightarrow 2I &= \int_0^{\frac{\pi}{2}} \frac{\cos^5 x + \sin^5 x}{\sin^5 x + \cos^5 x} dx = \int_0^{\frac{\pi}{2}} 1 dx = (x)_0^{\frac{\pi}{2}}
 \end{aligned}$$

$$\Rightarrow 2I = \frac{\pi}{2} \quad \Rightarrow I = \frac{\pi}{4}.$$

5.  $\int_{-5}^5 |x+2| dx$

**Sol.** Let  $I = \int_{-5}^5 |x+2| dx \quad \dots(i)$

We can evaluate this integral only if we can get rid of the modulus.

Putting expression within modulus equal to 0, we have

$$x + 2 = 0, \text{ i.e., } x = -2 \in (-5, 5)$$

$\therefore$  From (i),  $I = \int_{-5}^5 |x+2| dx$

$$= \int_{-5}^{-2} |x+2| dx + \int_{-2}^5 |x+2| dx$$

$$\begin{aligned}
 & \left[ \because \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \text{ where } a < c < b \right] \\
 &= \int_{-5}^{-2} -(x+2) dx + \int_{-2}^5 (x+2) dx \\
 & \quad \left[ \because \text{On } (-5, -2), x < -2 \Rightarrow x+2 < 0 \right. \\
 & \Rightarrow |x+2| = -(x+2) \text{ and on } (-2, 5); x > -2 \\
 & \Rightarrow x+2 > 0 \Rightarrow |x+2| = x+2, \text{ by definition of modulus function} \left. \right] \\
 &= - \left( \frac{x^2}{2} + 2x \right)_{-5}^{-2} + \left( \frac{x^2}{2} + 2x \right)_{-2}^5 \\
 &= - \left[ \left( \frac{4}{2} - 4 \right) - \left( \frac{25}{2} - 10 \right) \right] + \left[ \left( \frac{25}{2} + 10 \right) - \left( \frac{4}{2} - 4 \right) \right] \\
 &= - \left[ -2 - \frac{5}{2} \right] + \left[ \frac{45}{2} + 2 \right] = 2 + \frac{5}{2} + \frac{45}{2} + 2 \\
 &= 4 + \frac{50}{2} = 4 + 25 = 29.
 \end{aligned}$$

6.  $\int_2^8 |x-5| dx$

**Sol.** We know by definition of modulus function, that

$$|x-5| = \begin{cases} x-5 & \text{if } x-5 \geq 0, \text{ i.e., } x \geq 5 & \dots(i) \\ -(x-5) = 5-x, & \text{if } x < 5 & \dots(ii) \end{cases}$$

$$\begin{aligned}
 \therefore \int_2^8 |x-5| dx &= \int_2^5 |x-5| dx + \int_5^8 |x-5| dx \\
 &= \int_2^5 (5-x) dx + \int_5^8 (x-5) dx = \left( 5x - \frac{x^2}{2} \right)_2^5 + \left( \frac{x^2}{2} - 5x \right)_5^8 \\
 & \quad \text{[By (ii)]} \quad \quad \quad \text{[By (i)]} \\
 &= \left( 25 - \frac{25}{2} \right) - (10 - 2) + (32 - 40) - \left( \frac{25}{2} - 25 \right) \\
 &= 25 - \frac{25}{2} - 8 - 8 - \frac{25}{2} + 25 = 34 - \frac{50}{2} = 34 - 25 = 9
 \end{aligned}$$

**By using the properties of definite integrals, evaluate the integrals in Exercises 7 to 11:**

7.  $\int_0^1 x(1-x)^n dx$

**Sol.** Let  $I = \int_0^1 x(1-x)^n dx$

$$\therefore I = \int_0^1 (1-x) (1-(1-x))^n dx \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\text{or } I = \int_0^1 (1-x) (1-1+x)^n dx$$

$$\begin{aligned}
 \text{or } I &= \int_0^1 (1-x) x^n dx = \int_0^1 (x^n - x^{n+1}) dx \\
 &= \left( \frac{x^{n+1}}{n+1} - \frac{x^{n+2}}{n+2} \right)_0^1 = \frac{1}{n+1} - \frac{1}{n+2} - (0-0) \\
 &= \frac{n+2-n-1}{(n+1)(n+2)} = \frac{1}{(n+1)(n+2)}.
 \end{aligned}$$

8.  $\int_0^{\frac{\pi}{4}} \log(1 + \tan x) dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{4}} \log(1 + \tan x) dx$  ... (i)

Changing  $x$  to  $\frac{\pi}{4} - x$   $\left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$

$$\begin{aligned}
 I &= \int_0^{\frac{\pi}{4}} \log \left[ 1 + \tan \left( \frac{\pi}{4} - x \right) \right] dx = \int_0^{\frac{\pi}{4}} \log \left[ 1 + \frac{1 - \tan x}{1 + \tan x} \right] dx \\
 &\quad \left[ \because \tan \left( \frac{\pi}{4} - x \right) = \frac{\tan \frac{\pi}{4} - \tan x}{1 + \tan \frac{\pi}{4} \tan x} = \frac{1 - \tan x}{1 + \tan x} \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\frac{\pi}{4}} \log \left( \frac{1 + \tan x + 1 - \tan x}{1 + \tan x} \right) dx \\
 &= \int_0^{\frac{\pi}{4}} \log \left( \frac{2}{1 + \tan x} \right) dx \quad \dots (ii)
 \end{aligned}$$

Adding Eqns. (i) and (ii), we have

$$\begin{aligned}
 2I &= \int_0^{\frac{\pi}{4}} \left[ \log(1 + \tan x) + \log \left( \frac{2}{1 + \tan x} \right) \right] dx \\
 &= \int_0^{\frac{\pi}{4}} \log \left[ (1 + \tan x) \frac{2}{(1 + \tan x)} \right] dx = \int_0^{\frac{\pi}{4}} \log 2 dx
 \end{aligned}$$

or  $2I = (\log 2) [x]_0^{\frac{\pi}{4}} = \frac{\pi}{4} \log 2$  Dividing by 2,  $I = \frac{\pi}{8} \log 2$ .

9.  $\int_0^2 x \sqrt{2-x} dx$

**Sol.** Let  $I = \int_0^2 x \sqrt{2-x} dx$

Changing  $x$  to  $2-x$   $\left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$

$$\begin{aligned}
 I &= \int_0^2 (2-x) \sqrt{2-(2-x)} dx \\
 &= \int_0^2 (2-x) \sqrt{x} dx = \int_0^2 (2x^{1/2} - x^{3/2}) dx
 \end{aligned}$$

$$= \left[ 2 \cdot \frac{x^{3/2}}{3/2} - \frac{x^{5/2}}{5/2} \right]_0^2 = \left( \frac{4}{3} \cdot 2^{3/2} - \frac{2}{5} \cdot 2^{5/2} \right) - (0 - 0)$$

$$= \frac{4}{3} \times 2\sqrt{2} - \frac{2}{5} \times 4\sqrt{2} = \left( \frac{8}{3} - \frac{8}{5} \right) \sqrt{2}$$

$$(\because 2^{3/2} = (2^{1/2})^3 = (\sqrt{2})^3 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 2\sqrt{2}$$

$$\text{and } 2^{5/2} = (2^{1/2})^5 = (\sqrt{2})^5 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 2.2.\sqrt{2}$$

$$= 4\sqrt{2}) = \frac{16\sqrt{2}}{15}.$$

10.  $\int_0^{\pi/2} (2 \log \sin x - \log \sin 2x) dx$

**Sol.** Let  $I = \int_0^{\pi/2} (2 \log \sin x - \log \sin 2x) dx$

$$= \int_0^{\pi/2} (\log \sin^2 x - \log \sin 2x) dx$$

$$= \int_0^{\pi/2} \log \left( \frac{\sin^2 x}{\sin 2x} \right) dx = \int_0^{\pi/2} \log \left( \frac{\sin^2 x}{2 \sin x \cos x} \right) dx$$

$$\text{or } I = \int_0^{\pi/2} \log \left( \frac{1}{2} \tan x \right) dx \quad \dots(i)$$

$$\therefore I = \int_0^{\pi/2} \log \left( \frac{1}{2} \tan \left( \frac{\pi}{2} - x \right) dx \right) \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\text{or } I = \int_0^{\pi/2} \log \left( \frac{1}{2} \cot x \right) dx \quad \dots(ii)$$

Adding Eqns. (i) and (ii),

$$2I = \int_0^{\pi/2} \left[ \log \left( \frac{1}{2} \tan x \right) + \log \left( \frac{1}{2} \cot x \right) \right] dx$$

$$\Rightarrow 2I = \int_0^{\pi/2} \log \left( \frac{1}{2} \tan x \cdot \frac{1}{2} \cot x \right) dx = \int_0^{\pi/2} \log \frac{1}{4} dx = \log \frac{1}{4} (x)_0^{\pi/2}$$

$$= (\log 1 - \log 4) \frac{\pi}{2} = -\frac{\pi}{2} \log 4 \quad (\because \log 1 = 0)$$

$$\therefore I = -\frac{\pi}{4} \log 4 = -\frac{\pi}{4} \log 2^2 = -\frac{2\pi}{4} \log 2 = -\frac{\pi}{2} \log 2.$$

11.  $\int_{-\pi/2}^{\pi/2} \sin^2 x dx$

**Sol.** Let  $I = \int_{-\pi/2}^{\pi/2} \sin^2 x dx$  or  $I = 2 \int_0^{\pi/2} \sin^2 x dx \quad \dots(i)$

$$[\because \text{For } f(x) = \sin^2 x, f(-x) = \sin^2(-x) = (-\sin x)^2 = \sin^2 x = f(x)]$$

$$\therefore f(x) \text{ is an even function of } x \text{ and hence}$$

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

$$\therefore I = 2 \int_0^{\frac{\pi}{2}} \sin^2 \left( \frac{\pi}{2} - x \right) dx \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\text{or } I = 2 \int_0^{\frac{\pi}{2}} \cos^2 x dx \quad \dots(ii)$$

Adding Eqns. (i) and (ii), we have

$$2I = 2 \int_0^{\frac{\pi}{2}} (\sin^2 x + \cos^2 x) dx$$

$$\text{or } 2I = 2 \int_0^{\frac{\pi}{2}} 1 dx = 2 \left( x \right)_0^{\frac{\pi}{2}} = 2 \cdot \frac{\pi}{2} = \pi \therefore I = \frac{\pi}{2}.$$

**Using properties of definite integrals, evaluate the following integrals in Exercises 12 to 18:**

12.  $\int_0^{\pi} \frac{x dx}{1 + \sin x}$

**Sol.** Let  $I = \int_0^{\pi} \frac{x}{1 + \sin x} dx \quad \dots(i)$

Changing  $x$  to  $\pi - x$ ,  $I = \int_0^{\pi} \frac{\pi - x}{1 + \sin(\pi - x)} dx$

or  $I = \int_0^{\pi} \frac{\pi - x}{1 + \sin x} dx \quad \dots(ii) \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$

Adding Eqns. (i) and (ii), we have

$$2I = \int_0^{\pi} \left( \frac{x}{1 + \sin x} + \frac{\pi - x}{1 + \sin x} \right) dx = \int_0^{\pi} \frac{x + \pi - x}{1 + \sin x} dx$$

$$= \int_0^{\pi} \frac{\pi}{1 + \sin x} dx = \pi \int_0^{\pi} \frac{1}{1 + \sin x} dx$$

or  $2I = 2\pi \int_0^{\pi/2} \frac{dx}{1 + \sin x}$

$$\left[ \because \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(2a-x) = f(x) \right]$$

$$= 2\pi \int_0^{\pi/2} \frac{dx}{1 + \sin \left( \frac{\pi}{2} - x \right)} \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= 2\pi \int_0^{\pi/2} \frac{dx}{1 + \cos x}$$

$$\Rightarrow I = \pi \int_0^{\pi/2} \frac{dx}{2 \cos^2 \frac{x}{2}} = \frac{\pi}{2} \int_0^{\pi/2} \sec^2 \frac{x}{2} dx = \frac{\pi}{2} \left[ \frac{\tan \frac{x}{2}}{\frac{1}{2}} \right]_0^{\pi/2}$$

$$= \pi \left( \tan \frac{\pi}{4} - \tan 0 \right) = \pi(1 - 0) = \pi.$$

13.  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^7 x \, dx$

**Sol.** Let  $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^7 x \, dx$

Here Integrand  $f(x) = \sin^7 x$

$$\therefore f(-x) = \sin^7(-x) = (-\sin x)^7 = -\sin^7 x = -f(x)$$

$\therefore f(x)$  is an odd function of  $x$ .

$$\therefore I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^7 x \, dx = 0.$$

$$\left[ \because \text{If } f(x) \text{ is an odd function of } x, \text{ then } \int_{-a}^a f(x) \, dx = 0 \right]$$

14.  $\int_0^{2\pi} \cos^5 x \, dx$

**Sol.**  $\int_0^{2\pi} \cos^5 x \, dx = 2 \int_0^{\pi} \cos^5 x \, dx$

$$\left[ \because \int_0^{2a} f(x) \, dx = \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x) \right]$$

Here  $f(x) = \cos^5 x \quad \therefore f(2\pi - x) = \cos^5(2\pi - x) = \cos^5 x$   
 $= f(x) = 2(0) = 0$

$$\left[ \because \int_0^{2a} f(x) \, dx = 0, \text{ if } f(2a-x) = -f(x). \text{ Here } f(x) = \cos^5 x \right]$$

$$\therefore f(\pi - x) = \cos^5(\pi - x) = (-\cos x)^5 = -\cos^5 x = -f(x)$$

**Alternatively.** To evaluate  $\int_0^{2\pi} \cos^5 x \, dx$ , put  $\sin x = t$ .

**Remark.** In fact  $\int_0^{2\pi} \cos^n x \, dx$  or  $\int_0^{\pi} \cos^n x \, dx$  for all positive odd integers  $n$  is equal to zero.

This is a very important result for I.I.T. Entrance Examination.

15.  $\int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 + \sin x \cos x} \, dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 + \sin x \cos x} \, dx \quad \dots(i)$

Changing  $x$  to  $\frac{\pi}{2} - x$  in integrand of (i),

$$\left[ \because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin\left(\frac{\pi}{2} - x\right) - \cos\left(\frac{\pi}{2} - x\right)}{1 + \sin\left(\frac{\pi}{2} - x\right) \cos\left(\frac{\pi}{2} - x\right)} \, dx = \int_0^{\frac{\pi}{2}} \frac{\cos x - \sin x}{1 + \cos x \sin x} \, dx$$

$$= - \int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 + \sin x \cos x} dx \quad \dots(ii)$$

Adding equations (i) and (ii), we have  $2I = 0$  or  $I = 0$ .

### 16. $\int_0^{\pi} \log(1 + \cos x) dx$

**Sol.** Let  $I = \int_0^{\pi} \log(1 + \cos x) dx \quad \dots(i)$

$$\therefore I = \int_0^{\pi} \log(1 + \cos(\pi - x)) dx \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a - x) dx \right]$$

or  $I = \int_0^{\pi} \log(1 - \cos x) dx \quad \dots(ii)$

Adding Eqns. (i) and (ii), we have

$$\begin{aligned} 2I &= \int_0^{\pi} [\log(1 + \cos x) + \log(1 - \cos x)] dx \\ &= \int_0^{\pi} \log((1 + \cos x)(1 - \cos x)) dx = \int_0^{\pi} \log(1 - \cos^2 x) dx \end{aligned}$$

$$\Rightarrow 2I = \int_0^{\pi} \log \sin^2 x dx = 2 \int_0^{\pi} \log \sin x dx \quad (\because \log m^n = n \log m)$$

Dividing by 2,  $I = \int_0^{\pi} \log \sin x dx = 2 \int_0^{\frac{\pi}{2}} \log \sin x dx \quad \dots(iii)$

$$\left[ \because \text{For } f(x) = \log \sin x, f(\pi - x) = \log \sin(\pi - x) = \log \sin x = f(x) \text{ and if } f(2a - x) = f(x); \text{ then } \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx \right]$$

$$\therefore I = 2 \int_0^{\frac{\pi}{2}} \log \sin\left(\frac{\pi}{2} - x\right) dx \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a - x) dx \right]$$

or  $I = 2 \int_0^{\frac{\pi}{2}} \log \cos x dx \quad \dots(iv)$

Adding Eqns. (iii) and (iv), we have

$$2I = 2 \int_0^{\frac{\pi}{2}} (\log \sin x + \log \cos x) dx$$

Dividing by 2,  $I = \int_0^{\frac{\pi}{2}} (\log \sin x \cos x) dx$

$$= \int_0^{\frac{\pi}{2}} \log\left(\frac{2 \sin x \cos x}{2}\right) dx = \int_0^{\frac{\pi}{2}} \log\left(\frac{\sin 2x}{2}\right) dx$$

or  $I = \int_0^{\frac{\pi}{2}} (\log \sin 2x - \log 2) dx$

or  $I = \int_0^{\frac{\pi}{2}} \log \sin 2x dx - \int_0^{\frac{\pi}{2}} \log 2 dx$

or  $I = \int_0^{\frac{\pi}{2}} \log \sin 2x dx - \log 2 \left(x\right)_0^{\frac{\pi}{2}}$

$$\text{or } I = \int_0^{\frac{\pi}{2}} \log \sin 2x \, dx - \frac{\pi}{2} \log 2$$

$$\text{or } I = I_1 - \frac{\pi}{2} \log 2 \quad \dots(v)$$

$$\text{where } I_1 = \int_0^{\frac{\pi}{2}} \log \sin 2x \, dx \quad \dots(vi)$$

**Put  $2x = t$**  to make  $I_1$  look as  $I$  given by (iii)

$$\therefore 2 = \frac{dt}{dx} \quad \text{or} \quad 2 \, dx = dt \quad \text{or} \quad dx = \frac{dt}{2}$$

To change the limits: When  $x = 0$ ,  $t = 2x = 0$

$$\text{When } x = \frac{\pi}{2}, \quad t = 2x = \pi$$

$$\therefore \text{ From (vi), } I_1 = \int_0^{\pi} \log \sin t \, \frac{dt}{2} = \frac{1}{2} \int_0^{\pi} \log \sin t \, dt$$

$$\text{or} \quad I_1 = \frac{1}{2} \times 2 \int_0^{\frac{\pi}{2}} \log \sin t \, dt$$

(For reason see Explanation within brackets below Eqn. (iii))

$$\text{or } I_1 = \int_0^{\frac{\pi}{2}} \log \sin t \, dt = \int_0^{\frac{\pi}{2}} \log \sin x \, dx \left[ \because \int_a^b f(t) \, dt = \int_a^b f(x) \, dx \right]$$

$$\text{or } I_1 = \frac{I}{2} \quad [\text{By Eqn. (iii)}]$$

$$\text{Putting this value of } I_1 \text{ in Eqn. (v), } I = \frac{I}{2} - \frac{\pi}{2} \log 2$$

Multiplying by L.C.M. = 2,  $2I = I - \pi \log 2$

$$\text{or } 2I - I = -\pi \log 2 \quad \text{or } I = -\pi \log 2.$$

$$17. \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} \, dx$$

$$\text{Sol. Let } I = \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} \, dx \quad \dots(i)$$

$$\therefore I = \int_0^a \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{a-(a-x)}} \, dx = \int_0^a \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{x}} \, dx \quad \dots(ii)$$

Adding Eqns. (i) and (ii), we have

$$2I = \int_0^a \left( \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} + \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{x}} \right) dx = \int_0^a \left( \frac{\sqrt{x} + \sqrt{a-x}}{\sqrt{x} + \sqrt{a-x}} \right) dx$$

$$\text{or } 2I = \int_0^a 1 \, dx = (x)_0^a = a \therefore I = \frac{a}{2}.$$

18.  $\int_0^4 |x-1| dx$



**Sol.** Let  $I = \int_0^4 |x-1| dx$  ... (i)

Putting the expression  $(x-1)$  within modulus equal to zero, we have  
 $x = 1 \in (0, 4)$

$$\begin{aligned} \therefore \text{From (i), } I &= \int_0^4 |x-1| dx = \int_0^1 |x-1| dx + \int_1^4 |x-1| dx \\ &= - \int_0^1 (x-1) dx + \int_1^4 (x-1) dx \end{aligned}$$

[ $\because$  On  $(0, 1)$ ;  $x < 1 \Rightarrow x-1 < 0$  and hence  $|x-1| = -(x-1)$  and on  $(1, 4)$ ,  $x > 1 \Rightarrow x-1 > 0$  and hence  
 $|x-1| = (x-1)$  by definition of modulus function]

$$\begin{aligned} &= - \left( \frac{x^2}{2} - x \right)_0^1 + \left( \frac{x^2}{2} - x \right)_1^4 = - \left( \left( \frac{1}{2} - 1 \right) - 0 \right) + \left( \frac{16}{2} - 4 - \left( \frac{1}{2} - 1 \right) \right) \\ &= - \frac{1}{2} + 1 + 8 - 4 - \frac{1}{2} + 1 = 6 - \frac{2}{2} = 6 - 1 = 5. \end{aligned}$$

19. Show that  $\int_0^a f(x) g(x) dx = 2 \int_0^a f(x) dx$ , if  $f$  and  $g$  are defined as  $f(x) = f(a-x)$  and  $g(x) + g(a-x) = 4$ .

**Sol. Given:**  $f(x) = f(a-x)$  ... (i)

and  $g(x) + g(a-x) = 4$  ... (ii)

Let  $I = \int_0^a f(x) g(x) dx$  ... (iii)

$$\therefore I = \int_0^a f(a-x) g(a-x) dx \quad \left[ \because \int_0^a F(x) dx = \int_0^a F(a-x) dx \right]$$

Putting  $f(a-x) = f(x)$  from (i),

$$I = \int_0^a f(x) g(a-x) dx \quad \dots (iv)$$

Adding Eqns. (iii) and (iv), we have

$$2I = \int_0^a (f(x) g(x) + f(x) g(a-x)) dx = \int_0^a f(x) (g(x) + g(a-x)) dx$$

$$\text{or } 2I = \int_0^a f(x) (4) dx \quad [\text{By (ii)}] = 4 \int_0^a f(x) dx$$

$$\text{Dividing by 2, } I = 2 \int_0^a f(x) dx = \text{R.H.S.}$$

**Choose the correct answer in Exercises 20 and 21:**

20. The value of  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (x^3 + x \cos x + \tan^5 x + 1) dx$  is

(A) 0

(B) 2

(C)  $\pi$

(D) 1

**Sol.** Let  $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (x^3 + x \cos x + \tan^5 x + 1) dx$

$$= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x^3 dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x \cos x dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \tan^5 x dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 dx$$

$$= 0 + 0 + 0 + \left( x \right)_{-\frac{\pi}{2}}^{\frac{\pi}{2}} = \frac{\pi}{2} - \left( -\frac{\pi}{2} \right) = \frac{\pi}{2} + \frac{\pi}{2} = \pi$$

$\left[ \because \text{Each of the three functions } x^3, x \cos x \text{ and } \tan^5 x \text{ is an odd function of } x \text{ as } f(-x) = -f(x) \text{ for each of them and } \int_{-a}^a f(x) dx = 0 \text{ for each odd function } f(x) \right]$

$\therefore$  Option (C) is the correct option.

**21. The value of  $\int_0^{\frac{\pi}{2}} \log \left( \frac{4+3 \sin x}{4+3 \cos x} \right) dx$  is**

(A) 2

(B)  $\frac{3}{4}$

(C) 0

(D) -2

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \log \left( \frac{4+3 \sin x}{4+3 \cos x} \right) dx$  ... (i)

$$\therefore I = \int_0^{\frac{\pi}{2}} \log \left( \frac{4+3 \sin \left( \frac{\pi}{2} - x \right)}{4+3 \cos \left( \frac{\pi}{2} - x \right)} \right) dx$$

or  $I = \int_0^{\frac{\pi}{2}} \log \left( \frac{4+3 \cos x}{4+3 \sin x} \right) dx$  ... (ii)

Adding Eqns. (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^{\frac{\pi}{2}} \left[ \log \left( \frac{4+3 \sin x}{4+3 \cos x} \right) + \log \left( \frac{4+3 \cos x}{4+3 \sin x} \right) \right] dx \\
 &= \int_0^{\frac{\pi}{2}} \log \left[ \frac{4+3 \sin x}{4+3 \cos x} \cdot \frac{4+3 \cos x}{4+3 \sin x} \right] dx = \int_0^{\frac{\pi}{2}} \log 1 dx
 \end{aligned}$$

$$\Rightarrow 2I = \int_0^{\frac{\pi}{2}} 0 dx = 0 \quad \Rightarrow \quad I = \frac{0}{2} = 0.$$



## MISCELLANEOUS EXERCISE

Integrate the functions in Exercises 1 to 11:

1.  $\frac{1}{x - x^3}$

**Sol.** The integrand  $\frac{1}{x - x^3}$  is a rational function of  $x$  and the

denominator  $x - x^3 = x(1 - x^2) = x(1 - x)(1 + x)$  is the product of more than one factor. So, will form partial fractions.

$$\begin{aligned}\frac{1}{x - x^3} &= \frac{1}{x(1 - x^2)} = \frac{1}{x(1 - x)(1 + x)} \\ &= \frac{A}{x} + \frac{B}{1 - x} + \frac{C}{1 + x} \quad \dots(i)\end{aligned}$$

Multiplying every term of Eqn. (i) by L.C.M.  $= x(1 - x)(1 + x)$ ,

$$1 = A(1 - x)(1 + x) + Bx(1 + x) + Cx(1 - x)$$

$$\text{or } 1 = A(1 - x^2) + B(x + x^2) + C(x - x^2)$$

$$\Rightarrow 1 = A - Ax^2 + Bx + Bx^2 + Cx - Cx^2$$

Comparing coefficients of like powers on both sides,

$$\mathbf{x^2:} -A + B - C = 0 \quad \dots(ii)$$

$$\mathbf{x:} \quad B + C = 0 \quad \dots(iii)$$

$$\mathbf{Constants:} \quad A = 1$$

$$\text{Putting } A = 1 \text{ in (ii), } -1 + B - C = 0 \quad \text{or} \quad B - C = 1 \quad \dots(iv)$$

$$\text{Adding Eqns. (iii) and (iv), } 2B = 1 \quad \Rightarrow \quad B = \frac{1}{2}$$

$$\text{From (iii), } C = -B = \frac{-1}{2}$$

Putting these values of A, B, C in (i),

$$\begin{aligned}\frac{1}{x - x^3} &= \frac{1}{x} + \frac{\frac{1}{2}}{1 - x} - \frac{\frac{1}{2}}{1 + x} \\ \therefore \int \frac{1}{x - x^3} dx &= \int \frac{1}{x} dx + \frac{1}{2} \int \frac{1}{1 - x} dx - \frac{1}{2} \int \frac{1}{1 + x} dx \\ &= \log |x| + \frac{1}{2} \frac{\log |1 - x|}{-1} - \frac{1}{2} \log |1 + x| \\ &= \frac{1}{2} [2 \log |x| - \log |1 - x| - \log |1 + x|] + C \\ &= \frac{1}{2} [\log |x|^2 - (\log |1 - x| + \log |1 + x|)] + C \\ &= \frac{1}{2} [\log |x|^2 - \log |1 - x| |1 + x|] + C \\ &= \frac{1}{2} [\log |x|^2 - \log |1 - x^2|] + C = \frac{1}{2} \log \left| \frac{x^2}{1 - x^2} \right| + C.\end{aligned}$$

$$2. \frac{1}{\sqrt{x+a} + \sqrt{x+b}}$$

$$\text{Sol. } \int \frac{1}{\sqrt{x+a} + \sqrt{x+b}} dx$$

$$\begin{aligned}
 \text{Rationalising, } &= \int \frac{\sqrt{x+a} - \sqrt{x+b}}{(\sqrt{x+a} + \sqrt{x+b})(\sqrt{x+a} - \sqrt{x+b})} dx \\
 &= \int \frac{(\sqrt{x+a} - \sqrt{x+b})}{x+a - (x+b)} dx = \int \frac{\sqrt{x+a} - \sqrt{x+b}}{a-b} dx \\
 &\quad [\because x+a - (x+b) = x+a - x - b = a-b] \\
 &= \frac{1}{a-b} \int (\sqrt{x+a} - \sqrt{x+b}) dx \\
 &= \frac{1}{a-b} \left[ \int (x+a)^{1/2} dx - \int (x+b)^{1/2} dx \right] \\
 &= \frac{1}{a-b} \left[ \frac{(x+a)^{3/2}}{\frac{3}{2}(1)} - \frac{(x+b)^{3/2}}{\frac{3}{2}(1)} \right] + C \\
 &= \frac{1}{a-b} \left[ \frac{2}{3} (x+a)^{3/2} - \frac{2}{3} (x+b)^{3/2} \right] + C \\
 &= \frac{2}{3(a-b)} [(x+a)^{3/2} - (x+b)^{3/2}] + C.
 \end{aligned}$$

3.  $\frac{1}{x\sqrt{ax-x^2}}$

Sol.  $I = \int \frac{dx}{x\sqrt{ax-x^2}} \quad \left[ \begin{array}{l} \text{Form } \int \frac{dx}{\text{Linear} \sqrt{\text{Quadratic}}} \end{array} \right]$

**Put Linear**  $= \frac{1}{t}$ , i.e.,  $x = \frac{1}{t} = t^{-1}$ .

Differentiating both sides  $dx = -\frac{1}{t^2} dt$

$$\begin{aligned}
 \therefore I &= \int \frac{-\frac{1}{t^2} dt}{\frac{1}{t} \sqrt{\frac{a}{t} - \frac{1}{t^2}}} = - \int \frac{dt}{\sqrt{at-1}} \\
 &= - \int (at-1)^{-1/2} dt = - \frac{(at-1)^{1/2}}{\frac{1}{2} \times a} + c \\
 &= - \frac{2}{a} \sqrt{\frac{a}{t}-1} + c = - \frac{2}{a} \sqrt{\frac{a-x}{x}} + c.
 \end{aligned}$$

4.  $\frac{1}{x^2(x^4+1)^{3/4}}$

Sol.  $I = \int \frac{dx}{x^2(x^4+1)^{3/4}} = \int \frac{dx}{x^2 \left[ x^4 \left( 1 + \frac{1}{x^4} \right) \right]^{3/4}} = \int \frac{dx}{x^2 \cdot x^3 \left( 1 + \frac{1}{x^4} \right)^{3/4}} \quad \left[ \because (x^4)^{\frac{3}{4}} = x^3 \right]$

$$= \int \frac{1}{x^5} \left(1 + \frac{1}{x^4}\right)^{-3/4} dx$$

Put  $1 + \frac{1}{x^4} = t$  or  $1 + x^{-4} = t$ .

Differentiating both sides,  $-4x^{-5} dx = dt$

or  $-\frac{4}{x^5} dx = dt$  or  $\frac{1}{x^5} dx = -\frac{1}{4} dt$

$$\therefore I = -\frac{1}{4} \int t^{-3/4} dt = -\frac{1}{4} \cdot \frac{t^{1/4}}{1/4} + c = -\left(1 + \frac{1}{x^4}\right)^{1/4} + c.$$

5.  $\frac{1}{x^{1/2} + x^{1/3}}$

**Sol.** Here the denominators of fractional powers  $\frac{1}{2}$  and  $\frac{1}{3}$  of  $x$  are 2 and 3. L.C.M. of 2 and 3 is 6.

**Put  $x = t^6$ .** Differentiating both sides,  $dx = 6t^5 dt$

$$\begin{aligned} \therefore I &= \int \frac{dx}{x^{1/2} + x^{1/3}} = \int \frac{6t^5}{t^3 + t^2} dt = 6 \int \frac{t^5}{t^2(t+1)} dt \\ &= 6 \int \frac{t^3}{t+1} dt = 6 \int \frac{(t^3+1)-1}{t+1} dt = 6 \int \left[ \frac{t^3+1}{t+1} - \frac{1}{t+1} \right] dt \\ &= 6 \int \left[ \frac{(t+1)(t^2-t+1)}{t+1} - \frac{1}{t+1} \right] dt = 6 \int \left( t^2 - t + 1 - \frac{1}{t+1} \right) dt \\ &\quad [\because a^3 + b^3 = (a+b)(a^2 - ab + b^2)] \\ &= 6 \left[ \frac{t^3}{3} - \frac{t^2}{2} + t - \log |t+1| \right] + c \\ &= 2t^3 - 3t^2 + 6t - 6 \log |t+1| + c \end{aligned}$$

Putting  $t = x^{1/6}$  ( $\because x = t^6 \Rightarrow t = x^{1/6}$ )

$$= 2\sqrt{x} - 3x^{1/3} + 6x^{1/6} - 6 \log |x^{1/6} + 1| + c.$$

6.  $\frac{5x}{(x+1)(x^2+9)}$

**Sol.** Let  $I = \int \frac{5x}{(x+1)(x^2+9)} dx$  ... (i)

Let  $\frac{5x}{(x+1)(x^2+9)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+9}$  ... (ii)

L.C.M. =  $(x+1)(x^2+9)$

Multiplying every term of (ii) by L.C.M.,

$$5x = A(x^2+9) + (Bx+C)(x+1)$$

or  $5x = Ax^2 + 9A + Bx^2 + Bx + Cx + C$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides,

**$x^2$ :**  $A + B = 0$  ... (iii)

**$x$ :**  $B + C = 5$  ... (iv)

$$\text{Constant terms : } 9A + C = 0 \quad \dots(v)$$

Let us solve Eqns. (iii), (iv) and (v) for A, B, C.

$$(iii) - (iv) \text{ gives, (to eliminate B), } A - C = -5 \quad \dots(vi)$$

$$\text{Adding (v) and (vi),} \quad 10A = -5$$

$$\therefore A = \frac{-5}{10} = \frac{-1}{2}$$

$$\text{Putting } A = \frac{-1}{2} \text{ in (iii), } \frac{-1}{2} + B = 0 \Rightarrow B = \frac{1}{2}$$

$$\text{Putting } B = \frac{1}{2} \text{ in (iv), } \frac{1}{2} + C = 5 \Rightarrow C = 5 - \frac{1}{2} = \frac{9}{2}$$

Putting these values of A, B, C in (ii),

$$\frac{5x}{(x+1)(x^2+9)} = \frac{-1}{x+1} + \frac{\frac{1}{2}x + \frac{9}{2}}{x^2+9}$$

$$\therefore \int \frac{5x}{(x+1)(x^2+9)} dx$$

$$= \frac{-1}{2} \int \frac{1}{x+1} dx + \frac{1}{2} \int \frac{x}{x^2+9} dx + \frac{9}{2} \int \frac{1}{x^2+3^2} dx$$

$$= \frac{-1}{2} \log |x+1| + \frac{1}{4} \int \frac{2x}{x^2+9} dx + \frac{9}{2} \cdot \frac{1}{3} \tan^{-1} \frac{x}{3} + c$$

$$= \frac{-1}{2} \log |x+1| + \frac{1}{4} \log |x^2+9| + \frac{3}{2} \tan^{-1} \frac{x}{3} + c$$

$$\left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| \right]$$

$$= \frac{-1}{2} \log |x+1| + \frac{1}{4} \log (x^2+9) + \frac{3}{2} \tan^{-1} \frac{x}{3} + c.$$

$$(\because x^2+9 \geq 9 > 0 \text{ and hence } |x^2+9| = x^2+9)$$

## 7. $\frac{\sin x}{\sin(x-a)}$

$$\text{Sol. } \int \frac{\sin x}{\sin(x-a)} dx = \int \frac{\sin(x-a+a)}{\sin(x-a)} dx$$

$$= \int \frac{\sin(x-a) \cos a + \cos(x-a) \sin a}{\sin(x-a)} dx$$

$$[\because \sin(A+B) = \sin A \cos B + \cos A \sin B]$$

$$= \int \left[ \frac{\sin(x-a) \cos a}{\sin(x-a)} + \frac{\cos(x-a) \sin a}{\sin(x-a)} \right] dx \quad \left[ \because \frac{a+b}{c} = \frac{a}{c} + \frac{b}{c} \right]$$

$$= \int [\cos a + \sin a \cot(x-a)] dx = \int \cos a dx + \int \sin a \cot(x-a) dx$$

$$= \cos a \int 1 dx + \sin a \int \cot(x-a) dx$$

$$= (\cos a)x + \sin a \frac{\log |\sin(x-a)|}{1} + c [\because \int \cot x dx = \log |\sin x|]$$

$$= x \cos a + \sin a \log |\sin(x-a)| + c.$$

$$8. \frac{e^{5 \log x} - e^{4 \log x}}{e^{3 \log x} - e^{2 \log x}}$$

$$\begin{aligned} \text{Sol. } \int \frac{e^{5 \log x} - e^{4 \log x}}{e^{3 \log x} - e^{2 \log x}} dx &= \int \frac{e^{\log x^5} - e^{\log x^4}}{e^{\log x^3} - e^{\log x^2}} dx \quad [\because n \log m = \log m^n] \\ &= \int \frac{x^5 - x^4}{x^3 - x^2} dx \quad [\because e^{\log f(x)} = f(x)] \\ &= \int \frac{x^4(x-1)}{x^2(x-1)} dx = \int x^2 dx = \frac{x^3}{3} + c. \end{aligned}$$

$$9. \frac{\cos x}{\sqrt{4 - \sin^2 x}}$$

$$\text{Sol. Let } I = \int \frac{\cos x}{\sqrt{4 - \sin^2 x}} dx \quad \dots(i)$$

$$\text{Put } \sin x = t. \text{ Therefore } \cos x = \frac{dt}{dx} \Rightarrow \cos x dx = dt$$

$$\begin{aligned} \therefore \text{ From (i), } I &= \int \frac{dt}{\sqrt{4 - t^2}} = \int \frac{dt}{\sqrt{2^2 - t^2}} \\ &= \sin^{-1} \left( \frac{t}{2} \right) + c \quad \left[ \because \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a} \right] \\ &= \sin^{-1} \left[ \frac{1}{2} \sin x \right] + c. \end{aligned}$$

$$10. \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x}$$

$$\text{Sol. Let } I = \int \frac{\sin^8 x - \cos^8 x}{1 - 2 \sin^2 x \cos^2 x} dx \quad \dots(i)$$

$$\begin{aligned} \text{Now numerator of integrand} &= \sin^8 x - \cos^8 x \\ &= (\sin^4 x)^2 - (\cos^4 x)^2 \\ &= (\sin^4 x - \cos^4 x)(\sin^4 x + \cos^4 x) \quad [\because a^2 - b^2 = (a - b)(a + b)] \\ &= [(\sin^2 x)^2 - (\cos^2 x)^2] [(\sin^2 x)^2 + (\cos^2 x)^2] \\ &= (\sin^2 x + \cos^2 x)(\sin^2 x - \cos^2 x) \\ &\quad [(\sin^2 x + \cos^2 x)^2 - 2 \sin^2 x \cos^2 x] \\ &\quad [\because a^2 + b^2 = a^2 + b^2 + 2ab - 2ab = (a + b)^2 - 2ab] \\ &= 1 [-(\cos^2 x - \sin^2 x)] (1 - 2 \sin^2 x \cos^2 x) \\ &\Rightarrow \sin^8 x - \cos^8 x = -\cos 2x (1 - 2 \sin^2 x \cos^2 x) \end{aligned}$$

Putting this value of  $\sin^8 x - \cos^8 x$  in numerator of (i),

$$I = \int \frac{-\cos 2x (1 - 2 \sin^2 x \cos^2 x)}{1 - 2 \sin^2 x \cos^2 x} dx = \int -\cos 2x dx = -\frac{\sin 2x}{2} + c.$$

11.  $\frac{1}{\cos(x+a)\cos(x+b)}$

**Sol.** Let  $I = \int \frac{1}{\cos(x+a)\cos(x+b)} dx$  ... (i)

We know that  $(x+a) - (x+b) = x+a-x-b = a-b$  ... (ii)

Dividing and multiplying by  $\sin(a-b)$  in (i),

$$I = \frac{1}{\sin(a-b)} \int \frac{\sin(a-b)}{\cos(x+a)\cos(x+b)} dx$$

Replacing  $(a-b)$  by  $(x+a) - (x+b)$  in  $\sin(a-b)$

[Using (ii)],

$$= \frac{1}{\sin(a-b)} \int \frac{\sin[(x+a) - (x+b)]}{\cos(x+a)\cos(x+b)} dx$$

$$= \frac{1}{\sin(a-b)} \int \frac{\sin(x+a)\cos(x+b) - \cos(x+a)\sin(x+b)}{\cos(x+a)\cos(x+b)} dx$$

[ $\because \sin(A-B) = \sin A \cos B - \cos A \sin B$ ]

$$= \frac{1}{\sin(a-b)} \int \left[ \frac{\sin(x+a)\cos(x+b)}{\cos(x+a)\cos(x+b)} - \frac{\cos(x+a)\sin(x+b)}{\cos(x+a)\cos(x+b)} \right] dx$$

$$\left( \because \frac{a-b}{c} = \frac{a}{c} - \frac{b}{c} \right)$$

$$= \frac{1}{\sin(a-b)} \int [\tan(x+a) - \tan(x+b)] dx$$

$$= \frac{1}{\sin(a-b)} [-\log|\cos(x+a)| + \log|\cos(x+b)|] + c$$

$$\left[ \because \int \tan x dx = -\log|\cos x| \right]$$

$$= \frac{1}{\sin(a-b)} \log \left| \frac{\cos(x+b)}{\cos(x+a)} \right| + c. \quad \left[ \because \log m - \log n = \log \frac{m}{n} \right]$$

**Integrate the functions in Exercises 12 to 22:**

12.  $\frac{x^3}{\sqrt{1-x^8}}$

**Sol.** Let  $I = \int \frac{x^3}{\sqrt{1-x^8}} dx = \frac{1}{4} \int \frac{4x^3}{\sqrt{1-(x^4)^2}} dx$  ... (i)

**Put  $x^4 = t$ .** Therefore  $4x^3 = \frac{dt}{dx} \Rightarrow 4x^3 dx = dt$

$\therefore$  From (i),  $I = \frac{1}{4} \int \frac{dt}{\sqrt{1-t^2}} = \frac{1}{4} \sin^{-1} t + c$

or  $I = \frac{1}{4} \sin^{-1}(x^4) + c.$

$$13. \frac{e^x}{(1+e^x)(2+e^x)}$$

**Sol.** Let  $I = \int \frac{e^x}{(1+e^x)(2+e^x)} dx$  ... (i)

[Rule to evaluate  $\int f(e^x) dx$ , put  $e^x = t$ ]

Put  $e^x = t$ . Therefore  $e^x = \frac{dt}{dx} \Rightarrow e^x dx = dt$

$\therefore$  From (i),  $I = \int \frac{dt}{(1+t)(2+t)} = \int \frac{1}{(t+1)(t+2)} dt$  ... (ii)

Now  $t + 2 - (t + 1) = t + 2 - t - 1 = 1$

Replacing 1 in the numerator of integrand in (ii) by (this)

$$(t + 2) - (t + 1),$$

$$I = \int \frac{(t+2) - (t+1)}{(t+1)(t+2)} dt = \int \left( \frac{t+2}{(t+1)(t+2)} - \frac{t+1}{(t+1)(t+2)} \right) dt$$

$$= \int \left( \frac{1}{t+1} - \frac{1}{t+2} \right) dt$$

$$= \log |t+1| - \log |t+2| + c = \log \left| \frac{t+1}{t+2} \right| + c$$

Putting  $t = e^x$ ,  $= \log \left| \frac{e^x+1}{e^x+2} \right| + c = \log \left( \frac{e^x+1}{e^x+2} \right) + c$ .

[ $\because e^x + 1 > 0$  and  $e^x + 2 > 0$  and  $|t| = t$  if  $t \geq 0$ ]

$$14. \frac{1}{(x^2+1)(x^2+4)}$$

**Sol.** Let  $I = \int \frac{1}{(x^2+1)(x^2+4)} dx$  ... (i)

Put  $x^2 = y$  only in the integrand.

Now the integrand is  $\frac{1}{(y+1)(y+4)}$

Let  $\frac{1}{(y+1)(y+4)} = \frac{A}{y+1} + \frac{B}{y+4}$  ... (ii)

Multiplying by L.C.M. =  $(y+1)(y+4)$ ,

$$1 = A(y+4) + B(y+1)$$

$$\text{or } 1 = Ay + 4A + By + B$$

comparing coefficient of  $y$ ,  $A + B = 0$  ... (iii)

comparing constants,  $4A + B = 1$  ... (iv)

Let us solve (iii) and (iv) for  $A$  and  $B$ .

(iv) - (iii) gives  $3A = 1 \therefore A = \frac{1}{3}$

From (iii)  $B = -A = -\frac{1}{3}$

Putting values of A, B and y in (ii),

$$\frac{1}{(x^2+1)(x^2+4)} = \frac{\frac{1}{3}}{x^2+1} + \frac{-\frac{1}{3}}{x^2+4} = \frac{1}{3} \left( \frac{1}{x^2+1} - \frac{1}{x^2+4} \right)$$

Putting this value in (i),

$$\begin{aligned} I &= \frac{1}{3} \int \left( \frac{1}{x^2+1} - \frac{1}{x^2+2^2} \right) dx = \frac{1}{3} \left[ \int \frac{1}{x^2+1} dx - \int \frac{1}{x^2+2^2} dx \right] \\ &= \frac{1}{3} \left[ \tan^{-1} x - \frac{1}{2} \tan^{-1} \frac{x}{2} \right] + c. \end{aligned}$$

**15.  $\cos^3 x e^{\log \sin x}$**

**Sol.** Let  $I = \int \cos^3 x e^{\log \sin x} dx = \int \cos^3 x \sin x dx$   
 $= - \int \cos^3 x (-\sin x) dx \quad \dots(i)$

**Put  $\cos x = t$ .**  $\therefore -\sin x = \frac{dt}{dx} \Rightarrow -\sin x dx = dt$

$\therefore$  From (i),  $I = - \int t^3 dt = \frac{-t^4}{4} + c = \frac{-1}{4} \cos^4 x + c.$

**16.  $e^{3 \log x} (x^4 + 1)^{-1}$**

**Sol.** Let  $I = \int e^{3 \log x} (x^4 + 1)^{-1} dx = \int \frac{e^{\log x^3}}{x^4 + 1} dx = \int \frac{x^3}{x^4 + 1} dx$   
 $[\because e^{\log f(x)} = f(x)]$

$\Rightarrow I = \frac{1}{4} \int \frac{4x^3}{x^4 + 1} dx \quad \dots(i)$

**Put  $x^4 + 1 = t$ .** Therefore  $4x^3 = \frac{dt}{dx} \Rightarrow 4x^3 dx = dt$

$\therefore$  From (i),  $I = \frac{1}{4} \int \frac{dt}{t} = \frac{1}{4} \log |t| + c$

Putting  $t = x^4 + 1$ ,  $= \frac{1}{4} \log |x^4 + 1| + c = \frac{1}{4} \log (x^4 + 1) + c.$   
 $[\because x^4 + 1 > 0 \Rightarrow |x^4 + 1| = x^4 + 1]$

**17.  $\int f'(ax+b)(f(ax+b))^n dx$**

**Sol.** Let  $I = \int f'(ax+b)(f(ax+b))^n dx$   
 $= \frac{1}{a} \int (f(ax+b))^n af'(ax+b) dx \quad \dots(i)$

Put  $f(ax+b) = t$ . Therefore  $f'(ax+b) \frac{d}{dx} (ax+b) = \frac{dt}{dx}$   
 $\Rightarrow af'(ax+b) dx = dt$

$\therefore$  From (i),  $I = \frac{1}{a} \int t^n dt = \frac{1}{a} \frac{t^{n+1}}{n+1} + c$  if  $n \neq -1$

$$\begin{aligned}\text{and if } n = -1, \text{ then } I &= \frac{1}{a} \int t^{-1} dt = \frac{1}{a} \int \frac{1}{t} dt \\ &= \frac{1}{a} \log |t| + c.\end{aligned}$$

$$\text{Putting } t = f(ax + b), I = \frac{(f(ax + b))^{n+1}}{a(n+1)} + c \text{ if } n \neq -1$$

$$\text{and } I = \frac{1}{a} |\log f(ax + b)| + c \text{ if } n = -1.$$

18.  $\frac{1}{\sqrt{\sin^3 x \sin(x + \alpha)}}$

$$\begin{aligned}\text{Sol. } I &= \int \frac{dx}{\sqrt{\sin^3 x \sin(x + \alpha)}} = \int \frac{dx}{\sqrt{\sin^3 x (\sin x \cos \alpha + \cos x \sin \alpha)}} \\ &= \int \frac{dx}{\sqrt{\sin^3 x \cdot \sin x (\cos \alpha + \cot x \sin \alpha)}} \\ &= \int \frac{dx}{\sin^2 x \sqrt{\cos \alpha + \cot x \sin \alpha}} = \int \frac{\operatorname{cosec}^2 x dx}{\sqrt{\cos \alpha + \cot x \sin \alpha}}\end{aligned}$$

Put  $\cos \alpha + \cot x \sin \alpha = t$ . Differentiating both sides  
 $-\operatorname{cosec}^2 x \sin \alpha dx = dt$

$$\text{or } \operatorname{cosec}^2 x dx = -\frac{dt}{\sin \alpha}$$

$$\begin{aligned}\therefore I &= \int -\frac{dt}{\sin \alpha \sqrt{t}} = -\frac{1}{\sin \alpha} \int t^{-1/2} dt \\ &= -\frac{1}{\sin \alpha} \cdot \frac{t^{1/2}}{1/2} + c = -\frac{2}{\sin \alpha} \sqrt{\cos \alpha + \cot x \sin \alpha} + c \\ &= -\frac{2}{\sin \alpha} \sqrt{\cos \alpha + \frac{\cos x}{\sin x} \sin \alpha} + c \\ &= -\frac{2}{\sin \alpha} \sqrt{\frac{\sin x \cos \alpha + \cos x \sin \alpha}{\sin x}} + c \\ &= -\frac{2}{\sin \alpha} \sqrt{\frac{\sin(x + \alpha)}{\sin x}} + c.\end{aligned}$$

19.  $\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}}, x \in [0, 1]$

$$\text{Sol. We know that } \sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x} = \frac{\pi}{2}$$

$$\therefore \cos^{-1} \sqrt{x} = \frac{\pi}{2} - \sin^{-1} \sqrt{x}$$

$$\begin{aligned}
 \therefore I &= \int \frac{\sin^{-1} \sqrt{x} - \left( \frac{\pi}{2} - \sin^{-1} \sqrt{x} \right)}{\frac{\pi}{2}} dx \\
 &= \frac{2}{\pi} \int \left( 2 \sin^{-1} \sqrt{x} - \frac{\pi}{2} \right) dx = \frac{4}{\pi} \int \sin^{-1} \sqrt{x} dx - \int 1 dx \\
 &= \frac{4}{\pi} \int \sin^{-1} \sqrt{x} dx - x + c \quad \dots(i)
 \end{aligned}$$

**Now let us evaluate  $\int \sin^{-1} \sqrt{x} dx$**

Put  $\sqrt{x} = \sin \theta$ .  $\therefore x = \sin^2 \theta$ .

Differentiating both sides,  $dx = 2 \sin \theta \cos \theta d\theta = \sin 2\theta d\theta$

$$\therefore \int \sin^{-1} \sqrt{x} dx = \int \sin^{-1}(\sin \theta) \cdot \sin 2\theta d\theta = \int \underset{\text{I}}{\theta} \underset{\text{II}}{\sin 2\theta} d\theta$$

Applying Product Rule

$$\begin{aligned}
 &= \theta \left( \frac{-\cos 2\theta}{2} \right) - \int 1 \cdot \left( \frac{-\cos 2\theta}{2} \right) d\theta \\
 &= -\frac{1}{2} \theta \cos 2\theta + \frac{1}{2} \int \cos 2\theta d\theta = -\frac{1}{2} \theta \cos 2\theta + \frac{1}{2} \frac{\sin 2\theta}{2} \\
 &= -\frac{1}{2} \theta (1 - 2 \sin^2 \theta) + \frac{1}{4} 2 \sin \theta \cos \theta \\
 &= -\frac{1}{2} \theta (1 - 2 \sin^2 \theta) + \frac{1}{2} \sin \theta \sqrt{1 - \sin^2 \theta}
 \end{aligned}$$

Putting  $\sin \theta = \sqrt{x}$

$$= -\frac{1}{2} (\sin^{-1} \sqrt{x}) (1 - 2x) + \frac{1}{2} \sqrt{x} \sqrt{1 - x}$$

Putting this value of  $\int \sin^{-1} \sqrt{x} dx$  in (i),

$$\begin{aligned}
 I &= \frac{4}{\pi} \left[ -\frac{1}{2} (1 - 2x) \sin^{-1} \sqrt{x} + \frac{1}{2} \sqrt{x} \sqrt{1 - x} \right] - x + c \\
 &= -\frac{2}{\pi} (1 - 2x) \sin^{-1} \sqrt{x} + \frac{2}{\pi} \sqrt{x} \sqrt{1 - x} - x + c.
 \end{aligned}$$

20.  $\sqrt{\frac{1 - \sqrt{x}}{1 + \sqrt{x}}}$

**Sol.** Let  $I = \int \sqrt{\frac{1 - \sqrt{x}}{1 + \sqrt{x}}} dx$

Put  $\sqrt{x} = t$ , i.e.,  $\sqrt{\text{Linear}} = t$ .  $\therefore x = t^2$

Differentiating both sides,  $dx = 2t dt$

$$\therefore I = \int \sqrt{\frac{1 - t}{1 + t}} 2t dt = 2 \int t \sqrt{\frac{1 - t}{1 + t}} dt$$

$$= 2 \int t \sqrt{\frac{1-t}{1+t} \times \frac{1-t}{1-t}} dt \quad (\text{Rationalising})$$

$$= 2 \int \frac{t(1-t)}{\sqrt{1-t^2}} dt = 2 \int \frac{t-t^2}{\sqrt{1-t^2}} dt \quad \dots(i)$$

$$= 2 \int \frac{(1-t^2) + t - 1}{\sqrt{1-t^2}} dt$$

$$= 2 \left[ \int \sqrt{1-t^2} dt + \int \frac{t}{\sqrt{1-t^2}} dt - \int \frac{1}{\sqrt{1-t^2}} dt \right]$$

$$= 2 \left[ \frac{t}{2} \sqrt{1-t^2} + \frac{1}{2} \sin^{-1} t + \int \frac{t}{\sqrt{1-t^2}} dt - \sin^{-1} t \right] + c$$

$$\left[ \because \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]$$

$$\text{or } I = 2 \left[ \frac{1}{2} t \sqrt{1-t^2} - \frac{1}{2} \sin^{-1} t + \int \frac{t}{\sqrt{1-t^2}} dt \right] + c \quad \dots(ii)$$

$$\text{To evaluate } \int \frac{t}{\sqrt{1-t^2}} dt$$

$$\text{Put } 1 - t^2 = z$$

$$\text{Differentiating both sides } -2t dt = dz \quad \text{or } t dt = -\frac{1}{2} dz.$$

$$\begin{aligned} \therefore \int \frac{t}{\sqrt{1-t^2}} dt &= \int \frac{-\frac{1}{2} dz}{\sqrt{z}} = -\frac{1}{2} \int z^{-1/2} dz \\ &= -\frac{1}{2} \frac{z^{1/2}}{\frac{1}{2}} = -\sqrt{1-t^2} \quad \dots(iii) \end{aligned}$$

$$\text{Putting the value of } \int \frac{t}{\sqrt{1-t^2}} dt = -\sqrt{1-t^2} \text{ from (iii) in (ii),}$$

$$\begin{aligned} \text{We have } I &= 2 \left[ \frac{1}{2} t \sqrt{1-t^2} - \frac{1}{2} \sin^{-1} t - \sqrt{1-t^2} \right] + c \\ &= t \sqrt{1-t^2} - \sin^{-1} t - 2\sqrt{1-t^2} + c \\ &= (t-2) \sqrt{1-t^2} - \sin^{-1} t + c \end{aligned}$$

$$\text{Putting } t = \sqrt{x} = (\sqrt{x} - 2) \sqrt{1-x} - \sin^{-1} \sqrt{x} + c.$$

**Remark.** Second method to integrate after arriving at equation

$$(i) \text{ namely } I = 2 \int \frac{t-t^2}{\sqrt{1-t^2}} dt, \text{ is put } t = \sin \theta.$$

21.  $\frac{2 + \sin 2x}{1 + \cos 2x} e^x$

**Sol.** Let  $I = \int \frac{2 + \sin 2x}{1 + \cos 2x} e^x dx = \int e^x \frac{(2 + 2 \sin x \cos x)}{2 \cos^2 x} dx$

$$= \int e^x \left( \frac{2}{2 \cos^2 x} + \frac{2 \sin x \cos x}{2 \cos^2 x} \right) dx$$

$$= \int e^x \left( \frac{1}{\cos^2 x} + \frac{\sin x}{\cos x} \right) dx = \int e^x (\sec^2 x + \tan x) dx$$

$$= \int e^x (\tan x + \sec^2 x) dx = \int e^x (f(x) + f'(x)) dx$$

where  $f(x) = \tan x$  and  $f'(x) = \sec^2 x$

$$= e^x f(x) + c = e^x \tan x + c. \quad \left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) + c \right]$$

22.  $\frac{x^2 + x + 1}{(x+1)^2 (x+2)}$

**Sol.** Let  $I = \int \frac{x^2 + x + 1}{(x+1)^2 (x+2)} dx \quad \dots(i)$

The integrand  $\frac{x^2 + x + 1}{(x+1)^2 (x+2)}$  is a rational function of  $x$  and degree of numerator is less than degree of denominator. So we can form partial fractions of integrand.

Let integrand  $\frac{x^2 + x + 1}{(x+1)^2 (x+2)} = \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{x+2} \quad \dots(ii)$

Multiplying both sides of (ii) L.C.M. =  $(x+1)^2 (x+2)$ , we have

$$\begin{aligned} x^2 + x + 1 &= A(x+1)(x+2) + B(x+2) + C(x+1)^2 \\ \text{or } x^2 + x + 1 &= A(x^2 + 3x + 2) + B(x+2) + C(x^2 + 1 + 2x) \\ &= Ax^2 + 3Ax + 2A + Bx + 2B + Cx^2 + C + 2Cx \end{aligned}$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides, we have

$$x^2: \quad A + C = 1 \quad \dots(iii)$$

$$x: \quad 3A + B + 2C = 1 \quad \dots(iv)$$

$$\text{Constant terms: } 2A + 2B + C = 1 \quad \dots(v)$$

Let us solve Eqns. (iii), (iv) and (v) for A, B, C.

Eqn. (iv) - 2 × Eqn. (iii) gives (to eliminate C)

$$3A + B + 2C - 2A - 2C = 1 - 2$$

$$\text{or } A + B = -1 \quad \dots(vi)$$

Eqn. (v) - Eqn. (iii) gives (To eliminate C)

$$A + 2B = 0 \quad \dots(vii)$$

$$\text{Eqn. (vii) - Eqn. (vi) gives } B = 0 + 1 = 1.$$

$$\text{Putting } B = 1 \text{ in (vi), } A + 1 = -1 \Rightarrow A = -2$$

$$\text{Putting } A = -2 \text{ in (iii), } -2 + C = 1 \Rightarrow C = 3$$

Putting values of A, B, C in (ii)

$$\begin{aligned}
 \frac{x^2 + x + 1}{(x+1)^2(x+2)} &= \frac{-2}{x+1} + \frac{1}{(x+1)^2} + \frac{3}{x+2} \\
 \therefore \int \frac{x^2 + x + 1}{(x+1)^2(x+2)} dx \\
 &= -2 \int \frac{1}{x+1} dx + \int (x+1)^{-2} dx + 3 \int \frac{1}{x+2} dx \\
 &= -2 \log |x+1| + \frac{(x+1)^{-2+1}}{-2+1} + 3 \log |x+2| + c \\
 &= -2 \log |x+1| - \frac{1}{x+1} + 3 \log |x+2| + c \left( \because \frac{(x+1)^{-1}}{-1} = \frac{-1}{x+1} \right)
 \end{aligned}$$

**Evaluate the integrals in Exercises 23 and 24:**

**23.**  $\tan^{-1} \sqrt{\frac{1-x}{1+x}}$

**Sol.** Let  $I = \int \tan^{-1} \sqrt{\frac{1-x}{1+x}} dx \quad \dots(i)$

**Put**  $x = \cos 2\theta \Rightarrow \frac{dx}{d\theta} = -2 \sin 2\theta$   
 $\Rightarrow dx = -2 \sin 2\theta d\theta$

and  $\tan^{-1} \sqrt{\frac{1-x}{1+x}} = \tan^{-1} \sqrt{\frac{1-\cos 2\theta}{1+\cos 2\theta}} = \tan^{-1} \sqrt{\frac{2 \sin^2 \theta}{2 \cos^2 \theta}}$   
 $= \tan^{-1} \sqrt{\tan^2 \theta} = \tan^{-1} \tan \theta = \theta$

$\therefore$  From (i),  $I = \int \theta (-2 \sin 2\theta d\theta) = -2 \int \theta \sin 2\theta d\theta$   
I      II

Applying Product Rule of Integration,

$$\begin{aligned}
 \left( \int I \cdot II \, dx = I \int II \, dx - \int \left( \frac{d}{dx} (I) \int II \, dx \right) dx \right) \\
 I = -2 \left[ \theta \left( \frac{-\cos 2\theta}{2} \right) - \int 1 \left( \frac{-\cos 2\theta}{2} \right) d\theta \right] \\
 = -2 \left[ \frac{-1}{2} \theta \cos 2\theta + \frac{1}{2} \int \cos 2\theta d\theta \right] = \theta \cos 2\theta - \frac{\sin 2\theta}{2} + c \\
 = \theta \cos 2\theta - \frac{1}{2} \sqrt{1 - \cos^2 2\theta} + c \left( \because \sin^2 \alpha + \cos^2 \alpha = 1 \right) \\
 = \frac{1}{2} (\cos^{-1} x) x - \frac{1}{2} \sqrt{1 - x^2} + c \\
 \left[ \because \cos 2\theta = x \Rightarrow 2\theta = \cos^{-1} x \Rightarrow \theta = \frac{1}{2} \cos^{-1} x \right] \\
 = \frac{1}{2} x \cos^{-1} x - \frac{1}{2} \sqrt{1 - x^2} + c \\
 = \frac{1}{2} [x \cos^{-1} x - \sqrt{1 - x^2}] + c.
 \end{aligned}$$

$$24. \frac{\sqrt{x^2+1} [\log(x^2+1) - 2 \log x]}{x^4}$$

$$\text{Sol. } I = \int \frac{\sqrt{x^2+1} [\log(x^2+1) - 2 \log x]}{x^4} dx$$

$$= \int \frac{\sqrt{x^2+1}}{x^4} [\log(x^2+1) - \log x^2] dx$$

$$= \int \frac{\sqrt{x^2 \left(1 + \frac{1}{x^2}\right)}}{x^4} \log \left( \frac{x^2+1}{x^2} \right) dx$$

$$= \int \frac{\sqrt{1 + \frac{1}{x^2}}}{x^3} \log \left( 1 + \frac{1}{x^2} \right) dx = \int \sqrt{1 + \frac{1}{x^2}} \log \left( 1 + \frac{1}{x^2} \right) \cdot \frac{dx}{x^3}$$

$$\text{Put } 1 + \frac{1}{x^2} = t \quad \text{or} \quad 1 + x^{-2} = t.$$

$$\text{Differentiating both sides, } -\frac{2}{x^3} dx = dt \quad \text{or} \quad \frac{dx}{x^3} = -\frac{1}{2} dt$$

$$\therefore I = -\frac{1}{2} \int \sqrt{t} \log t \, dt = -\frac{1}{2} \int (\log t) \cdot t^{1/2} dt$$

$\quad \quad \quad I \quad \quad \quad II$

Integrating by Product Rule,

$$= -\frac{1}{2} \left[ (\log t) \cdot \frac{t^{3/2}}{3/2} - \int \frac{1}{t} \cdot \frac{t^{3/2}}{3/2} dt \right] = -\frac{1}{3} t^{3/2} \log t + \frac{1}{3} \int t^{1/2} dt$$

$$= -\frac{1}{3} t^{3/2} \log t + \frac{1}{3} \cdot \frac{t^{3/2}}{3/2} + c$$

$$= \frac{2}{9} t^{3/2} - \frac{1}{3} t^{3/2} \log t + c = \frac{1}{3} t^{3/2} \left[ \frac{2}{3} - \log t \right] + c$$

$$\text{Putting } t = 1 + \frac{1}{x^2}, \text{ we have } = \frac{1}{3} \left( 1 + \frac{1}{x^2} \right)^{3/2} \left[ \frac{2}{3} - \log \left( 1 + \frac{1}{x^2} \right) \right] + c.$$

**Evaluate the definite integrals in Exercises 25 to 33:**

$$25. \int_{\frac{\pi}{2}}^{\pi} e^x \left( \frac{1 - \sin x}{1 - \cos x} \right) dx$$

$$\text{Sol. Let } I = \int_{\frac{\pi}{2}}^{\pi} e^x \left( \frac{1 - \sin x}{1 - \cos x} \right) dx = \int_{\frac{\pi}{2}}^{\pi} e^x \left[ \frac{1 - 2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \sin^2 \frac{x}{2}} \right] dx$$

$$= \int_{\frac{\pi}{2}}^{\pi} e^x \left[ \frac{1}{2 \sin^2 \frac{x}{2}} - \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \sin^2 \frac{x}{2}} \right] dx = \int_{\frac{\pi}{2}}^{\pi} e^x \left[ \frac{1}{2} \operatorname{cosec}^2 \frac{x}{2} - \cot \frac{x}{2} \right] dx$$

$$= \int_{\frac{\pi}{2}}^{\pi} e^x \left[ -\cot \frac{x}{2} + \frac{1}{2} \operatorname{cosec}^2 \frac{x}{2} \right] dx = \int_{\frac{\pi}{2}}^{\pi} e^x (f(x) + f'(x)) dx$$

where  $f(x) = -\cot \frac{x}{2}$ . Therefore  $f'(x) = \frac{1}{2} \operatorname{cosec}^2 \frac{x}{2}$

$$= \left( e^x f(x) \right)_{\frac{\pi}{2}}^{\pi} = \left( -e^x \cot \frac{x}{2} \right)_{\frac{\pi}{2}}^{\pi} \quad \left[ \because \int e^x (f(x) + f'(x)) dx = e^x f(x) \right]$$

$$= -e^{\pi} \cot \frac{\pi}{2} - \left( -e^{\frac{\pi}{2}} \cot \frac{\pi}{4} \right)$$

$$= -e^{\pi} (0) + e^{\pi/2} (1) \quad \left[ \because \cot \frac{\pi}{2} = \frac{\cos \frac{\pi}{2}}{\sin \frac{\pi}{2}} = \frac{0}{1} = 0 \right]$$

$$= e^{\pi/2}.$$

26.  $\int_0^{\frac{\pi}{4}} \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{4}} \frac{\sin x \cos x}{\cos^4 x + \sin^4 x} dx$

Dividing every term by  $\cos^4 x$ ,

$$I = \int_0^{\frac{\pi}{4}} \frac{\frac{\sin x \cos x}{\cos^4 x}}{1 + \frac{\sin^4 x}{\cos^4 x}} dx = \int_0^{\frac{\pi}{4}} \frac{\tan x \sec^2 x}{1 + \tan^4 x} dx$$

Dividing and multiplying by 2,

$$I = \frac{1}{2} \int_0^{\frac{\pi}{4}} \frac{2 \tan x \sec^2 x}{1 + \tan^4 x} dx \quad \dots(i)$$

**Put  $\tan^2 x = t$ .**

$$\therefore 2 \tan x \frac{d}{dx} (\tan x) = \frac{dt}{dx} \Rightarrow 2 \tan x \sec^2 x dx = dt.$$

**To change the limits of integration**

When  $x = 0$ ,  $t = \tan^2 x = \tan^2 0 = 0$

When  $x = \frac{\pi}{4}$ ,  $t = \tan^2 \frac{\pi}{4} = 1$

$$\therefore \text{From (i), } I = \frac{1}{2} \int_0^1 \frac{dt}{1+t^2} = \frac{1}{2} \left( \tan^{-1} t \right)_0^1$$

$$= \frac{1}{2} (\tan^{-1} 1 - \tan^{-1} 0) = \frac{1}{2} \left( \frac{\pi}{4} - 0 \right) = \frac{\pi}{8}. \quad \left[ \because \tan \frac{\pi}{4} = 1 \right]$$

$$27. \int_0^{\frac{\pi}{2}} \frac{\cos^2 x \, dx}{\cos^2 x + 4 \sin^2 x}$$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{\cos^2 x + 4 \sin^2 x} \, dx$

Dividing every term of integrand by  $\cos^2 x$ ,

$$I = \int_0^{\frac{\pi}{2}} \frac{1}{(1 + 4 \tan^2 x)} \, dx \quad \dots(i)$$

**Put  $\tan x = t$ .**

$$\therefore \sec^2 x = \frac{dt}{dx} \Rightarrow \sec^2 x \, dx = dt$$

$$\Rightarrow dx = \frac{dt}{\sec^2 x} = \frac{dt}{1 + \tan^2 x} = \frac{dt}{1 + t^2}$$

**To change the limits:**

When  $x = 0$ ,  $t = \tan 0 = 0$

When  $x = \frac{\pi}{2}$ ,  $t = \tan \frac{\pi}{2} = \infty$

$$\begin{aligned} \therefore \text{From (i), } I &= \int_0^{\infty} \frac{1}{1 + 4t^2} \cdot \frac{dt}{1 + t^2} \\ &= \int_0^{\infty} \frac{1}{(4t^2 + 1)(t^2 + 1)} \, dt \quad \dots(ii) \end{aligned}$$

Put  $t^2 = y$  only in the integrand of (ii) to form partial fractions.

The new integrand is  $\frac{1}{(4y + 1)(y + 1)}$

$$\text{Let } \frac{1}{(4y + 1)(y + 1)} = \frac{A}{4y + 1} + \frac{B}{y + 1} \quad \dots(iii)$$

Multiplying by L.C.M. =  $(4y + 1)(y + 1)$

$$1 = A(y + 1) + B(4y + 1)$$

$$\text{or } 1 = Ay + A + 4By + B$$

$$\text{Comparing coefficient of } y \text{ on both sides, } A + 4B = 0 \quad \dots(iv)$$

$$\text{Comparing constants, } A + B = 1 \quad \dots(v)$$

$$(iv) - (v) \text{ gives } 3B = -1 \Rightarrow B = -\frac{1}{3}$$

$$\therefore \text{From (iv) } A = -4B = -4 \left( -\frac{1}{3} \right) = \frac{4}{3}$$

Putting values of A, B and y in (iii), we have

$$\frac{1}{(4t^2 + 1)(t^2 + 1)} = \frac{\frac{4}{3}}{4t^2 + 1} - \frac{\frac{1}{3}}{t^2 + 1} = \frac{1}{3} \left( \frac{4}{(4t^2 + 1)} - \frac{1}{(t^2 + 1)} \right)$$

Putting this value in (ii)

$$\begin{aligned}
 I &= \frac{1}{3} \left[ 4 \int_0^\infty \frac{1}{(4t^2+1)} dt - \int_0^\infty \frac{1}{t^2+1} dt \right] \\
 &= \frac{1}{3} \left[ 4 \int_0^\infty \frac{1}{(2t)^2+1^2} dt - \left( \tan^{-1} t \right)_0^\infty \right] \\
 &= \frac{1}{3} \left[ 4 \frac{\left( \frac{1}{1} \tan^{-1} \frac{2t}{1} \right)_0^\infty}{2 \rightarrow \text{Coeff. of } t} - \left( \tan^{-1} t \right)_0^\infty \right] \\
 &= \frac{1}{3} [2 (\tan^{-1} \infty - \tan^{-1} 0) - (\tan^{-1} \infty - \tan^{-1} 0)] \\
 &= \frac{1}{3} \left[ 2 \cdot \left( \frac{\pi}{2} - 0 \right) - \left( \frac{\pi}{2} - 0 \right) \right] = \frac{1}{3} \left( \frac{2\pi}{2} - \frac{\pi}{2} \right) = \frac{1}{3} \times \frac{\pi}{2} = \frac{\pi}{6}.
 \end{aligned}$$

28.  $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$

**Sol.** Let  $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$  ... (i)

**Put  $\sin x - \cos x = t$ .** Differentiating both sides w.r.t.  $x$ ,

$$(\cos x + \sin x) dx = dt$$

Also, squaring  $\sin^2 x + \cos^2 x - 2 \sin x \cos x = t^2$

$$\Rightarrow 1 - \sin 2x = t^2$$

$$\Rightarrow \sin 2x = 1 - t^2$$

**To change the limits of Integration**

When  $x = \frac{\pi}{6}$ ,  $t = \sin \frac{\pi}{6} - \cos \frac{\pi}{6}$

$$= \frac{1}{2} - \frac{\sqrt{3}}{2} = \frac{1-\sqrt{3}}{2} = \frac{-(\sqrt{3}-1)}{2} = -\alpha \text{ (say)}$$

where  $\alpha = \frac{\sqrt{3}-1}{2}$

... (ii)

When  $x = \frac{\pi}{3}$ ,  $t = \sin \frac{\pi}{3} - \cos \frac{\pi}{3} = \frac{\sqrt{3}}{2} - \frac{1}{2} = \frac{\sqrt{3}-1}{2} = \alpha$

$$\therefore \text{ From (i), } I = \int_{-\alpha}^{\alpha} \frac{dt}{\sqrt{1-t^2}} = \left[ \sin^{-1} t \right]_{-\alpha}^{\alpha}$$

$$= \sin^{-1} \alpha - \sin^{-1} (-\alpha)$$

$$= \sin^{-1} \alpha + \sin^{-1} \alpha = 2 \sin^{-1} \left( \frac{\sqrt{3}-1}{2} \right). \quad [\text{By (ii)}]$$

29.  $\int_0^1 \frac{dx}{\sqrt{1+x-\sqrt{x}}}$

**Sol.** Let  $I = \int_0^1 \frac{1}{\sqrt{1+x-\sqrt{x}}} dx$

$$\begin{aligned}
 \text{Rationalising} &= \int_0^1 \frac{\sqrt{1+x} + \sqrt{x}}{(\sqrt{1+x} + \sqrt{x})(\sqrt{1+x} - \sqrt{x})} dx \\
 &= \int_0^1 \frac{\sqrt{1+x} + \sqrt{x}}{1+x-x} dx = \int_0^1 (\sqrt{1+x} + \sqrt{x}) dx \quad (\because 1+x-x=1) \\
 &= \int_0^1 (1+x)^{1/2} dx + \int_0^1 x^{1/2} dx = \left[ \frac{(1+x)^{3/2}}{3/2} \right]_0^1 + \left[ \frac{x^{3/2}}{3/2} \right]_0^1 \\
 &= \frac{2}{3} [(2)^{3/2} - (1)^{3/2}] + \frac{2}{3} [(1)^{3/2} - 0] = \frac{2}{3} (2\sqrt{2} - 1) + \frac{2}{3} (1 - 0) \\
 &\quad \left[ \because x^{3/2} = x^{\frac{2+1}{2}} = x^{1+\frac{1}{2}} = x^1 \cdot x^{\frac{1}{2}} = x\sqrt{x} \right] \\
 &= \frac{4\sqrt{2}}{3} - \frac{2}{3} + \frac{2}{3} = \frac{4\sqrt{2}}{3}.
 \end{aligned}$$

30.  $\int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{9 + 16 \sin 2x} dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{9 + 16 \sin 2x} dx$

**Put  $\sin x - \cos x = t$ .** Differentiating both sides

$$(\cos x + \sin x) dx = dt$$

Also  $(\sin x - \cos x)^2 = t^2 \therefore \sin^2 x + \cos^2 x - 2 \sin x \cos x = t^2$   
or  $1 - t^2 = \sin 2x$

**Let us change the limits of Integration**

When  $x = 0$ ,  $t = 0 - 1 = -1$

When  $x = \frac{\pi}{4}$ ,  $t = \sin \frac{\pi}{4} - \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = 0$

$$\begin{aligned}
 \therefore I &= \int_{-1}^0 \frac{dt}{9 + 16(1-t^2)} = \int_{-1}^0 \frac{dt}{25 - 16t^2} \\
 &= \int_{-1}^0 \frac{dt}{16 \left( \frac{25}{16} - t^2 \right)} = \frac{1}{16} \int_{-1}^0 \frac{dt}{\left( \frac{5}{4} \right)^2 - t^2} \\
 &= \frac{1}{16} \times \left[ \frac{1}{2 \times 5/4} \log \left| \frac{5/4 + t}{5/4 - t} \right| \right]_{-1}^0 \left[ \because \int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| \right] \\
 &= \frac{1}{40} \left[ \log 1 - \log \frac{1/4}{9/4} \right] = \frac{1}{40} \left[ 0 - \log \frac{1}{9} \right] \\
 &= \frac{1}{40} [- (\log 1 - \log 9)] = \frac{1}{40} \log 9
 \end{aligned}$$

$$= \frac{1}{40} \log 3^2 = \frac{2}{40} \log 3 = \frac{1}{20} \log 3.$$

31.  $\int_0^{\frac{\pi}{2}} \sin 2x \tan^{-1}(\sin x) dx$

**Sol.** Let  $I = \int_0^{\frac{\pi}{2}} \sin 2x \tan^{-1}(\sin x) dx = \int_0^{\frac{\pi}{2}} 2 \sin x \cos x \tan^{-1}(\sin x) dx$

**Put  $\sin x = t$ .** Differentiating both sides  $\cos x dx = dt$

**To change the limits of Integration**

When  $x = 0, t = 0$

When  $x = \frac{\pi}{2}, t = \sin \frac{\pi}{2} = 1 \quad \therefore I = 2 \int_0^1 t \tan^{-1} t dt \quad \dots(i)$

Now  $\int t \tan^{-1} t dt = \int (\tan^{-1} t) t dt$  Integrating by parts

$$\begin{aligned} &= \tan^{-1} t \cdot \frac{t^2}{2} - \int \frac{1}{1+t^2} \cdot \frac{t^2}{2} dt \\ &= \frac{t^2}{2} \tan^{-1} t - \frac{1}{2} \int \frac{(1+t^2)-1}{1+t^2} dt \\ &= \frac{t^2}{2} \tan^{-1} t - \frac{1}{2} \int \left( 1 - \frac{1}{1+t^2} \right) dt = \frac{t^2}{2} \tan^{-1} t - \frac{1}{2} (t - \tan^{-1} t) \\ &= \frac{t^2}{2} \tan^{-1} t - \frac{1}{2} t + \frac{1}{2} \tan^{-1} t + c = \frac{1}{2} [(t^2 + 1) \tan^{-1} t - t] \end{aligned}$$

From (i),  $I = 2 \left[ \frac{1}{2} \{(t^2 + 1) \tan^{-1} t - t\} \right]_0^1 = (2 \tan^{-1} 1 - 1) - (0 - 0)$

$$= 2 \times \frac{\pi}{4} - 1 = \frac{\pi}{2} - 1.$$

32.  $\int_0^{\pi} \frac{x \tan x}{\sec x + \tan x} dx$

**Sol.** Let  $I = \int_0^{\pi} \frac{x \tan x}{\sec x + \tan x} dx = \int_0^{\pi} \frac{x \frac{\sin x}{\cos x}}{\frac{1}{\cos x} + \frac{\sin x}{\cos x}} dx$

$$= \int_0^{\pi} \frac{x \sin x}{1 + \sin x} dx \quad \dots(ii)$$

Using  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

$\therefore I = \int_0^{\pi} \frac{(\pi-x) \sin(\pi-x)}{1 + \sin(\pi-x)} dx = \int_0^{\pi} \frac{(\pi-x) \sin x}{1 + \sin x} dx \quad \dots(ii)$

Adding Eqns. (i) and (ii), we have

$$2I = \int_0^{\pi} \frac{x \sin x + (\pi-x) \sin x}{1 + \sin x} dx = \int_0^{\pi} \frac{x \sin x + \pi \sin x - x \sin x}{1 + \sin x} dx$$

$$\begin{aligned}
&= \int_0^\pi \frac{\pi \sin x}{1 + \sin x} dx = \pi \int_0^\pi \frac{\sin x}{1 + \sin x} dx \\
\text{or } 2I &= \pi \int_0^\pi \frac{(1 + \sin x) - 1}{1 + \sin x} dx \\
\Rightarrow 2I &= \pi \int_0^\pi \left( 1 - \frac{1}{1 + \sin x} \right) dx = \pi \int_0^\pi dx - \pi \int_0^\pi \frac{dx}{1 + \sin x} \\
&= \pi \left[ x \right]_0^\pi - 2\pi \int_0^{\frac{\pi}{2}} \frac{dx}{1 + \sin x} \\
&\quad \left[ \because \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(2a - x) = f(x) \right] \\
&= \pi(\pi) - 2\pi \int_0^{\frac{\pi}{2}} \frac{dx}{1 + \sin\left(\frac{\pi}{2} - x\right)} = \pi^2 - 2\pi \int_0^{\frac{\pi}{2}} \frac{dx}{1 + \cos x} \\
&= \pi^2 - 2\pi \int_0^{\frac{\pi}{2}} \frac{dx}{2 \cos^2 \frac{x}{2}} = \pi^2 - \pi \int_0^{\frac{\pi}{2}} \sec^2 \frac{x}{2} dx \\
\text{or } 2I &= \pi^2 - \pi \left[ \frac{\tan \frac{x}{2}}{\frac{1}{2}} \right]_0^{\pi/2} = \pi^2 - 2\pi(1)
\end{aligned}$$

Dividing both sides by 2,  $I = \frac{\pi^2}{2} - \pi = \pi \left( \frac{\pi}{2} - 1 \right) = \pi \left( \frac{\pi - 2}{2} \right)$ .

33.  $\int_1^4 [|x - 1| + |x - 2| + |x - 3|] dx$

**Sol.** Let  $I = \int_1^4 (|x - 1| + |x - 2| + |x - 3|) dx$  ... (i)

Putting each expression within modulus equal to 0, we have

$$x - 1 = 0, x - 2 = 0, x - 3 = 0 \quad \text{i.e.,} \quad x = 1, x = 2, x = 3$$

Here 2 and 3  $\in$  (1, 4)

$$\begin{aligned}
\therefore \text{ From (i), } I &= \int_1^2 (|x - 1| + |x - 2| + |x - 3|) dx \\
&+ \int_2^3 (|x - 1| + |x - 2| + |x - 3|) dx + \int_3^4 (|x - 1| + |x - 2| + |x - 3|) dx \\
&\quad \dots (ii)
\end{aligned}$$

$$\left[ \because \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^d f(x) dx + \int_d^b f(x) dx \quad \text{where } a < c < d < b \right]$$

Let  $I_1 = \int_1^2 (|x - 1| + |x - 2| + |x - 3|) dx$

On this interval (1, 2) (for example taking  $x = 1.3$ ;  $(x - 1)$  is positive,  $(x - 2)$  is negative and  $(x - 3)$  is negative and hence

$$|x - 1| = (x - 1), |x - 2| = -(x - 2) \text{ and } |x - 3| = -(x - 3).$$

$$\begin{aligned}
 \text{Therefore } I_1 &= \int_1^2 ((x-1) - (x-2) - (x-3)) \, dx \\
 &= \int_1^2 (x-1-x+2-x+3) \, dx = \int_1^2 (4-x) \, dx \\
 &= \left( 4x - \frac{x^2}{2} \right)_1^2 = (8-2) - \left( 4 - \frac{1}{2} \right) \\
 &= 6-4 + \frac{1}{2} = 2 + \frac{1}{2} = \frac{5}{2} \quad \dots(iii)
 \end{aligned}$$

Let  $I_2 = \int_2^3 (|x-1| + |x-2| + |x-3|) \, dx$   
 On this interval (2, 3) (for example taking  $x = 2.8$ ;  $(x-1)$  is positive,  $(x-2)$  is positive and  $(x-3)$  is negative and hence  $|x-1| = x-1$ ,  $|x-2| = x-2$  and  $|x-3| = -(x-3)$ )

$$\begin{aligned}
 \text{Therefore } I_2 &= \int_2^3 ((x-1) + (x-2) - (x-3)) \, dx = \int_2^3 (2x-3-x+3) \, dx \\
 &= \int_2^3 x \, dx = \left( \frac{x^2}{2} \right)_2^3 = \frac{9}{2} - \frac{4}{2} = \frac{5}{2} \quad \dots(iv)
 \end{aligned}$$

Let  $I_3 = \int_3^4 (|x-1| + |x-2| + |x-3|) \, dx$   
 On this interval (3, 4), (for example taking  $x = 3.4$ ;  $(x-1)$  is positive,  $(x-2)$  is positive and  $(x-3)$  is positive and hence  $|x-1| = x-1$ ,  $|x-2| = x-2$  and  $|x-3| = x-3$ )

$$\begin{aligned}
 \text{Therefore } I_3 &= \int_3^4 (x-1+x-2+x-3) \, dx = \int_3^4 (3x-6) \, dx \\
 &= \left( \frac{3x^2}{2} - 6x \right)_3^4 = (24-24) - \left( \frac{27}{2} - 18 \right) \\
 &= 0 - \left( \frac{27-36}{2} \right) = - \left( -\frac{9}{2} \right) = \frac{9}{2} \quad \dots(v)
 \end{aligned}$$

Putting values of  $I_1$ ,  $I_2$ ,  $I_3$  from (iii), (iv) and (v) in (ii),

$$I = \frac{5}{2} + \frac{5}{2} + \frac{9}{2} = \frac{19}{2}.$$

**Prove the following (Exercises 34 to 40):**

$$34. \int_1^3 \frac{dx}{x^2(x+1)} = \frac{2}{3} + \log \frac{2}{3}$$

$$\text{Sol. Let } I = \int_1^3 \frac{dx}{x^2(x+1)} = \int_1^3 \frac{1}{x^2(x+1)} \, dx \quad \dots(i)$$

$$\text{Let integrand } \frac{1}{x^2(x+1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x+1} \quad \dots(ii)$$

(Partial fractions)

Multiplying by L.C.M. =  $x^2(x+1)$

$$1 = Ax(x+1) + B(x+1) + Cx^2$$

$$\Rightarrow 1 = Ax^2 + Ax + Bx + B + Cx^2$$

Comparing coefficients of  $x^2$ ,  $x$  and constant terms on both sides, we have

$$x^2: \quad A + C = 0 \quad \dots(iii)$$

$$x: \quad A + B = 0 \quad \dots(iv)$$

$$\text{Constants:} \quad B = 1 \quad \dots(v)$$

Let us solve (iii), (iv), (v) for A, B, C.

Putting B = 1 from (v) in (iv),  $A + 1 = 0$  or  $A = -1$

Putting A = -1 in (iii),  $-1 + C = 0 \Rightarrow C = 1$

Putting values of A, B, C in (ii),

$$\frac{1}{x^2(x+1)} = \frac{-1}{x} + \frac{1}{x^2} + \frac{1}{x+1}$$

$$\therefore \text{From (i), } I = \int_1^3 \frac{dx}{x^2(x+1)}$$

$$= - \int_1^3 \frac{1}{x} dx + \int_1^3 \frac{1}{x^2} dx + \int_1^3 \frac{1}{x+1} dx$$

$$= - (\log |x|)_1^3 + \int_1^3 x^{-2} dx + (\log |x+1|)_1^3$$

$$= - (\log |3| - \log |1|) + \left( \frac{x^{-1}}{-1} \right)_1^3 + (\log |4| - \log |2|)$$

$$= - \log 3 + 0 - \left( \frac{1}{x} \right)_1^3 + \log 4 - \log 2$$

$$= - \log 3 - \left( \frac{1}{3} - 1 \right) + \log 2^2 - \log 2$$

$$= - \log 3 - \left( \frac{1-3}{3} \right) + 2 \log 2 - \log 2$$

$$= - \log 3 + \frac{2}{3} + \log 2 = \frac{2}{3} + \log 2 - \log 3$$

$$= \frac{2}{3} + \log \frac{2}{3}.$$

$$35. \int_0^1 x e^x dx = 1$$

$$\text{Sol. } \int_0^1 x e^x$$

I II

Applying Product Rule of definite Integration

$$\begin{aligned} \left( \int I \cdot II dx = \left( I \int II dx \right)_a^b - \int_a^b \left( \frac{d}{dx} (I) \int II dx \right) dx \right) \\ = \left( x e^x \right)_0^1 - \int_0^1 1 \cdot e^x dx \\ = e - 0 - \int_0^1 e^x dx = e - \left( e^x \right)_0^1 = e - (e - e^0) = e - e + e^0 = 1. \end{aligned}$$

$$36. \int_{-1}^1 x^{17} \cos^4 x \, dx = 0$$

$$\text{Sol. Let } I = \int_{-1}^1 x^{17} \cos^4 x \, dx \quad \dots(i)$$

Here the integrand  $f(x) = x^{17} \cos^4 x$

$$\begin{aligned} \therefore f(-x) &= (-x)^{17} \cos^4(-x) \\ &= -x^{17} \cos^4 x = -f(x) \end{aligned}$$

$\therefore f(x)$  is an odd function of  $x$ .

$$\therefore \text{From (i), } I = \int_{-1}^1 x^{17} \cos^4 x \, dx = 0$$

[ $\because$  If  $f(x)$  is an odd function of  $x$ , then  $\int_{-a}^a f(x) \, dx = 0$ ]

$$37. \int_0^{\frac{\pi}{2}} \sin^3 x \, dx = \frac{2}{3}$$

$$\text{Sol. } \int_0^{\frac{\pi}{2}} \sin^3 x \, dx = \int_0^{\frac{\pi}{2}} \frac{1}{4} (3 \sin x - \sin 3x) \, dx$$

$$\left[ \because \sin 3A = 3 \sin A - 4 \sin^3 A \Rightarrow \sin^3 A = \frac{1}{4} (3 \sin A - \sin 3A) \right]$$

$$= \frac{1}{4} \left[ 3(-\cos x) - \left( -\frac{\cos 3x}{3} \right) \right]_0^{\pi/2} = \frac{1}{4} \left( -3 \cos x + \frac{1}{3} \cos 3x \right)_0^{\pi/2}$$

$$= \frac{1}{4} \left[ \left( -3 \cos \frac{\pi}{2} + \frac{1}{3} \cos \frac{3\pi}{2} \right) - \left( -3 \cos 0 + \frac{1}{3} \cos 0 \right) \right]$$

$$= \frac{1}{4} \left[ -3 \times 0 + \frac{1}{3} \times 0 + 3 \times 1 - \frac{1}{3} \times 1 \right] = \frac{1}{4} \left( 3 - \frac{1}{3} \right)$$

$$= \frac{1}{4} \times \frac{8}{3} = \frac{2}{3}.$$

$$\left[ \because \cos \frac{3\pi}{2} = \cos 270^\circ = \cos (180^\circ + 90^\circ) = -\cos 90^\circ = 0 \right]$$

$$38. \int_0^{\frac{\pi}{4}} 2 \tan^3 x \, dx = 1 - \log 2$$

$$\text{Sol. Let } I = \int_0^{\frac{\pi}{4}} 2 \tan^3 x \, dx = 2 \int_0^{\frac{\pi}{4}} \tan x \cdot \tan^2 x \, dx$$

Replacing  $\tan^2 x$  by  $(\sec^2 x - 1)$  in the integrand,

$$I = 2 \int_0^{\frac{\pi}{4}} \tan x (\sec^2 x - 1) \, dx = 2 \left[ \int_0^{\frac{\pi}{4}} (\tan x \sec^2 x - \tan x) \, dx \right]$$

$$= 2 \left[ \int_0^{\frac{\pi}{4}} \tan x \sec^2 x \, dx - \int_0^{\frac{\pi}{4}} \tan x \, dx \right] \quad \dots(i)$$

$$\text{Let } I_1 = \int_0^{\frac{\pi}{4}} \tan x \sec^2 x \, dx$$

**Put  $\tan x = t$ .** Therefore  $\sec^2 x = \frac{dt}{dx} \therefore \sec^2 x \, dx = dt$

**To change the limits of Integration**

When  $x = 0$ ,  $t = \tan x = \tan 0 = 0$

When  $x = \frac{\pi}{4}$ ,  $t = \tan \frac{\pi}{4} = 1$

$$\therefore I_1 = \int_0^1 t \, dt = \left( \frac{t^2}{2} \right)_0^1 = \frac{1}{2} - 0 = \frac{1}{2}$$

Putting this value of  $I_1$  in (i),

$$\begin{aligned} I &= 2 \left[ \frac{1}{2} - \left( \log |\sec x| \right)_0^{\pi/4} \right] = 1 - 2 \left( \log \sec \frac{\pi}{4} - \log \sec 0 \right) \\ &= 1 - 2 (\log \sqrt{2} - \log 1) = 1 - 2 (\log 2^{1/2} - 0) \\ &= 1 - 2 \left( \frac{1}{2} \log 2 \right) = 1 - \log 2. \end{aligned}$$

39.  $\int_0^1 \sin^{-1} x \, dx = \frac{\pi}{2} - 1$

**Sol.** Put  $x = \sin \theta$ . Differentiating both sides  $dx = \cos \theta \, d\theta$

**To change the limits of Integration**

When  $x = 0$ ,  $\theta = 0$ ,

When  $x = 1$ ,  $\sin \theta = 1$  and therefore  $\theta = \frac{\pi}{2}$

$$\therefore \int_0^1 \sin^{-1} x \, dx = \int_0^{\frac{\pi}{2}} \theta \cos \theta \, d\theta$$

Integrating by parts

$$\begin{aligned} &= \left[ \theta \sin \theta \right]_0^{\pi/2} - \int_0^{\pi/2} 1 \cdot \sin \theta \, d\theta = \left( \frac{\pi}{2} - 0 \right) + \left[ \cos \theta \right]_0^{\pi/2} \\ &= \frac{\pi}{2} + \left( \cos \frac{\pi}{2} - \cos 0 \right) = \frac{\pi}{2} + (0 - 1) = \frac{\pi}{2} - 1. \end{aligned}$$

40. Evaluate  $\int_0^1 e^{2-3x} \, dx$  as a limit of a sum.

**Sol. Step I.** Comparing  $\int_0^1 e^{2-3x} \, dx$  with  $\int_a^b f(x) \, dx$ , we have

$$a = 0, b = 1, f(x) = e^{2-3x}$$

$$\therefore nh = b - a = 1$$

**Step II.** Putting  $x = a$ ,  $a + h$ ,  $a + 2h$ ,  $a + (n - 1)h$  in  $f(x)$ , we have

$$f(a) = f(0) = e^2$$

$$f(a + h) = f(h) = e^{2-3h}$$

$$f(a + 2h) = f(2h) = e^{2-6h}$$

$$f(a + (n - 1)h) = f((n - 1)h) = e^{2-3(n-1)h}$$

**Step III.** Putting these values in

$$\int_a^b f(x) = \lim_{\substack{n \rightarrow \infty \\ h \rightarrow 0}} h[f(a) + f(a+h) + f(a+2h) + \dots + f(a+(n-1)h)],$$

$$\begin{aligned} \text{we have } \int_0^1 e^{2-3x} dx &= \lim_{\substack{n \rightarrow \infty \\ h \rightarrow 0}} h[e^2 + e^{2-3h} + e^{2-6h} + \dots + e^{2-3(n-1)h}] \\ &= \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h \cdot e^2 [1 + e^{-3h} + e^{-6h} + \dots + e^{-3(n-1)h}] \end{aligned}$$

**The series within brackets is a G.P. series of  $n$  terms**

$$\begin{aligned} \text{with } a = 1, r = e^{-3h} \text{ and using } S_n \text{ of G.P.} &= a \frac{(r^n - 1)}{r - 1} \\ &= e^2 \lim_{\substack{h \rightarrow 0 \\ n \rightarrow \infty}} h \cdot \left[ \frac{e^{-3nh} - 1}{e^{-3h} - 1} \right] \quad \left[ \because (e^{-3h})^n = e^{-3nh} \right] \end{aligned}$$

**Step IV.** Putting  $nh = 1$

$$= e^2 \lim_{h \rightarrow 0} h \cdot \left[ \frac{e^{-3} - 1}{e^{-3h} - 1} \right]$$

**Step V.** Taking limits as  $h \rightarrow 0$ ,

$$\begin{aligned} &= e^2 (e^{-3} - 1) \lim_{h \rightarrow 0} \frac{-3h}{e^{-3h} - 1} \times \left( -\frac{1}{3} \right) \\ &= (e^{-1} - e^2) \times 1 \times \left( -\frac{1}{3} \right) \quad \left[ \because \lim_{x \rightarrow 0} \frac{x}{e^x - 1} = 1 \right] \\ &= \frac{1}{3} \left( e^2 - \frac{1}{e} \right) \end{aligned}$$

**41. Choose the correct answer:**  $\int \frac{dx}{e^x + e^{-x}}$  is equal to

(A)  $\tan^{-1}(e^x) + c$

(B)  $\tan^{-1}(e^{-x}) + c$

(C)  $\log(e^x - e^{-x}) + c$

(D)  $\log(e^x + e^{-x}) + c$

**Sol.** Let  $I = \int \frac{dx}{e^x + e^{-x}} = \int \frac{1}{e^x + \left(\frac{1}{e^x}\right)} dx$

$$= \int \frac{1}{\left(\frac{e^{2x} + 1}{e^x}\right)} dx = \int \frac{e^x}{e^{2x} + 1} dx \quad \dots(i)$$

$$[\because e^x \cdot e^x = e^{x+x} = e^{2x}]$$

Put  $e^x = t$ .

$\left[ \because \text{For } \int f(e^x) dx, \text{ put } e^x = t \right]$

Therefore  $e^x = \frac{dt}{dx}$ . Therefore  $e^x dx = dt$

$$\begin{aligned} \therefore \text{From (i), } I &= \int \frac{dt}{t^2 + 1} = \tan^{-1} t + c \\ &= \tan^{-1} (e^x) + c \end{aligned}$$

$\therefore$  Option (A) is the correct answer.

42. Choose the correct answer:

$\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx$  is equal to

(A)  $\frac{-1}{\sin x + \cos x} + c$  (B)  $\log |\sin x + \cos x| + c$

(C)  $\log |\sin x - \cos x| + c$  (D)  $\frac{1}{(\sin x + \cos x)^2} + c$

Sol. Let  $I = \int \frac{\cos 2x}{(\sin x + \cos x)^2} dx = \int \frac{\cos^2 x - \sin^2 x}{(\sin x + \cos x)^2} dx$

$$= \int \frac{(\cos x + \sin x)(\cos x - \sin x)}{(\sin x + \cos x)(\sin x + \cos x)} dx = \int \frac{\cos x - \sin x}{\sin x + \cos x} dx$$
$$= \log |\sin x + \cos x| + c, \left[ \because \int \frac{f'(x)}{f(x)} dx = \log |f(x)| + c \right]$$

OR

Put denominator  $\sin x + \cos x = t$ .

$\therefore$  Option (B) is the correct answer.

43. Choose the correct answer:

If  $f(a + b - x) = f(x)$ , then  $\int_a^b x f(x) dx$  is equal to

(A)  $\frac{a+b}{2} \int_a^b f(b-x) dx$  (B)  $\frac{a+b}{2} \int_a^b f(b+x) dx$

(C)  $\frac{b-a}{2} \int_a^b f(x) dx$  (D)  $\frac{a+b}{2} \int_a^b f(x) dx$ .

Sol. Given:  $f(a + b - x) = f(x)$  ... (i)

Let  $I = \int_a^b x f(x) dx$  ... (ii)

Changing  $x$  to  $(a + b - x)$  in the Integrand on Right side (ii).

$$I = \int_a^b (a + b - x) f(a + b - x) dx \quad \dots (iii)$$

$$\left[ \because \text{By Property of definite integrals, } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx \right]$$

Putting  $f(a + b - x) = f(x)$  from (i) in integrand of (iii),

$$I = \int_a^b f(a + b - x) f(x) dx \quad \dots(iv)$$

Adding (ii) and (iv), we have  $2I = \int_a^b [x f(x) + (a + b - x) f(x)] dx$

$$2I = \int_a^b (x + a + b - x) f(x) dx = \int_a^b (a + b) f(x) dx = (a + b) \int_a^b f(x) dx$$

$$\text{Dividing by 2, } I = \left( \frac{a + b}{2} \right) \int_a^b f(x) dx$$

$$\text{or } \int_a^b x f(x) dx = \left( \frac{a + b}{2} \right) \int_a^b f(x) dx$$

$\therefore$  Option (D) is the correct answer.

**44. The value of  $\int_0^1 \tan^{-1} \left( \frac{2x-1}{1+x-x^2} \right) dx$  is**

(A) 1

(B) 0

(C) -1

(D)  $\frac{\pi}{4}$

$$\begin{aligned} \text{Sol. Let } I &= \int_0^1 \tan^{-1} \left( \frac{2x-1}{1+x-x^2} \right) dx = \int_0^1 \tan^{-1} \left( \frac{x+x-1}{1-x^2+x} \right) dx \\ &= \int_0^1 \tan^{-1} \left( \frac{x+(x-1)}{1-x(x-1)} \right) dx = \int_0^1 (\tan^{-1} x + \tan^{-1} (x-1)) dx \\ &\quad \left[ \because \tan^{-1} \frac{x+y}{1-xy} = \tan^{-1} x + \tan^{-1} y \right] \\ &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1} (x-1) \, dx \end{aligned}$$

Changing  $x$  to  $(1-x)$  in integrand of second integral

$$\begin{aligned} &\quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \\ &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1} (1-x-1) \, dx \\ &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1} (-x) \, dx = \int_0^1 \tan^{-1} x \, dx - \int_0^1 \tan^{-1} x \, dx \\ &\quad \left[ \because \tan^{-1} (-x) = -\tan^{-1} x \right] \\ &= 0. \end{aligned}$$

$\therefore$  Option (B) is the correct answer.