


RD Sharma  
Solutions  
Class 12 Maths  
Chapter  
Ex 18.5





### Maxima and Minima 18.5 Q1

Let  $x$  and  $y$  be the two numbers.

$$\text{Given that } x + y = 16 \quad \text{---(i)}$$

$$\text{Let } s = x^2 + y^2 \quad \text{---(ii)}$$

From (i) and (ii)

$$s = x^2 + (15 - x)^2$$

$$\begin{aligned}\therefore \frac{ds}{dx} &= 2x + 2(15 - x)(-1) \\ &= 2x - 30 + 2x \\ &= 4x - 30\end{aligned}$$

$$\text{Now, } \frac{ds}{dx} = 0$$

$$\Rightarrow 4x - 30 = 0$$

$$\Rightarrow x = \frac{15}{2}$$

Since,

$$\frac{d^2s}{dx^2} = 4 > 0$$

$$\therefore x = \frac{15}{2} \text{ is the point of local minima.}$$

So, from (i)

$$y = 15 - \frac{15}{2} = \frac{15}{2}$$

Hence, the required numbers are  $\frac{15}{2}, \frac{15}{2}$ .

## Maxima and Minima 18.5 Q2

Let  $x$  and  $y$  be the two parts of 64.

$$\therefore x + y = 64 \quad \text{---(i)}$$

$$\text{Let } S = x^3 + y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x^3 + (64 - x)^3$$

$$\begin{aligned} \therefore \frac{dS}{dx} &= 3x^2 + 3(64 - x)^2 \times (-1) \\ &= 3x^2 - 3(4096 - 128x + x^2) \\ &= -3(4096 - 128x) \end{aligned}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow -3(4096 - 128x) = 0$$

$$\Rightarrow x = 32$$

Now,

$$\frac{d^2S}{dx^2} = 384 > 0$$

$\therefore x = 32$  is the point of local minima.

Thus, the two parts of 64 are  $(32, 32)$ .

### Maxima and Minima 18.5 Q3

Let  $x$  and  $y$  be the two numbers, such that,  $x, y \geq -2$  and

$$x + y = \frac{1}{2} \quad \text{---(i)}$$

$$\text{Let } S = x + y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x + \left(\frac{1}{2} - x\right)^3$$

$$\begin{aligned}\therefore \frac{dS}{dx} &= 1 + 3\left(\frac{1}{2} - x\right)^2 \times (-1) \\ &= 1 - 3\left(\frac{1}{4} - x + x^2\right) \\ &= \frac{1}{4} + 3x - 3x^2\end{aligned}$$

For maximum and minimum,

$$\begin{aligned}\frac{dS}{dx} &= 0 \\ \Rightarrow \frac{1}{4} + 3x - 3x^2 &= 0 \\ \Rightarrow 1 + 12x - 12x^2 &= 0 \\ \Rightarrow 12x^2 - 12x - 1 &= 0 \\ \Rightarrow x &= \frac{12 \pm \sqrt{144 + 48}}{24} \\ \Rightarrow x &= \frac{1}{2} \pm \frac{8\sqrt{3}}{24} \\ \Rightarrow x &= \frac{1}{2} \pm \frac{1}{\sqrt{3}} \\ \Rightarrow x &= \frac{1}{2} - \frac{1}{\sqrt{3}}, \frac{1}{2} + \frac{1}{\sqrt{3}}\end{aligned}$$

Now,

$$\frac{d^2S}{dx^2} = 3 - 6x$$

$$\begin{aligned}\text{At } x &= \frac{1}{2} - \frac{1}{\sqrt{3}}, \quad \frac{d^2S}{dx^2} = 3 \left(1 - 2\left(\frac{1}{2} - \frac{1}{\sqrt{3}}\right)\right) \\ &= 3 \left(+\frac{2}{\sqrt{3}}\right) = 2\sqrt{3} > 0\end{aligned}$$

$$\therefore x = \frac{1}{2} - \frac{1}{\sqrt{3}} \text{ is point of local minima}$$

$\therefore$  from (i)

$$y = \frac{1}{2} - \left(\frac{1}{2} - \frac{1}{\sqrt{3}}\right) = \frac{1}{\sqrt{3}}$$

Hence, the required numbers are  $\frac{1}{2} - \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$ .

### Maxima and Minima 18.5 Q4

Let  $x$  and  $y$  be the two parts of 15, such that

$$\therefore x + y = 15 \quad \text{---(i)}$$

$$\text{Also, } S = x^2 y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x^2 (15 - x)^3$$

$$\begin{aligned}\therefore \frac{dS}{dx} &= 2x(15 - x)^3 - 3x^2(15 - x)^2 \\ &= (15 - x)^2 [30x - 2x^2 - 3x^2] \\ &= 5x(15 - x)^2(6 - x)\end{aligned}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 5x(15 - x)^2(6 - x) = 0$$

$$\Rightarrow x = 0, 15, 6$$

Now,

$$\frac{d^2S}{dx^2} = 5(15 - x)^2(6 - x) - 5x \times 2(15 - x)(6 - x) - 5x(15 - x)^2$$

$$\therefore \text{At } x = 0, \frac{d^2S}{dx^2} = 1125 > 0$$

$$\therefore x = 0 \text{ is point of local minima}$$

$$\text{At } x = 15, \frac{d^2S}{dx^2} = 0$$

$$\therefore x = 15 \text{ is an inflection point.}$$

$$\text{At } x = 6, \frac{d^2S}{dx^2} = -2430 < 0$$

$$\therefore x = 6 \text{ is the point of local maxima}$$

Thus the numbers are 6 and 9.

### Maxima and Minima 18.5 Q5

Let  $r$  and  $h$  be the radius and height of the cylinder respectively.

Then, volume ( $V$ ) of the cylinder is given by,

$$V = \pi r^2 h = 100 \quad (\text{given})$$

$$\therefore h = \frac{100}{\pi r^2}$$

Surface area ( $S$ ) of the cylinder is given by,

$$S = 2\pi r^2 + 2\pi r h = 2\pi r^2 + \frac{200}{r}$$

$$\therefore \frac{dS}{dr} = 4\pi r - \frac{200}{r^2}, \quad \frac{d^2S}{dr^2} = 4\pi + \frac{400}{r^3}$$

$$\frac{dS}{dr} = 0 \Rightarrow 4\pi r = \frac{200}{r^2}$$

$$\Rightarrow r^3 = \frac{200}{4\pi} = \frac{50}{\pi}$$

$$\Rightarrow r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$$

Now, it is observed that when  $r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ ,  $\frac{d^2S}{dr^2} > 0$ .

$\therefore$  By second derivative test, the surface area is the minimum when the radius of the cylinder

is  $\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$  cm.

$$\text{When } r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}, \quad h = \frac{100}{\pi \left(\frac{50}{\pi}\right)^{\frac{2}{3}}} = \frac{2 \times 50}{(50)^{\frac{2}{3}} (\pi)^{1-\frac{2}{3}}} = 2 \left(\frac{50}{\pi}\right)^{\frac{1}{3}} \text{ cm.}$$

Hence, the required dimensions of the can which has the minimum surface area is given by

radius =  $\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$  cm and height =  $2 \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$  cm.

We are given that the bending moment  $M$  at a distance  $x$  from one end of the beam is given by

$$(i) \quad M = \frac{WL}{2}x - \frac{W}{2}x^2$$

$$\therefore \quad \frac{dM}{dx} = \frac{WL}{2} - Wx$$

For maxima and minima,

$$\frac{dM}{dx} = 0 \Rightarrow \quad \frac{WL}{2} - Wx = 0 \Rightarrow \quad x = \frac{L}{2}$$

Now,

$$\frac{d^2M}{dx^2} = -W < 0$$

$$\therefore \quad x = \frac{L}{2} \text{ is point of local maxima.}$$

$$(ii) \quad M = \frac{Wx}{3} - \frac{Wx^3}{3L^2}$$

$$\therefore \quad \frac{dM}{dx} = \frac{W}{3} - \frac{Wx^2}{L^2}$$

For maxima and minima,

$$\frac{dM}{dx} = 0 \Rightarrow \quad \frac{W}{3} - \frac{Wx^2}{L^2} = 0 \Rightarrow \quad x = \frac{L}{\sqrt{3}}$$

Now,

$$\frac{d^2M}{dx^2} = -\frac{2xW}{L^2}$$

$$\text{At } x = \frac{L}{\sqrt{3}}, \quad \frac{d^2M}{dx^2} = -\frac{2W}{\sqrt{3}L} < 0$$

$$\therefore \quad x = \frac{L}{\sqrt{3}} \text{ is point of local maxima}$$

$$\Rightarrow \quad \begin{aligned} \frac{d^2s}{dx^2} &= -\frac{\sqrt{2}r}{\frac{r^2}{2}} \\ &= \frac{2\sqrt{2}}{r} < 0 \end{aligned}$$

$$\therefore \quad x = \frac{r}{\sqrt{2}} \text{ is the point of local maxima}$$

From (i)

$$y = \frac{r}{\sqrt{2}}$$

Hence,  $x = \frac{r}{\sqrt{2}}, y = \frac{r}{\sqrt{2}}$  is the required number.



Let a piece of length  $l$  be cut from the given wire to make a square.

Then, the other piece of wire to be made into a circle is of length  $(28 - l)$  m.

Now, side of square  $= \frac{l}{4}$ .

Let  $r$  be the radius of the circle. Then,  $2\pi r = 28 - l \Rightarrow r = \frac{1}{2\pi}(28 - l)$ .

The combined areas of the square and the circle ( $A$ ) is given by,

$$\begin{aligned} A &= (\text{side of the square})^2 + r^2 \\ &= \frac{l^2}{16} + \pi \left[ \frac{1}{2\pi}(28 - l) \right]^2 \\ &= \frac{l^2}{16} + \frac{1}{4\pi}(28 - l)^2 \\ \therefore \frac{dA}{dl} &= \frac{2l}{16} + \frac{2}{4\pi}(28 - l)(-1) = \frac{l}{8} - \frac{1}{2\pi}(28 - l) \\ \frac{d^2A}{dl^2} &= \frac{1}{8} + \frac{1}{2\pi} > 0 \\ \text{Now, } \frac{dA}{dl} = 0 &\Rightarrow \frac{l}{8} - \frac{1}{2\pi}(28 - l) = 0 \\ \Rightarrow \frac{\pi l - 4(28 - l)}{8\pi} &= 0 \\ \Rightarrow (\pi + 4)l - 112 &= 0 \\ \Rightarrow l &= \frac{112}{\pi + 4} \end{aligned}$$

Thus, when  $l = \frac{112}{\pi + 4}$ ,  $\frac{d^2A}{dl^2} > 0$ .

$\therefore$  By second derivative test, the area ( $A$ ) is the minimum when  $l = \frac{112}{\pi + 4}$ .

Hence, the combined area is the minimum when the length of the wire in making the square is  $\frac{112}{\pi + 4}$  cm while the length of the wire in making the circle is  $28 - \frac{112}{\pi + 4} = \frac{28\pi}{\pi + 4}$  cm.

Let the wire of length 20 m be cut into  $x$  cm and  $y$  cm and bent into a square and equilateral triangle, so that the sum of area of square and triangle is minimum.

Now,

$$\begin{aligned}x + y &= 20 \\x &= 4l \text{ and } y = 3a\end{aligned}\quad \text{---(i)}$$

Let  $s$  = sum of area of square and triangle

$$s = l^2 + \frac{\sqrt{3}}{4}a^2 \quad \text{---(ii)}$$

$$\left[ \because \text{area of equilateral } \Delta = \frac{\sqrt{3}}{4}(\text{one side})^2 \right]$$

We have,  $4l + 3a = 20$

$$\Rightarrow 4l = 20 - 3a$$

$$\Rightarrow l = \frac{20 - 3a}{4}$$

From (i), we have,

$$s = \left( \frac{20 - 3a}{4} \right)^2 + \frac{\sqrt{3}}{4}a^2$$

$$\frac{ds}{da} = 2 \left( \frac{20 - 3a}{4} \right) \left( \frac{-3}{4} \right) + 2a \times \frac{\sqrt{3}}{4}$$

To find the maximum or minimum,  $\frac{ds}{da} = 0$

$$\Rightarrow 2 \left( \frac{20 - 3a}{4} \right) \left( \frac{-3}{4} \right) + 2a \times \frac{\sqrt{3}}{4} = 0$$

$$\Rightarrow -3(20 - 3a) + 4a\sqrt{3} = 0$$

$$\Rightarrow -60 + 9a + 4a\sqrt{3} = 0$$

$$\Rightarrow 9a + 4a\sqrt{3} = 60$$

$$\Rightarrow a(9 + 4\sqrt{3}) = 60$$

$$\Rightarrow a = \frac{60}{9 + 4\sqrt{3}}$$

Differentiating once again, we have,

$$\frac{d^2s}{da^2} = \frac{9 + 4\sqrt{3}}{8} > 0$$

Thus, the sum of the areas of the square and triangle is minimum when  $a = \frac{60}{9 + 4\sqrt{3}}$

We know that,  $l = \frac{20 - 3a}{4}$

$$\Rightarrow l = \frac{20 - 3 \left( \frac{60}{9 + 4\sqrt{3}} \right)}{4}$$

$$\Rightarrow l = \frac{180 + 80\sqrt{3} - 180}{4(9 + 4\sqrt{3})}$$

$$\Rightarrow l = \frac{20\sqrt{3}}{9 + 4\sqrt{3}}$$

Let  $r$  be the radius of the circle and  $a$  be the side of the square.

Then, we have:

$$2\pi r + 4a = k \text{ (where } k \text{ is constant)}$$

$$\Rightarrow a = \frac{k - 2\pi r}{4}$$

The sum of the areas of the circle and the square ( $A$ ) is given by,

$$A = \pi r^2 + a^2 = \pi r^2 + \frac{(k - 2\pi r)^2}{16}$$

$$\therefore \frac{dA}{dr} = 2\pi r + \frac{2(k - 2\pi r)(-2\pi)}{16} = 2\pi r - \frac{\pi(k - 2\pi r)}{4}$$

$$\text{Now, } \frac{dA}{dr} = 0$$

$$\Rightarrow 2\pi r = \frac{\pi(k - 2\pi r)}{4}$$

$$8r = k - 2\pi r$$

$$\Rightarrow (8 + 2\pi)r = k$$

$$\Rightarrow r = \frac{k}{8 + 2\pi} = \frac{k}{2(4 + \pi)}$$

$$\text{Now, } \frac{d^2A}{dr^2} = 2\pi + \frac{\pi^2}{2} > 0$$

$$\therefore \text{ When } r = \frac{k}{2(4 + \pi)}, \frac{d^2A}{dr^2} > 0.$$

$$\therefore \text{ The sum of the areas is least when } r = \frac{k}{2(4 + \pi)}.$$

$$\text{When } r = \frac{k}{2(4 + \pi)}, a = \frac{k - 2\pi \left[ \frac{k}{2(4 + \pi)} \right]}{4} = \frac{k(4 + \pi) - k}{4(4 + \pi)} = \frac{4k}{4(4 + \pi)} = \frac{k}{4 + \pi} = 2r.$$

Hence, it has been proved that the sum of their areas is least when the side of the square is double the radius of the circle.

$ABC$  is a right angled triangle. Hypotenuse  $h = AC = 5$  cm.

Let  $x$  and  $y$  one the other two side of the triangle.

$$\therefore x^2 + y^2 = 25 \quad \text{---(i)}$$

$$\therefore \text{Area of } \triangle ABC = \frac{1}{2} BC \times AB$$

$$\Rightarrow S = \frac{1}{2} xy \quad \text{---(ii)}$$

From (i) and (ii)

$$S = \frac{1}{2} x \sqrt{25 - x^2}$$

$$\begin{aligned} \therefore \frac{ds}{dx} &= \frac{1}{2} \left[ \sqrt{25 - x^2} - \frac{2x^2}{2\sqrt{25 - x^2}} \right] \\ &= \frac{1}{2} \frac{[25 - x^2 - x^2]}{\sqrt{25 - x^2}} \\ &= \frac{1}{2} \left[ \frac{25 - 2x^2}{\sqrt{25 - x^2}} \right] \end{aligned}$$

For maxima and minima,

$$\frac{ds}{dx} = 0$$

$$\Rightarrow \frac{1}{2} \left[ \frac{25 - 2x^2}{\sqrt{25 - x^2}} \right] = 0$$

$$\Rightarrow x = 5\sqrt{2}$$

Now,

$$\frac{d^2s}{dx^2} = \frac{1}{2} \frac{\sqrt{25 - x^2} \times (-4x) + \frac{(25 - 2x^2) 2x}{2\sqrt{25 - x^2}}}{(25 - x^2)}$$

$$\text{At } x = \frac{5}{\sqrt{2}}, \frac{d^2s}{dx^2} = \frac{1}{2} \left[ \frac{-\frac{25}{\sqrt{2}} \times \frac{5}{\sqrt{2}} + 0}{\frac{25}{2}} \right]$$

$$= -\frac{5}{2} < 0$$

$$\therefore x = \frac{5}{\sqrt{2}} \text{ is a point local maxima,}$$

$ABC$  is a given triangle with  $AB = a, BC = b$  and  $\angle ABC = \theta$ .

$AD$  is perpendicular to  $BC$ .

$$\therefore BD = a \sin \theta$$

Now,

$$\text{Area of } \triangle ABC = \frac{1}{2} \times BC \times AD$$

$$\Rightarrow A = \frac{1}{2} b \times a \sin \theta \quad \text{---(i)}$$

$$\therefore \frac{dA}{d\theta} = \frac{1}{2} ab \cos \theta$$

For maxima and minima,

$$\frac{dA}{d\theta} = 0$$

$$\Rightarrow \frac{1}{2} ab \cos \theta = 0$$

$$\Rightarrow \cos \theta = 0$$

$$\Rightarrow \theta = \frac{\pi}{2}$$

Now,

$$\frac{d^2A}{d\theta^2} = -\frac{1}{2} ab \sin \theta$$

$$\text{At } \theta = \frac{\pi}{2}, \quad \frac{d^2A}{d\theta^2} = -\frac{1}{2} ab < 0$$

$$\therefore \theta = \frac{\pi}{2} \text{ is point of local maxima}$$

$$\therefore \text{Maximum area of } \Delta = \frac{1}{2} ab \sin \frac{\pi}{2} = \frac{1}{2} ab.$$

Let the side of the square to be cut off be  $x$  cm. Then, the length and the breadth of the box will be  $(18 - 2x)$  cm each and the height of the box is  $x$  cm.

Therefore, the volume  $V(x)$  of the box is given by,

$$V(x) = x(18 - 2x)^2$$

$$\begin{aligned}\therefore V'(x) &= (18 - 2x)^2 - 4x(18 - 2x) \\ &= (18 - 2x)[18 - 2x - 4x] \\ &= (18 - 2x)(18 - 6x) \\ &= 6 \times 2(9 - x)(3 - x) \\ &= 12(9 - x)(3 - x)\end{aligned}$$

$$\begin{aligned}\text{And, } V''(x) &= 12[-(9 - x) - (3 - x)] \\ &= -12(9 - x + 3 - x) \\ &= -12(12 - 2x) \\ &= -24(6 - x)\end{aligned}$$

Maximum volume is  $V_{x=3} = 3 \times (18 - 2 \times 3)^2$

$$\Rightarrow V = 3 \times 12^2$$

$$\Rightarrow V = 3 \times 144$$

$$\Rightarrow V = 432 \text{ cm}^3$$

### Maxima and Minima 18.5 Q13

Let the side of the square to be cut off be  $x$  cm. Then, the height of the box is  $x$ , the length is  $45 - 2x$ , and the breadth is  $24 - 2x$ .

Therefore, the volume  $V(x)$  of the box is given by,

$$\begin{aligned}V(x) &= x(45 - 2x)(24 - 2x) \\ &= x(1080 - 90x - 48x + 4x^2) \\ &= 4x^3 - 138x^2 + 1080x \\ \therefore V'(x) &= 12x^2 - 276x + 1080 \\ &= 12(x^2 - 23x + 90) \\ &= 12(x - 18)(x - 5) \\ V''(x) &= 24x - 276 = 12(2x - 23)\end{aligned}$$

$$\text{Now, } V'(x) = 0 \Rightarrow x = 18 \text{ and } x = 5$$

It is not possible to cut off a square of side 18 cm from each corner of the rectangular sheet.

Thus,  $x$  cannot be equal to 18.

$$\therefore x = 5$$

$$\text{Now, } V''(5) = 12(10 - 23) = 12(-13) = -156 < 0$$

$\therefore$  By second derivative test,  $x = 5$  is the point of maxima.

Hence, the side of the square to be cut off to make the volume of the box maximum possible is 5 cm.

### Maxima and Minima 18.5 Q14

Let  $l$ ,  $b$ , and  $h$  represent the length, breadth, and height of the tank respectively.

Then, we have height  $(h) = 2$  m

$$\text{Volume of the tank} = 8\text{m}^3$$

$$\text{Volume of the tank} = l \times b \times h$$

$$\therefore 8 = l \times b \times 2$$

$$\Rightarrow lb = 4 \Rightarrow b = \frac{4}{l}$$

$$\text{Now, area of the base} = lb = 4$$

$$\text{Area of the 4 walls } (A) = 2h(l + b)$$

$$\therefore A = 4\left(l + \frac{4}{l}\right)$$

$$\Rightarrow \frac{dA}{dl} = 4\left(1 - \frac{4}{l^2}\right)$$

$$\text{Now, } \frac{dA}{dl} = 0$$

$$\Rightarrow 1 - \frac{4}{l^2} = 0$$

$$\Rightarrow l^2 = 4$$

$$\Rightarrow l = \pm 2$$

However, the length cannot be negative.

Therefore, we have  $l = 4$ .

$$\therefore b = \frac{4}{l} = \frac{4}{2} = 2$$

$$\text{Now, } \frac{d^2 A}{dl^2} = \frac{32}{l^3}$$

$$\text{When } l = 2, \frac{d^2 A}{dl^2} = \frac{32}{8} = 4 > 0.$$

Thus, by second derivative test, the area is the minimum when  $l = 2$ .

We have  $l = b = h = 2$ .

$$\therefore \text{Cost of building the base} = \text{Rs } 70 \times (lb) = \text{Rs } 70 (4) = \text{Rs } 280$$

$$\text{Cost of building the walls} = \text{Rs } 2h(l + b) \times 45 = \text{Rs } 90 (2) (2 + 2)$$


$$= \text{Rs } 8 (90) = \text{Rs } 720$$

$$\text{Required total cost} = \text{Rs } (280 + 720) = \text{Rs } 1000$$

Hence, the total cost of the tank will be Rs 1000.



Radius of the semicircular opening =  $\frac{x}{2}$



It is given that the perimeter of the window is 10 m.

$$\therefore x + 2y + \frac{\pi x}{2} = 10$$

$$\Rightarrow x \left( 1 + \frac{\pi}{2} \right) + 2y = 10$$

$$\Rightarrow 2y = 10 - x \left( 1 + \frac{\pi}{2} \right)$$

$$\Rightarrow y = 5 - x \left( \frac{1}{2} + \frac{\pi}{4} \right)$$

$\therefore$  Area of the window ( $A$ ) is given by,

$$A = xy + \frac{\pi}{2} \left( \frac{x}{2} \right)^2$$

$$= x \left[ 5 - x \left( \frac{1}{2} + \frac{\pi}{4} \right) \right] + \frac{\pi}{8} x^2$$

$$= 5x - x^2 \left( \frac{1}{2} + \frac{\pi}{4} \right) + \frac{\pi}{8} x^2$$

$$\therefore \frac{dA}{dx} = 5 - 2x \left( \frac{1}{2} + \frac{\pi}{4} \right) + \frac{\pi}{4} x$$

$$= 5 - x \left( 1 + \frac{\pi}{2} \right) + \frac{\pi}{4} x$$

$$\therefore \frac{d^2 A}{dx^2} = - \left( 1 + \frac{\pi}{2} \right) + \frac{\pi}{4} = -1 - \frac{\pi}{4}$$

$$\text{Now, } \frac{dA}{dx} = 0$$

$$\Rightarrow 5 - x \left( 1 + \frac{\pi}{2} \right) + \frac{\pi}{4} x = 0$$

$$\Rightarrow 5 - x - \frac{\pi}{4} x = 0$$

$$\Rightarrow x \left( 1 + \frac{\pi}{4} \right) = 5$$

$$\Rightarrow x = \frac{5}{\left( 1 + \frac{\pi}{4} \right)} = \frac{20}{\pi + 4}$$

Thus, when  $x = \frac{20}{\pi + 4}$  then  $\frac{d^2 A}{dx^2} < 0$ .


Therefore, by second derivative test, the area is the maximum when length  $x = \frac{20}{\pi + 4}$  m.

Now,

$$y = 5 - \frac{20}{\pi + 4} \left( \frac{2 + \pi}{4} \right) = 5 - \frac{5(2 + \pi)}{\pi + 4} = \frac{10}{\pi + 4} \text{ m}$$

Hence, the required dimensions of the window to admit maximum light is given

by length  $= \frac{20}{\pi + 4}$  m and breadth  $= \frac{10}{\pi + 4}$  m.



The perimeter of the window = 12 m

$$\Rightarrow (l + 2b) + (l + l) = 12$$

$$\Rightarrow 3l + 2b = 12 \quad \text{----- (i)}$$

Let  $S$  = Area of the rectangle + Area of the equilateral  $\Delta$

From (i),

$$S = l \left( \frac{12 - 3l}{2} \right) + \frac{\sqrt{3}}{4} l^2$$

$$\therefore \frac{dS}{dl} = 6 - 3l + \frac{\sqrt{3}}{2} l = 6 - \sqrt{3} \left( \sqrt{3} - \frac{1}{2} \right) l$$

For maxima and minima,

$$\frac{dS}{dl} = 0$$

$$\Rightarrow 6 - \sqrt{3} \left( \sqrt{3} - \frac{1}{2} \right) l = 0$$

$$\Rightarrow l = \frac{6}{\sqrt{3} \left( \sqrt{3} - \frac{1}{2} \right)} = \frac{12}{6 - \sqrt{3}}$$

$$\text{Now, } \frac{d^2S}{dl^2} = -\sqrt{3} \left( \sqrt{3} - \frac{1}{2} \right) = -3 + \frac{\sqrt{3}}{2} < 0$$


$$\therefore l = \frac{12}{6 - \sqrt{3}} \text{ is the point of local maxima}$$

From (i),

$$b = \frac{12 - 3l}{2} = \frac{12 - 3 \left( \frac{12}{6 - \sqrt{3}} \right)}{2} = \frac{24 - 6\sqrt{3}}{6 - \sqrt{3}}$$

A sphere of fixed radius ( $R$ ) is given.

Let  $r$  and  $h$  be the radius and the height of the cylinder respectively.



From the given figure, we have  $h = 2\sqrt{R^2 - r^2}$ .

The volume ( $V$ ) of the cylinder is given by,

$$\begin{aligned} V &= \pi r^2 h = 2\pi r^2 \sqrt{R^2 - r^2} \\ \therefore \frac{dV}{dr} &= 4\pi r \sqrt{R^2 - r^2} + \frac{2\pi r^2 (-2r)}{2\sqrt{R^2 - r^2}} \\ &= 4\pi r \sqrt{R^2 - r^2} - \frac{2\pi r^3}{\sqrt{R^2 - r^2}} \\ &= \frac{4\pi r (R^2 - r^2) - 2\pi r^3}{\sqrt{R^2 - r^2}} \\ &= \frac{4\pi r R^2 - 6\pi r^3}{\sqrt{R^2 - r^2}} \end{aligned}$$

$$\text{Now, } \frac{dV}{dr} = 0 \Rightarrow 4\pi r R^2 - 6\pi r^3 = 0$$

$$\Rightarrow r^2 = \frac{2R^2}{3}$$

$$\begin{aligned}\text{Now, } \frac{d^2V}{dr^2} &= \frac{\sqrt{R^2 - r^2} (4\pi R^2 - 18\pi r^2) - (4\pi r R^2 - 6\pi r^3) \frac{(-2r)}{2\sqrt{R^2 - r^2}}}{(R^2 - r^2)} \\ &= \frac{(R^2 - r^2)(4\pi R^2 - 18\pi r^2) + r(4\pi r R^2 - 6\pi r^3)}{(R^2 - r^2)^{\frac{3}{2}}} \\ &= \frac{4\pi R^4 - 22\pi r^2 R^2 + 12\pi r^4 + 4\pi r^2 R^2}{(R^2 - r^2)^{\frac{3}{2}}}\end{aligned}$$

Now, it can be observed that at  $r^2 = \frac{2R^2}{3}$ ,  $\frac{d^2V}{dr^2} < 0$ .


$\therefore$  The volume is the maximum when  $r^2 = \frac{2R^2}{3}$ .

When  $r^2 = \frac{2R^2}{3}$ , the height of the cylinder is  $2\sqrt{R^2 - \frac{2R^2}{3}} = 2\sqrt{\frac{R^2}{3}} = \frac{2R}{\sqrt{3}}$ .

Hence, the volume of the cylinder is the maximum when the height of the cylinder is  $\frac{2R}{\sqrt{3}}$ .

### Maxima and Minima 18.5 Q18

Let  $EFGH$  be a rectangle inscribed in a semi-circle with radius  $r$ .



Let  $l$  and  $b$  are the length and width of rectangle.

In  $\triangle OHE$

$$HE^2 = OE^2 - OH^2$$

$$\Rightarrow HE = b = \sqrt{r^2 - \left(\frac{l}{2}\right)^2} \quad \text{---(i)}$$

Let  $S$  = Area of rectangle

$$= lb = l \times \sqrt{r^2 - \left(\frac{l}{2}\right)^2}$$

$$\therefore S = \frac{1}{2} l \sqrt{4r^2 - l^2}$$

$$\begin{aligned} \therefore \frac{ds}{dl} &= \frac{1}{2} \left[ \sqrt{4r^2 - l^2} - \frac{l^2}{\sqrt{4r^2 - l^2}} \right] \\ &= \frac{1}{2} \left[ \frac{4r^2 - l^2 - l^2}{\sqrt{4r^2 - l^2}} \right] \\ &= \frac{2r^2 - l^2}{\sqrt{4r^2 - l^2}} \end{aligned}$$

For maxima and minima,

$$\begin{aligned} \frac{ds}{dl} &= 0 \\ \Rightarrow \frac{2r^2 - l^2}{\sqrt{4r^2 - l^2}} &= 0 \\ \Rightarrow l &= \pm \sqrt{2}r \end{aligned}$$

Also,

$$\frac{d^2s}{dl^2} = 0 \text{ at } l = \sqrt{2}r$$

So, the dimension of the rectangle

$$l = \sqrt{2}r, \quad b = \sqrt{r^2 - \left(\frac{l}{2}\right)^2} = \frac{r}{\sqrt{2}}$$

$$\begin{aligned} \text{Area of rectangle} = lb &= \sqrt{2}r \times \frac{r}{\sqrt{2}} \\ &= r^2. \end{aligned}$$

Let  $r$  and  $h$  be the radius and the height (altitude) of the cone respectively.

Then, the volume ( $V$ ) of the cone is given as:

$$V = \frac{1}{3}\pi r^2 h \Rightarrow h = \frac{3V}{r^2}$$

The surface area ( $S$ ) of the cone is given by,

$S = \pi r l$  (where  $l$  is the slant height)

$$\begin{aligned} &= \pi r \sqrt{r^2 + h^2} \\ &= \pi r \sqrt{r^2 + \frac{9V^2}{r^4}} = \frac{\pi r \sqrt{r^6 + 9V^2}}{r^2} \\ &= \frac{1}{r} \sqrt{\pi^2 r^6 + 9V^2} \end{aligned}$$

$$\begin{aligned} \therefore \frac{dS}{dr} &= \frac{r \cdot \frac{6\pi^2 r^5}{2\sqrt{\pi^2 r^6 + 9V^2}} - \sqrt{\pi^2 r^6 + 9V^2}}{r^2} \\ &= \frac{3\pi^2 r^6 - \pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \\ &= \frac{2\pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \end{aligned}$$

$$\begin{aligned} &= \frac{2\pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \\ \text{Now, } \frac{dS}{dr} &= 0 \Rightarrow 2\pi^2 r^6 = 9V^2 \Rightarrow r^6 = \frac{9V^2}{2\pi^2} \end{aligned}$$

Thus, it can be easily verified that when  $r^6 = \frac{9V^2}{2\pi^2}$ ,  $\frac{d^2S}{dr^2} > 0$ .


$\therefore$  By second derivative test, the surface area of the cone is the least when  $r^6 = \frac{9V^2}{2\pi^2}$ .

$$\text{When } r^6 = \frac{9V^2}{2\pi^2}, h = \frac{3V}{r^2} = \frac{3}{r^2} \left( \frac{2\pi^2 r^6}{9} \right)^{\frac{1}{2}} = \frac{3}{r^2} \cdot \frac{\sqrt{2}\pi r^3}{3} = \sqrt{2}r.$$

Hence, for a given volume, the right circular cone of the least curved surface has an altitude equal to  $\sqrt{2}$  times the radius of the base.

We have a cone, which is inscribed in a sphere.

Let  $v$  be the volume of greatest cone  $ABC$ . It is obvious that, for maximum volume the axis of the cone must be along the diameter of sphere.



Let  $OD = x$  and  $AO = OB = R$

$$\Rightarrow BD = \sqrt{R^2 - x^2} \text{ and } AD = R + x$$

Now,

$$\begin{aligned} v &= \frac{1}{3} \pi r^2 h \\ &= \frac{1}{3} \pi BD^2 \times AD \\ &= \frac{1}{3} \pi (R^2 - x^2) \times (R + x) \end{aligned}$$

$$\begin{aligned} \therefore \frac{dv}{dx} &= \frac{\pi}{3} [-2x(R + x) + R^2 - x^2] \\ &= \frac{\pi}{3} [R^2 - 2xR - 3x^2] \end{aligned}$$

For maximum and minimum

$$\begin{aligned} \frac{dv}{dx} &= 0 \\ \Rightarrow \frac{\pi}{3} [R^2 - 2xR - 3x^2] &= 0 \\ \Rightarrow \frac{\pi}{3} [(R - 3x)(R + x)] &= 0 \\ \Rightarrow R - 3x = 0 \text{ or } x = -R \\ \Rightarrow x = \frac{R}{3} \end{aligned}$$

$\left[ \because x = -R \text{ is not possible as, } x = -R \text{ will make the altitude } 0 \right]$

Now,

$$\begin{aligned} \frac{d^2v}{dx^2} &= \frac{\pi}{3} [-2R - 6x] \\ \text{At } x = \frac{R}{3}, \frac{d^2v}{dx^2} &= \frac{\pi}{3} [-2R - 2R] \\ &= \frac{-4\pi R}{3} < 0 \end{aligned}$$

$\therefore x = \frac{R}{3}$  is the point of local maxima.



$$\text{Volume of the cone} = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow V = \frac{1}{3} \pi r^2 h$$

Squaring both the sides, we have,

$$\begin{aligned} V^2 &= \left( \frac{1}{3} \pi r^2 h \right)^2 \\ &= \frac{1}{9} \pi^2 r^4 h^2 \dots (1) \end{aligned}$$

$$\Rightarrow \pi^2 r^4 h^2 = \frac{9V^2}{r^2} \dots (2)$$

Consider the curved surface area of the cone.

Thus,

$$C = \pi r l$$

Squaring both the sides, we have,


$$C^2 = \pi^2 r^2 l^2$$


We know that  $l^2 = r^2 + h^2$

$$\Rightarrow C^2 = \pi^2 r^2 (r^2 + h^2)$$

$$\Rightarrow C^2 = \pi^2 r^4 + \pi^2 r^2 h^2$$

$$\Rightarrow C^2 = \pi^2 r^4 + \frac{9V^2}{r^2} \dots (\text{From equation (2)})$$





ABC is an isosceles triangle such that  $AB = AC$ .  
 The vertical angle  $\angle BAC = 2\theta$   
 Triangle is inscribed in the circle with center O and radius a.

Draw AM perpendicular to BC.  
 $\therefore \Delta ABC$  is an isosceles triangle the circumcentre of the circle will lie on the perpendicular from A to BC.

Let O be the circumcentre.  
 $\angle BOC = 2 \times 2\theta = 4\theta$  .....[Using central angle theorem]  
 $\angle COM = 2\theta$  .....[ $\therefore \Delta OMB$  and  $\Delta OMC$  are congruent triangles]  
 $OA = OB = OC = a$  .....[Radius of the circle]

In  $\Delta OMC$ ,  
 $CM = a \sin 2\theta$  and  $OM = a \cos 2\theta$   
 $BC = 2CM$ ...[Perpendicular from the center bisects the chord]  
 $BC = 2a \sin 2\theta$  ..... (1)  
 Height of  $\Delta ABC = AM = AO + OM$   
 $AM = a + a \cos 2\theta$  ..... (2)

Area of  $\Delta ABC$  is,

$$A = \frac{1}{2} \times BC \times AM$$

Differentiating equation (3) with respect to  $\theta$

$$\frac{dA}{d\theta} = a^2 \left( 2 \cos 2\theta + \frac{1}{2} \times 4 \cos 4\theta \right)$$

$$\frac{dA}{d\theta} = 2a^2 (\cos 2\theta + \cos 4\theta)$$

Differentiating again with respect to  $\theta$

$$\frac{d^2A}{d\theta^2} = 2a^2 (-2 \sin 2\theta - 4 \sin 4\theta)$$

For maximum value of area equating  $\frac{dA}{d\theta} = 0$

$$2a^2 (\cos 2\theta + \cos 4\theta) = 0$$

$$\cos 2\theta + \cos 4\theta = 0$$

$$\cos 2\theta + 2 \cos^2 2\theta - 1 = 0$$

$$(2 \cos 2\theta - 1)(2 \cos 2\theta + 1) = 0$$

$$\cos 2\theta = \frac{1}{2} \text{ or } \cos 2\theta = -1$$

$$2\theta = \frac{\pi}{3} \text{ or } 2\theta = \pi$$

$$\theta = \frac{\pi}{6} \text{ or } \theta = \frac{\pi}{2}$$

If  $2\theta = \pi$  it will not form a triangle.

$$\therefore \theta = \frac{\pi}{6}$$

Also  $\frac{d^2A}{d\theta^2}$  is negative for  $\theta = \frac{\pi}{6}$ .

**Maxima**

Thus the area of the triangle is maximum when  $\theta = \frac{\pi}{6}$ .

Here,  $ABCD$  is a rectangle with width  $AB = x$  cm and length  $AD = y$  cm.

The rectangle is rotated about  $AD$ . Let  $v$  be the volume of the cylinder so formed.

$$\therefore v = \pi r^2 y \quad \text{---(i)}$$

Again,

$$\text{Perimeter of } ABCD = 2(l + b) = 2(x + y) \quad \text{---(ii)}$$

$$\Rightarrow 36 = 2(x + y)$$

$$\Rightarrow y = 18 - x \quad \text{---(iii)}$$

From (i) and (ii), we get

$$v = \pi r^2 (18 - x) = \pi (18x^2 - x^3)$$

$$\Rightarrow \frac{dv}{dx} = \pi (36x - 3x^2)$$

For maxima or minima, we have,

$$\frac{dv}{dx} = 0$$

$$\Rightarrow \pi (36x - 3x^2) = 0$$

$$\Rightarrow 3\pi (12x - x^2) = 0$$

$$\Rightarrow x(12 - x) = 0$$

$$\Rightarrow x = 0 \text{ (Not possible) or } 12$$

$$\therefore x = 12 \text{ cm}$$

From (iii)

$$y = 18 - 12 = 6 \text{ cm}$$

Now,

$$\frac{d^2v}{dx^2} = \pi (36 - 6x)$$

$$\text{At } (x = 12, y = 6) \frac{d^2v}{dx^2} = \pi (36 - 72) = -36\pi < 0$$


$$\therefore (x = 12, y = 6) \text{ is the point of local maxima,}$$

Hence,

The dimension of rectangle, which wiout maximum value, when revolved about one of its side is width = 12 cm and length = 6 cm.

### Maxima and Minima 18.5 Q24

Let  $r$  and  $h$  be the radius of the base of cone and height of the cone respectively.



Let  $OD = x$

It is obvious that the axis of cone must be along the diameter of sphere for maximum volume of cone.

Now,

$$\begin{aligned}\text{In } \triangle BOD, BD &= \sqrt{R^2 - x^2} \\ &= \sqrt{144 - x^2}\end{aligned}$$

$$AD = AO + OD = R + x = 12 + x$$

$$v = \text{volume of cone} = \frac{1}{3} \pi r^2 h$$

$$\begin{aligned}\Rightarrow v &= \frac{1}{3} \pi BD^2 \times AD \\ &= \frac{1}{3} \pi (144 - x^2) (12 + x) \\ &= \frac{1}{3} \pi (1728 + 144x - 12x^2 - x^3)\end{aligned}$$

$$\therefore \frac{dv}{dx} = \frac{1}{3} \pi (144 - 24x - 3x^2)$$

For maximum and minimum of  $v$ ,

$$\frac{dv}{dx} = 0$$

$$\Rightarrow \frac{1}{3} \pi (144 - 24x - 3x^2) = 0$$

$$\Rightarrow x = -12, 4$$

$x = -12$  is not possible

$$\therefore x = 4$$

Now,

$$\frac{d^2v}{dx^2} = \frac{\pi}{3} (-24 - 6x)$$

$$\begin{aligned}\text{At } x = 4, \frac{d^2v}{dx^2} &= -2\pi (4 + x) \\ &= -2\pi \times 8 = -16\pi < 0\end{aligned}$$

$\therefore x = 4$  is point of local maxima.

Hence,

$$\begin{aligned}\text{Height of cone of maximum volume} &= R + x \\ &= 12 + 4 \\ &= 16 \text{ cm.}\end{aligned}$$

We have, a closed cylinder whose volume  $v = 2156 \text{ cm}^3$

Let  $r$  and  $h$  be the radius and the height of the cylinder. Then,

$$\therefore v = \pi r^2 h = 2156 \quad \text{---(i)}$$

$$\text{Total surface area} = S = 2\pi rh + 2\pi r^2$$

$$\Rightarrow S = 2\pi r(h + r) \quad \text{---(ii)}$$

From (i) and (ii)

$$S = \frac{2156 \times 2}{r} + 2\pi r^2$$

$$\therefore \frac{ds}{dr} = -\frac{4312}{r^2} + 4\pi r$$

For maximum and minimum

$$\frac{ds}{dr} = 0$$

$$\Rightarrow \frac{-4312 + 4\pi r^3}{r^2} = 0$$

$$\Rightarrow r^3 = \frac{4312}{4\pi}$$

$$\Rightarrow r = 7$$

Now,

$$\frac{d^2s}{dr^2} = \frac{8624}{r^3} + 4\pi > 0 \text{ for } r = 7.$$

$\therefore r = 7$  is the point of local minima


Hence,

The total surface area of closed cylinder will be minimum at  $r = 7 \text{ cm}$ .

### Maxima and Minima 18.5 Q26

Let  $r$  be the radius of the base of the cylinder and  $h$  be the height of the cylinder.

$$\therefore LM = h.$$



Let  $R = 5\sqrt{3}$  cm be the radius of the sphere.

It is obvious, that for maximum volume of cylinder  $ABCD$ , the axis of cylinder must be along the diameter of sphere.

Let  $OL = x$

$\therefore h = 2x$

Now,

$$\begin{aligned}\text{In } \triangle AOL, AL &= \sqrt{AO^2 - OL^2} \\ &= \sqrt{75 - x^2}\end{aligned}$$

Now,

$$v = \text{volume of cylinder} = \pi r^2 h$$

$$\begin{aligned}\Rightarrow v &= \pi AL^2 \times ML \\ &= \pi (75 - x^2) \times 2x\end{aligned}$$

For maxima and minima of  $v$ , we must have,

$$\begin{aligned}\frac{dv}{dx} &= \pi [150 - 6x^2] = 0 \\ \Rightarrow x &= 5 \text{ cm}\end{aligned}$$

$$\text{Also, } \frac{d^2v}{dx^2} = -12\pi x$$

$$\text{At } x = 5, \frac{d^2v}{dx^2} = -60\pi x < 0$$

$\therefore x = 5$  is point of local maxima.

Hence,

$$\text{The maximum volume of cylinder is} = \pi (75 - 25) \times 10 = 500\pi \text{ cm}^3.$$

## Maxima and Minima 18.5 Q27

Let  $x$  and  $y$  be two positive numbers with

$$x^2 + y^2 = r^2 \quad \text{--- (i)}$$

$$\text{Let } S = x + y \quad \text{--- (ii)}$$

$$\therefore S = x + \sqrt{r^2 - x^2} \quad \text{from (ii)}$$

$$\therefore \frac{dS}{dx} = 1 - \frac{x}{\sqrt{r^2 - x^2}}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 1 - \frac{x}{\sqrt{r^2 - x^2}} = 0$$

$$\Rightarrow x = \sqrt{r^2 - x^2}$$

$$\Rightarrow 2x^2 = r^2$$

$$\Rightarrow x = \frac{r}{\sqrt{2}}, \frac{r}{\sqrt{2}}$$

$\therefore x$  &  $y$  are positive numbers

$$\therefore x = \frac{r}{\sqrt{2}}$$

$$\text{Also, } \frac{d^2S}{dx^2} = \frac{-\left(\sqrt{r^2 - x^2} + \frac{x^2}{\sqrt{r^2 - x^2}}\right)}{r^2 - x^2}$$

$$\text{At, } x = \frac{r}{\sqrt{2}}, \frac{d^2S}{dx^2} = - \left[ \frac{\frac{r^2}{\sqrt{2}} + \frac{\frac{r^2}{2}}{\frac{r}{\sqrt{2}}}}{\frac{r^2}{2}} \right] < 0$$

Since  $\frac{d^2S}{dx^2} < 0$ , the sum is largest when  $x = y = \frac{r}{\sqrt{2}}$



The given equation of parabola is

$$x^2 = 4y \quad \text{---(i)}$$

Let  $P(x, y)$  be the nearest point on (i) from the point  $A(0, 5)$

Let  $S$  be the square of the distance of  $P$  from  $A$ .

$$\therefore S = x^2 + (y - 5)^2 \quad \text{---(ii)}$$

From (i),

$$S = 4y + (y - 5)^2$$

$$\Rightarrow \frac{dS}{dy} = 4 + 2(y - 5)$$

For maxima or minima, we have

$$\frac{dS}{dy} = 0$$

$$\Rightarrow 4 + 2(y - 5) = 0$$

$$\Rightarrow 2y = 6$$

$$\Rightarrow y = 3$$

From (i)

$$x^2 = 12$$

$$\therefore x = \pm 2\sqrt{3}$$

$$\Rightarrow P = (2\sqrt{3}, 3) \text{ and } P' = (-2\sqrt{3}, 3)$$

Now,

$$\frac{d^2S}{dy^2} = 2 > 0$$

$\therefore P$  and  $P'$  are the point of local minima

Hence, the nearest points are  $P(2\sqrt{3}, 3)$  and  $P'(-2\sqrt{3}, 3)$ .

Let  $P(x, y)$  be a point on  
 $y^2 = 4x$  --- (i)

Let  $S$  be the square of the distance between  $A(2, -8)$  and  $P$ .

$\therefore S = (x - 2)^2 + (y + 8)^2$  --- (ii)

Using (i),

$$S = \left(\frac{y^2}{4} - 2\right)^2 + (y + 8)^2$$

$$\therefore \frac{dS}{dy} = 2\left(\frac{y^2}{4} - 2\right) \times \frac{y}{2} + 2(y + 8)$$

$$= \frac{y^3 - 8y}{4} + 2y + 16$$

$$= \frac{y^3}{4} + 16$$

For maxima and minima,

$$\frac{dS}{dy} = 0$$

$$\Rightarrow \frac{y^3}{4} + 16 = 0$$

$$\Rightarrow y = -4$$

Now,

$$\frac{d^2S}{dy^2} = \frac{3y^2}{4}$$

At  $y = -4$ ,  $\frac{d^2S}{dy^2} = 12 > 0$

$\therefore y = -4$  is the point of local minima

From (i)

$$x = \frac{y^2}{4} = 4$$

Thus, the required point is  $(4, -4)$  nearest to  $(2, -8)$ .

Let  $P(x, y)$  be a point on the curve,  
 $x^2 = 8y$  ---(i)

Let  $A = (2, 4)$  be a point and

let  $S =$  square of the distance between  $P$  and  $A$

$$\therefore S = (x - 2)^2 + (y - 4)^2 \quad \text{---(ii)}$$

Using (i), we get

$$S = (x - 2)^2 + \left(\frac{x^2}{8} - 4\right)^2$$

$$\begin{aligned} \therefore \frac{dS}{dy} &= 2(x - 2) + 2\left(\frac{x^2}{8} - 4\right) \times \frac{2x}{8} \\ &= 2(x - 2) + \frac{(x^2 - 32)x}{16} \end{aligned}$$

$$\begin{aligned} \text{Also, } \frac{d^2S}{dx^2} &= 2 + \frac{1}{16}[x^2 - 32 + 2x^2] \\ &= 2 + \frac{1}{16}[3x^2 - 32] \end{aligned}$$

For maxima and minima,

$$\begin{aligned} \frac{dS}{dx} &= 0 \\ \Rightarrow 2(x - 2) + \frac{x(x^2 - 32)}{16} &= 0 \\ \Rightarrow 32x - 64 + x^3 - 32x &= 0 \\ \Rightarrow x^3 - 64 &= 0 \\ \Rightarrow x &= 4 \end{aligned}$$

Now,

$$\text{At } x = 4, \quad \frac{d^2S}{dx^2} = 2 + \frac{1}{16}[16 \times 3 - 32] = 2 + 1 = 3 > 0$$

$\therefore x = 4$  is point of local minima

From (i)

$$y = \frac{x^2}{8} = 2$$

Thus,  $P(4, 2)$  is the nearest point.

Let  $P(x, y)$  be a point on the curve  $x^2 = 2y$  which is closest to  $A(0, 5)$

Let  $S$  = square of the length of  $AP$

$$\Rightarrow S = x^2 + (y - 5)^2 \quad \text{---(ii)}$$

Using (i),

$$S = 2y + (y - 5)^2$$

$$\therefore \frac{dS}{dy} = 2 + 2(y - 5)$$

For maxima and minima,

$$\frac{dS}{dy} = 0$$

$$\Rightarrow 2 + 2y - 10 = 0$$

$$\Rightarrow y = 4$$

Now,

$$\frac{d^2S}{dy^2} = 2 > 0$$

$\therefore y = 4$  is the point of local minima

From (i)

$$r = \pm 2\sqrt{2}$$

Hence,  $(\pm 2\sqrt{2}, 4)$  is the closest point on the curve to  $A(0, 5)$ .

The given equations are

$$y = x^2 + 7x + 2 \quad \text{---(i)}$$

$$\text{and } y = 3x - 3 \quad \text{---(ii)}$$

Let  $P(x, y)$  be the point on parabola (i) which is closest to the line (ii)

Let  $S$  be the perpendicular distance from  $P$  to the line (ii).

$$\begin{aligned} \therefore S &= \frac{|y - 3x + 3|}{\sqrt{1^2 + (-3)^2}} \\ \Rightarrow S &= \frac{|x^2 + 7x + 2 - 3x + 3|}{\sqrt{10}} \quad \text{---(iii)} \\ \Rightarrow \frac{dS}{dx} &= \frac{2x + 4}{\sqrt{10}} \end{aligned}$$

For maxima or minima, we have

$$\begin{aligned} \frac{dS}{dx} &= 0 \\ \Rightarrow \frac{2x + 4}{\sqrt{10}} &= 0 \\ \Rightarrow x &= -2 \end{aligned}$$

From (i)

$$y = 4 - 14 + 2 = -8$$

Now,

$$\frac{d^2S}{dx^2} = \frac{2}{\sqrt{10}} > 0$$

$\therefore (x = -2, y = -8)$  is the point of local minima,

Hence,

The closest point on the parabola to the line  $y = 3x - 3$  is  $(-2, -8)$ .

Let  $P(x, y)$  be a point on the curve  $y^2 = 2x$  which is minimum distance from the point  $A(1, 4)$ .

Let

$S$  = square of the length of  $AP$

$$S = (x - 1)^2 + (y - 4)^2$$

Using this equation, we have

$$S = x^2 + 1 - 2x + y^2 + 16 - 8y$$

$$S = x^2 - 2x + 2x + 17 - 8y$$

$$S = \frac{y^4}{4} - 8y + 17 \quad \left[ \text{Since } x = \frac{y^2}{2} \right]$$

$$\frac{dS}{dy} = y^3 - 8$$

For maxima and minima, we have

$$\frac{dS}{dy} = 0$$

$$y^3 - 8 = 0$$

$$y^3 = 8$$

$$y = 2$$

Now,

$$\frac{d^2S}{dy^2} = 3y^2$$

$$\frac{d^2S}{dy^2} = 12 > 0$$

$\therefore y = 2$  is minimum point

We have

$$x = \frac{y^2}{2}$$

$$= \frac{4}{2}$$

$$= 2$$

Hence,  $(2, 2)$  is at a minimum distance from the point  $(1, 4)$ .

The given equation of curve is

$$y = x^3 + 3x^2 + 2x - 27 \quad \text{--- (i)}$$

Slope of (i)

$$m = \frac{dy}{dx} = -3x^2 + 6x + 2 \quad \text{--- (ii)}$$

Now,

$$\frac{dm}{dx} = -6x + 6$$

and  $\frac{d^2m}{dx^2} = -6 < 0$

For maxima and minima,

$$\frac{dm}{dx} = 0$$

$$\Rightarrow -6x + 6 = 0$$

$$\Rightarrow x = 1$$

$$\therefore \frac{d^2m}{dx^2} = -6 < 0$$

$\therefore x = 1$  is point of local maxima

Hence, maximum slope =  $-3 + 6 + 2 = 5$

**Maxima and Minima 18.5 Q35**

We have,

Cost of producing  $x$  radio sets is Rs.  $\frac{x^2}{4} + 35x + 25$

Selling price of  $x$  radio is Rs.  $x \left( 50 - \frac{x}{2} \right)$

So,

Profit on  $x$  radio sets is

$$P = \text{Rs} \left( 50x - \frac{x^2}{2} - \frac{x^2}{4} - 35x - 25 \right)$$

$$\begin{aligned} \therefore \frac{dP}{dx} &= 50 - x - \frac{x}{2} - 35 \\ &= 15 - \frac{3}{2}x \end{aligned}$$

For maxima and minima,

$$\frac{dP}{dx} = 0$$

$$\Rightarrow 15 - \frac{3}{2}x = 0$$

$$\Rightarrow x = 10$$

Also,

$$\frac{d^2P}{dx^2} = \frac{-3}{2} < 0$$

$\therefore x = 10$  is the point of local maxima

Hence, the daily output should be 10 radio sets.



### Maxima and Minima 18.5 Q35

We have,

Cost of producing  $x$  radio sets is Rs.  $\frac{x^2}{4} + 35x + 25$

Selling price of  $x$  radio is Rs.  $x \left( 50 - \frac{x}{2} \right)$

So,

Profit on  $x$  radio sets is

$$P = \text{Rs} \left( 50x - \frac{x^2}{2} - \frac{x^2}{4} - 35x - 25 \right)$$

$$\begin{aligned}\therefore \frac{dP}{dx} &= 50 - x - \frac{x}{2} - 35 \\ &= 15 - \frac{3}{2}x\end{aligned}$$

For maxima and minima,

$$\frac{dP}{dx} = 0$$

$$\Rightarrow 15 - \frac{3}{2}x = 0$$

$$\Rightarrow x = 10$$

Also,

$$\frac{d^2P}{dx^2} = \frac{-3}{2} < 0$$

$\therefore x = 10$  is the point of local maxima

Hence, the daily output should be 10 radio sets.

### Maxima and Minima 18.5 Q36

Let  $S(x)$  be the selling price of  $x$  items and let  $C(x)$  be the cost price of  $x$  items.

$$\text{Then, we have } S(x) = \left(5 - \frac{x}{100}\right)x = 5x - \frac{x^2}{100}$$

$$\text{and } C(x) = \frac{x}{5} + 500$$

Thus, the profit function  $P(x)$  is given by

$$P(x) = S(x) - C(x) = 5x - \frac{x^2}{100} - \frac{x}{5} - 500 = \frac{24}{5}x - \frac{x^2}{100} - 500$$

$$\therefore P'(x) = \frac{24}{5} - \frac{x}{50}$$

$$\text{Now, } P'(x) = 0$$

$$\Rightarrow \frac{24}{5} - \frac{x}{50} = 0$$

$$\Rightarrow x = \frac{24}{5} \times 50 = 240$$

$$\text{Also } P''(x) = -\frac{1}{50}$$

$$\text{So, } P''(240) = -\frac{1}{50} < 0$$

Thus,  $x = 240$  is a point of maxima.

Hence, the manufacturer can earn maximum profit, if he sells 240 items.

Let  $l$  be the length of side of square base of the tank and  $h$  be the height of tank.

Then,

$$\text{Volume of tank } (v) = l^2 h$$

$$\text{Total surface area } (s) = l^2 + 4lh$$

Since the tank holds a given quantity of water the volume ( $v$ ) is constant.

$$\therefore v = l^2 h \quad \text{---(i)}$$

Also, cost of lining with lead will be least if the total surface area is least.

So we need to minimise the surface area.

$$\therefore S = l^2 + 4lh \quad \text{---(ii)}$$

Now,

From (i) and (ii)

$$S = l^2 + \frac{4v}{l}$$

$$\therefore \frac{ds}{dl} = 2l - \frac{4v}{l^2}$$

For maximum and minimum

$$\frac{ds}{dl} = 0$$

$$\Rightarrow 2l - \frac{4v}{l^2} = 0$$

$$\Rightarrow 2l^3 - 4v = 0$$

$$\Rightarrow l^3 = 2v = 2l^2 h$$

$$\Rightarrow l^2[l - 2h] = 0$$

$$\Rightarrow l = 0 \text{ or } 2h$$

$l = 0$  is not possible.

$$\therefore l = 2h$$

Now,


$$\frac{d^2s}{dl^2} = 2 + \frac{8v}{l^3}$$

$$\text{At } l = 2h, \frac{d^2s}{dl^2} > 0 \quad \text{for all } h.$$

$$\therefore l = 2h \text{ is point of local minima}$$

$$\therefore S \text{ is minimum when } l = 2h$$

Let  $ABCDEFGH$  be a box of constant volume  $c$ . We are given that the box is twice as long as its width.



$$\therefore \text{Let } BF = x$$

$$\Rightarrow AB = 2x$$

Cost of material of top and front side =  $3 \times$  cost of material of the bottom of the box.

$$\Rightarrow 2x \times x + xh + xh + 2xh + 2xh = 3 \times 2x^2$$

$$\Rightarrow 2x^2 + 2xh + 4xh = 6x^2$$

$$\Rightarrow 4x^2 - 6xh = 0$$

$$\Rightarrow 2x(2x - 3h) = 0$$

$$\Rightarrow x = \frac{3h}{2} \text{ or } h = \frac{2x}{3}$$

Volume of box =  $2x \times x \times h$

$$\Rightarrow c = 2x^2h$$

$$\Rightarrow h = \frac{c}{2x^2} \quad \text{---(ii)}$$

Now,

$$S = \text{Surface area of box} = 2(2x^2 + 2xh + xh)$$

$$\Rightarrow S = 2(2x^2 + 3xh)$$

From (i)

$$S = 2\left(2x^2 + \frac{3xc}{2x^2}\right)$$

$$\Rightarrow S = 2\left(2x^2 + \frac{3c}{2x}\right)$$

For maxima and minima,

$$\frac{dS}{dx} = 2\left(4x - \frac{3c}{2x^2}\right) = 0$$

$$\Rightarrow 8x^3 - 3c = 0$$

$$\Rightarrow x = \left(\frac{3c}{8}\right)^{\frac{1}{3}}$$

Now,

$$\frac{d^2s}{dx^2} = 2 \left( 4 + 3 \frac{c}{x^3} \right) > 0 \text{ as } x = \left( \frac{3c}{8} \right)^{\frac{1}{3}}$$

$$x = \left( \frac{3c}{8} \right)^{\frac{1}{3}} \text{ is point of local minima}$$

∴ Most economic dimension will be

$$x = \text{width} = \left( \frac{3c}{8} \right)^{\frac{1}{3}}$$

$$2x = \text{length} = 2 \left( \frac{3c}{8} \right)^{\frac{1}{3}}$$

$$h = \text{height} = \frac{2x}{3} = \frac{2}{3} \left( \frac{3c}{8} \right)^{\frac{1}{3}}.$$

**Maxima and Minima 18.5 Q39**

Let  $s$  be the sum of the surface areas of a sphere and a cube.

$$\therefore s = 4\pi r^2 + 6l^2 \quad \text{---(i)}$$

Let  $v$  = volume of sphere + volume of cube

$$\Rightarrow v = \frac{4}{3}\pi r^3 + l^3 \quad \text{---(ii)}$$

From (i)

$$l = \sqrt{\frac{s - 4\pi r^2}{6}}$$

$$\therefore v = \frac{4}{3}\pi r^3 + \left(\frac{s - 4\pi r^2}{6}\right)^{\frac{3}{2}}$$

$$\therefore \frac{dv}{dr} = 4\pi r^2 + \frac{3}{2} \left(\frac{s - 4\pi r^2}{6}\right)^{\frac{1}{2}} \times \left(\frac{-4\pi}{6}\right)^{\frac{1}{2}}$$

For maxima and minima,

$$\frac{dv}{dr} = 0$$

$$\Rightarrow 4\pi r^2 = \frac{\pi}{6} (s - 4\pi r^2)^{\frac{1}{2}} \times 2r = 0$$

$$\Rightarrow 2r\pi [2r - l] = 0$$

$$\therefore r = 0, \quad \frac{l}{2}$$

Now,

$$\frac{d^2v}{dr^2} = 8\pi r - \frac{2\pi}{\sqrt{6}} \left[ (s - 4\pi r^2)^{\frac{1}{2}} \right] - \frac{8\pi r^2}{2(s - 4\pi r^2)^{\frac{1}{2}}}$$

$$\text{At } r = \frac{l}{2}$$

$$\frac{d^2v}{dr^2} = \pi \frac{l}{2} - \frac{2\pi}{\sqrt{6}} \left[ \sqrt{6}l - \frac{8\pi \frac{l^2}{4}}{2\sqrt{6}l} \right] = 4\pi l - \frac{2\pi}{\sqrt{6}} \left[ \frac{12l^2 - 2\pi l^2}{2\sqrt{6}l} \right]$$

Let ABCDEF be a half cylinder with rectangular base and semi-circular ends.

Here AB = height of the cylinder

AB = h

Let r be the radius of the cylinder.

Volume of the half cylinder is  $V = \frac{1}{2} \pi r^2 h$

$$\Rightarrow \frac{2V}{\pi r^2} = h$$

$\therefore$  TSA of the half cylinder is

S = LSA of the half cylinder + area of two semi-circular ends + area of the rectangle (base)

$$S = \pi r h + \frac{\pi r^2}{2} + \frac{\pi r^2}{2} + h \times 2r$$

$$S = (\pi r + 2r)h + \pi r^2$$

$$S = (\pi r + 2r) \frac{2V}{\pi r^2} + \pi r^2$$

$$S = (\pi + 2) \frac{2V}{\pi r} + \pi r^2$$

Differentiate S wrt r we get,

$$\frac{ds}{dr} = \left[ (\pi + 2) \times \frac{2V}{\pi} \left( \frac{-1}{r^2} \right) + 2\pi r \right]$$

For maximum and minimum values of S, we have  $\frac{ds}{dr} = 0$

$$\Rightarrow (\pi + 2) \times \frac{2V}{\pi} \left( \frac{-1}{r^2} \right) + 2\pi r = 0$$

$$\Rightarrow (\pi + 2) \times \frac{2V}{\pi r^2} = 2\pi r$$

But  $2r = D$

$$\therefore h : D = \pi : \pi + 2$$

Differentiate  $\frac{ds}{dr}$  wrt r we get,

$$\frac{d^2s}{dr^2} = (\pi + 2) \times \frac{2V}{\pi} \times \frac{2}{r^3} + 2\pi > 0$$


Thus S will be minimum when h : D is  $\pi : \pi + 2$ .

Height of the cylinder : Diameter of the circular end

$$\pi : \pi + 2$$

## Maxima and Minima 18.5 Q41

Let ABCD be the cross-sectional area of the beam which is cut from a circular log of radius a.



$$\therefore AO = a \Rightarrow AC = 2a$$

Let  $x$  be the width of log and  $y$  be the depth of log  $ABCD$

Let  $S$  be the strength of the beam according to the question,

$$S = xy^2 \quad \text{---(i)}$$

In  $\triangle ABC$

$$x^2 + y^2 = (2a)^2$$

$$\Rightarrow y = (2a)^2 - x^2 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x \left( (2a)^2 - x^2 \right)$$

$$\Rightarrow \frac{dS}{dx} = (4a^2 - x^2) - 2x^2$$

$$\Rightarrow \frac{dS}{dx} = 4a^2 - 3x^2$$

For maxima or minima

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 4a^2 - 3x^2 = 0$$

$$\Rightarrow x^2 = \frac{4a^2}{3}$$

$$\therefore x = \frac{2a}{\sqrt{3}}$$

From (ii),

$$y^2 = 4a^2 - \frac{4a^2}{3} = \frac{8a^2}{3}$$

$$\therefore y = 2a \times \sqrt{\frac{2}{3}}$$

Now,

$$\frac{d^2S}{dx^2} = -6x$$

$$\text{At } x = \frac{2a}{\sqrt{3}}, y = \sqrt{\frac{2}{3}} 2a, \quad \frac{d^2S}{dx^2} = -\frac{12a}{\sqrt{3}} < 0$$

$$\therefore \left( x = \frac{2a}{\sqrt{3}}, y = \sqrt{\frac{2}{3}} 2a \right) \text{ is the point of local maxima.}$$

Hence,

$$\text{The dimension of strongest beam is width } = x = \frac{2a}{\sqrt{3}} \text{ and depth } = y = \sqrt{\frac{2}{3}} 2a.$$



### Maxima and Minima 18.5 Q42

Let  $l$  be a line through the point  $P(1, 4)$  that cuts the  $x$ -axis and  $y$ -axis.

Now, equation of  $l$  is

$$y - 4 = m(x - 1)$$

$$\therefore x\text{-Intercept is } \frac{m-4}{m} \text{ and } y\text{-Intercept is } 4-m$$

$$\text{Let } S = \frac{m-4}{m} + 4 - m$$

$$\therefore \frac{dS}{dm} = +\frac{4}{m^2} - 1$$

For maxima and minima,

$$\frac{dS}{dm} = 0$$

$$\Rightarrow \frac{4}{m^2} - 1 = 0$$

$$\Rightarrow m = \pm 2$$

Now,

$$\frac{d^2S}{dm^2} = -\frac{8}{m^3}$$

$$\text{At } m = 2, \frac{d^2S}{dm^2} = -1 < 0$$

$$m = -2, \frac{d^2S}{dm^2} = 1 > 0$$

$$\therefore m = -2 \text{ is point of local minima.}$$

$\therefore$  least value of sum of intercept is


$$\begin{aligned} & \frac{m-4}{m} + 4 - m \\ &= 3 + 6 = 9 \end{aligned}$$

### Maxima and Minima 18.5 Q43

The area of the page  $PQRS$  is  $150 \text{ cm}^2$

Also,  $AB + CD = 3 \text{ cm}$

$EF + GH = 2 \text{ cm}$



Let  $x$  and  $y$  be the combined width of margin at the top and bottom and the sides respectively.

$\therefore x = 3 \text{ cm}$  and  $y = 2 \text{ cm}$ .

Now, area of printed matter = area of  $P'Q'R'S'$

$$\Rightarrow A = P'Q' \times Q'R'$$

$$\Rightarrow A = (b - y)(l - x)$$

$$\Rightarrow A = (b - 2)(l - 3) \quad \text{---(i)}$$

Also,

$$\text{Area of } PQRS = 150 \text{ cm}^2$$

$$\Rightarrow lb = 150 \quad \text{---(ii)}$$

From (i) and (ii)

$$A = (b - 2) \left( \frac{150}{b} - 3 \right)$$

$\therefore$  For maximum and minimum,

$$\frac{dA}{db} = \left( \frac{150}{b} - 3 \right) + (b - 2) \left( -\frac{150}{b^2} \right) = 0$$

$$\Rightarrow \frac{(150 - 3b)}{b} + (-150) \frac{(b - 2)}{b^2} = 0$$

$$\Rightarrow 150b - 3b^2 - 150b + 300 = 0$$

$$\Rightarrow -3b^2 + 300 = 0$$

$$\Rightarrow b = 10$$

From (ii)

$$l = 15$$

Now,

$$\frac{d^2A}{db^2} = \frac{-150}{b^2} - 150 \left[ -\frac{1}{b^2} + \frac{4}{b^3} \right]$$

At  $b = 10$

$$\begin{aligned}\frac{d^2A}{db^2} &= -\frac{15}{10} - 150\left[-\frac{1}{100} + \frac{4}{1000}\right] \\ &= -1.5 - .15[-10 + 4] \\ &= -1.5 + .9 \\ &= -0.6 < 0\end{aligned}$$

$\therefore b = 10$  is point of local maxima.

Hence,

The required dimension will be  $l = 15$  cm,  $b = 10$  cm.

### Maxima and Minima 18.5 Q44

The space  $s$  described in time  $t$  by a moving particle is given by

$$s = t^5 - 40t^3 + 30t^2 + 80t - 250$$

$$\therefore \text{velocity} = \frac{ds}{dt} = 5t^4 - 120t^2 + 60t + 80$$

$$\text{Acceleration} = a = \frac{d^2s}{dt^2} = 20t^3 - 240t + 60t \quad \text{---(i)}$$

Now,

$$\frac{da}{dt} = 60t^2 - 240$$

For maxima and minima,

$$\frac{da}{dt} = 0$$

$$\Rightarrow 60t^2 - 240 = 0$$

$$\Rightarrow 60(t^2 - 4) = 0$$

$$\Rightarrow t = 2$$

Now,

$$\frac{d^2a}{dt^2} = 120t$$

$$\text{At } t = 2, \frac{d^2a}{dt^2} = 240 > 0$$

$\therefore t = 2$  is point of local minima

Hence, minimum acceleration is  $160 - 480 + 60 = -260$ .

## Maxima and Minima 18.5 Q45

We have,

$$\text{Distance, } s = \frac{t^4}{4} - 2t^3 + 4t^2 - 7$$

$$\text{Velocity, } v = \frac{ds}{dt} = t^3 - 6t^2 + 8t$$

$$\text{Acceleration, } a = \frac{d^2s}{dt^2} = 3t^2 - 12t + 8$$

For velocity to be maximum and minimum,

$$\frac{dv}{dt} = 0$$

$$\Rightarrow 3t^2 - 12t + 8 = 0$$

$$\Rightarrow t = \frac{12 \pm \sqrt{144 - 96}}{6}$$

$$= 2 \pm \frac{4\sqrt{3}}{6}$$

$$\therefore t = 2 + \frac{2}{\sqrt{3}}, 2 - \frac{2}{\sqrt{3}}$$

Now,

$$\frac{d^2v}{dt^2} = 6t - 12$$

$$\text{At } t = 2 - \frac{2}{\sqrt{3}}, \frac{d^2v}{dt^2} = 6\left(2 - \frac{2}{\sqrt{3}}\right) - 12 = \frac{-12}{\sqrt{3}} < 0$$

$$t = 2 + \frac{2}{\sqrt{3}}, \frac{d^2v}{dt^2} = 6\left(2 + \frac{2}{\sqrt{3}}\right) - 12 = \frac{12}{\sqrt{3}} > 0$$

$$\therefore \text{At } t = 2 - \frac{2}{\sqrt{3}}, \text{ velocity is maximum}$$

For acceleration to be maximum and minimum

$$\frac{da}{dt} = 0$$

$$\Rightarrow 6t - 12 = 0$$

$$\Rightarrow t = 2$$

Now,

$$\frac{d^2a}{dt^2} = 6 > 0$$

$$\therefore \text{At } t = 2 \text{ Acceleration is minimum.}$$

