


RD Sharma
Solutions
Class 12 Maths
Chapter 18
Ex 18.1



Maxima and Minima 18.1 Q1

$$f(x) = 4x^2 - 4x + 4 \quad \text{on } \mathbb{R}$$

$$= 4x^2 - 4x + 1 + 3$$

$$= (2x - 1)^2 + 3$$

$$\therefore (2x - 1)^2 \geq 0$$

$$\Rightarrow (2x - 1)^2 + 3 \geq 3$$

$$\Rightarrow f(x) \geq f\left(\frac{1}{2}\right)$$

Thus, the minimum value of $f(x)$ is 3 at $x = \frac{1}{2}$

Since, $f(x)$ can be made as large as we please. Therefore maximum value does not exist

Maxima and Minima 18.1 Q2

The given function is $f(x) = -(x - 1)^2 + 2$

It can be observed that $(x - 1)^2 \geq 0$ for every $x \in \mathbb{R}$.

Therefore, $f(x) = -(x - 1)^2 + 2 \leq 2$ for every $x \in \mathbb{R}$.

The maximum value of f is attained when $(x - 1) = 0$.

$$(x - 1) = 0 \Rightarrow x = 1$$

$$\therefore \text{Maximum value of } f = f(1) = -(1 - 1)^2 + 2 = 2$$

Hence, function f does not have a minimum value.

Maxima and Minima 18.1 Q3

$$f(x) = |x + 2| \quad \text{on } \mathbb{R}$$

$$\therefore |x + 2| \geq 0 \quad \text{for } x \in \mathbb{R}$$

$$\Rightarrow f(x) \geq 0 \quad \text{for all } x \in \mathbb{R}$$

So, the minimum value of $f(x)$ is 0, which attains at $x = -2$

Clearly, $f(x) = |x + 2|$ does not have the maximum value.

Maxima and Minima 18.1 Q4

$$h(x) = \sin 2x + 5$$

We know that $-1 \leq \sin 2x \leq 1$.

$$\Rightarrow -1 + 5 \leq \sin 2x + 5 \leq 1 + 5$$

$$\Rightarrow 4 \leq \sin 2x + 5 \leq 6$$

Hence, the maximum and minimum values of h are 6 and 4 respectively.

Maxima and Minima 18.1 Q5

$$f(x) = |\sin 4x + 3|$$

We know that $-1 \leq \sin 4x \leq 1$.

$$\Rightarrow 2 \leq \sin 4x + 3 \leq 4$$

$$\Rightarrow 2 \leq |\sin 4x + 3| \leq 4$$

Hence, the maximum and minimum values of f are 4 and 2 respectively.

Maxima and Minima 18.1 Q6

$$f(x) = 2x^3 + 5 \text{ on } \mathbb{R}$$

Here, we observe that the values of $f(x)$ increase when the values of x are increased and $f(x)$ can be made as large as possible, we please.

So, $f(x)$ does not have the maximum value.

Similarly $f(x)$ can be made as small as we please by giving smaller values to x .

So, $f(x)$ does not have the minimum value.

Maxima and Minima 18.1 Q7

$$g(x) = -|x+1| + 3$$

We know that $-|x+1| \leq 0$ for every $x \in \mathbb{R}$.

Therefore, $g(x) = -|x+1| + 3 \leq 3$ for every $x \in \mathbb{R}$.

The maximum value of g is attained when $|x+1| = 0$

$$|x+1| = 0$$

$$\Rightarrow x = -1$$

$$\therefore \text{Maximum value of } g = g(-1) = -|-1+1| + 3 = 3$$

Hence, function g does not have a minimum value.

Maxima and Minima 18.1 Q8

$$f(x) = 16x^2 - 16x + 28 \text{ on } R$$

$$= 16x^2 - 16x + 4 + 24$$

$$= (4x - 2)^2 + 24$$

Now,

$$(4x - 2)^2 \geq 0 \text{ for all } x \in R$$

$$\Rightarrow (4x - 2)^2 + 24 \geq 24 \text{ for all } x \in R$$

$$\Rightarrow f(x) \geq f\left(\frac{1}{2}\right)$$

Thus, the minimum value of $f(x)$ is 24 at $x = \frac{1}{2}$

Since $f(x)$ can be made as large as possible by giving difference values to x .

Thus, maximum values does not exist.

Maxima and Minima 18.1 Q9

$$f(x) = x^3 - 1 \text{ on } R$$


Here, we observe that the values of $f(x)$ increases when the values of x are increased and $f(x)$ can be made as large as we please by giving large values to x .

So, $f(x)$ does not have the maximum value.

Similarly, $f(x)$ can be made as small as we please by giving smaller values to x .

So, $f(x)$ does not have the minimum value.

RD Sharma
Solutions
Class 12 Maths
Chapter 18
Ex 18.2



Maxima and Minima Ex 18.2 Q1

$$f(x) = (x - 5)^4$$

$$\therefore f'(x) = 4(x - 5)^3$$

For local maxima and minima

$$f'(x) = 0$$

$$\Rightarrow 4(x - 5)^3 = 0$$

$$\Rightarrow x - 5 = 0$$

$$\Rightarrow x = 5$$

$f'(x)$ changes from -ve to +ve as x passes through 5.

So, $x = 5$ is the point of local minima

Thus, local minimum value is $f(5) = 0$

Maxima and Minima Ex 18.2 Q2

$$g(x) = x^3 - 3x$$

$$\therefore g'(x) = 3x^2 - 3$$

Now,

$$g'(x) = 0 \Rightarrow 3x^2 = 3 \Rightarrow x = \pm 1$$

$$g''(x) = 6x$$

$$g''(1) = 6 > 0$$

$$g''(-1) = -6 < 0$$

By second derivative test, $x = 1$ is a point of local minima and local minimum value of g at $x = 1$ is $g(1) = 1^3 - 3 = 1 - 3 = -2$. However,

$x = -1$ is a point of local maxima and local maximum value of g at

$$x = -1 \text{ is } g(-1) = (-1)^3 - 3(-1) = -1 + 3 = 2.$$

Maxima and Minima Ex 18.2 Q3

$$f(x) = x^3(x-1)^2$$

$$\begin{aligned} \therefore f'(x) &= 3x^2(x-1)^2 + 2x^3(x-1) \\ &= (x-1)(3x^2(x-1) + 2x^3) \\ &= (x-1)(3x^3 - 3x^2 + 2x^3) \\ &= (x-1)(5x^3 - 3x^2) \\ &= x^2(x-1)(5x-3) \end{aligned}$$

For all maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow x^2(x-1)(5x-3) = 0$$

$$\Rightarrow x = 0, 1, \frac{3}{5}$$

At $x = \frac{3}{5}$ $f'(x)$ changes from +ve to -ve

$\therefore x = \frac{3}{5}$ is point of minima.

At $x = 1$ $f'(x)$ changes from -ve to +ve

$\therefore x = 1$ is point of maxima

Maxima and Minima Ex 18.2 Q4

$$f(x) = (x-1)(x+2)^2$$

$$\begin{aligned}\therefore f'(x) &= (x+2)^2 + 2(x-1)(x+2) \\ &= (x+2)(x+2+2x-2) \\ &= (x+2)(3x)\end{aligned}$$

For point of maxima and minima

$$f'(x) = 0$$

$$\Rightarrow (x+2) \times 3x = 0$$

$$\Rightarrow x = 0, -2$$

At $x = -2$ $f'(x)$ changes from +ve to -ve

$\therefore x = -2$ is point of local maxima

At $x = 0$ $f'(x)$ changes from -ve to +ve

$\therefore x = 0$ is point of local minima

Thus, local min value = $f(0) = -4$

local max value = $f(-2) = 0$.

Maxima and Minima Ex 18.2 Q5

$$f(x) = (x-1)^3(x+1)^2$$

$$\begin{aligned}\therefore f'(x) &= 3(x-1)^2(x+1)^2 + 2(x-1)^3(x+1) \\ &= (x-1)^2(x+1)(3(x+1) + 2(x-1)) \\ &= (x-1)^2(x+1)(5x+1)\end{aligned}$$

For the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow (x-1)^2(x+1)(5x+1) = 0$$

$$\Rightarrow x = 1, -1, -\frac{1}{5}$$

Here,

At $x = -1$ $f'(x)$ changes from +ve to -ve so $x = -1$ is point of maxima.

At $x = -\frac{1}{5}$, $f'(x)$ changes from -ve to +ve so $x = -\frac{1}{5}$ is point of minima

Hence, local max value = 0

$$\text{local min value} = -\frac{3456}{3125}.$$

Maxima and Minima Ex 18.2 Q6

$$f(x) = x^3 - 6x^2 + 9x + 15$$

$$\begin{aligned}\therefore f'(x) &= 3x^2 - 12x + 9 \\ &= 3(x^2 - 4x + 3) \\ &= 3(x - 3)(x - 1)\end{aligned}$$

For, the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 3(x - 3)(x - 1) = 0$$

$$\Rightarrow x = 3, 1$$

At $x = 1$, $f'(x)$ changes from +ve to -ve

$\therefore x = 1$ is point of local maxima

At $x = 3$, $f'(x)$ changes from -ve to +ve

$\therefore x = 3$ is point of local minima

Hence, local max value = $f(1) = 19$

local min value = $f(3) = 15$.

Maxima and Minima Ex 18.2 Q7

$$f(x) = \sin 2x, \quad 0 < x, \pi$$

$$\therefore f'(x) = 2 \cos 2x$$

For, the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 2x = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$\Rightarrow x = \frac{\pi}{4}, \frac{3\pi}{4}$$

At $x = \frac{\pi}{4}$, $f'(x)$ changes from +ve to -ve

$\therefore x = \frac{\pi}{4}$ is point of local maxima

At $x = \frac{3\pi}{4}$, $f'(x)$ changes from -ve to +ve

$\therefore x = \frac{3\pi}{4}$ is point of local minima,

Hence, local max value = $f\left(\frac{\pi}{4}\right) = 1$

local min value = $f\left(\frac{3\pi}{4}\right) = -1$.

Maxima and Minima Ex 18.2 Q8

$$f(x) = \sin x - \cos x, 0 < x < 2\pi$$

$$\therefore f'(x) = \cos x + \sin x$$

$$f'(x) = 0 \Rightarrow \cos x = -\sin x \Rightarrow \tan x = -1 \Rightarrow x = \frac{3\pi}{4}, \frac{7\pi}{4} \in (0, 2\pi)$$

$$f''(x) = -\sin x + \cos x$$

$$f''\left(\frac{3\pi}{4}\right) = -\sin \frac{3\pi}{4} + \cos \frac{3\pi}{4} = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = -\sqrt{2} < 0$$

$$f''\left(\frac{7\pi}{4}\right) = -\sin \frac{7\pi}{4} + \cos \frac{7\pi}{4} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2} > 0$$

Therefore, by second derivative test, $x = \frac{3\pi}{4}$ is a point of local maxima and the local maximum value of f at $x = \frac{3\pi}{4}$ is

$$f\left(\frac{3\pi}{4}\right) = \sin \frac{3\pi}{4} - \cos \frac{3\pi}{4} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}. \text{ However, } x = \frac{7\pi}{4} \text{ is a point of local minima and the}$$

$$\text{local minimum value of } f \text{ at } x = \frac{7\pi}{4} \text{ is } f\left(\frac{7\pi}{4}\right) = \sin \frac{7\pi}{4} - \cos \frac{7\pi}{4} = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = -\sqrt{2}.$$

Maxima and Minima Ex 18.2 Q9

$$f(x) = \cos x, 0 < x < \pi$$

$$\therefore f'(x) = -\sin x$$

For, the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow -\sin x = 0$$

$$\Rightarrow x = 0, \text{ and } \pi$$

But, these two points lies outside the interval $(0, \pi)$

So, no local maxima and minima will exist in the interval $(0, \pi)$.

Maxima and Minima Ex 18.2 Q10

$$\therefore f'(x) = 2 \cos 2x - 1$$

For, the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 2 \cos 2x - 1 = 0$$

$$\Rightarrow \cos 2x = \frac{1}{2} = \cos \frac{\pi}{3}$$

$$\Rightarrow 2x = \frac{\pi}{3}, -\frac{\pi}{3}$$

$$\Rightarrow x = \frac{\pi}{6}, -\frac{\pi}{6}$$

At $x = -\frac{\pi}{6}$, $f'(x)$ changes from -ve to +ve

$\therefore x = -\frac{\pi}{6}$ is point of local minima

At $x = \frac{\pi}{6}$, $f'(x)$ changes from +ve to -ve

$\therefore x = \frac{\pi}{6}$ is point of local maxima

$$\text{Hence, local max value} = f\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} - \frac{\pi}{6}$$

$$\text{local min value} = f\left(-\frac{\pi}{6}\right) = -\frac{\sqrt{3}}{2} + \frac{\pi}{6}$$

Maxima and Minima Ex 18.2 Q11

$$f(x) = 2 \sin x - x, \quad -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$$

For checking the minima and maxima, we have

$$f'(x) = 2 \cos x - 1 = 0$$

$$\Rightarrow \cos x = \frac{1}{2} = \cos \frac{\pi}{3}$$

$$\Rightarrow x = -\frac{\pi}{3}, \frac{\pi}{3}$$

At $x = -\frac{\pi}{3}$, $f(x)$ changes from -ve to +ve

$$\Rightarrow x = -\frac{\pi}{3} \text{ is point of local minima with value} = -\sqrt{3} - \frac{\pi}{3}$$

At $x = \frac{\pi}{3}$, $f(x)$ changes from +ve to +ve

$$\Rightarrow x = \frac{\pi}{3} \text{ is point of local maxima with value} = \sqrt{3} - \frac{\pi}{3}$$

Maxima and Minima Ex 18.2 Q12

$$\begin{aligned}\therefore f'(x) &= \sqrt{1-x} + x \cdot \frac{1}{2\sqrt{1-x}}(-1) = \sqrt{1-x} - \frac{x}{2\sqrt{1-x}} \\ &= \frac{2(1-x) - x}{2\sqrt{1-x}} = \frac{2-3x}{2\sqrt{1-x}}\end{aligned}$$

$$f'(x) = 0 \Rightarrow \frac{2-3x}{2\sqrt{1-x}} = 0 \Rightarrow 2-3x = 0 \Rightarrow x = \frac{2}{3}$$

$$f''(x) = \frac{1}{2} \left[\frac{\sqrt{1-x}(-3) - (2-3x)\left(\frac{-1}{2\sqrt{1-x}}\right)}{1-x} \right]$$

$$= \frac{\sqrt{1-x}(-3) + (2-3x)\left(\frac{1}{2\sqrt{1-x}}\right)}{2(1-x)}$$

$$= \frac{-6(1-x) + (2-3x)}{4(1-x)^{\frac{3}{2}}}$$

$$= \frac{3x-4}{4(1-x)^{\frac{3}{2}}}$$

$$f''\left(\frac{2}{3}\right) = \frac{3\left(\frac{2}{3}\right) - 4}{4\left(1 - \frac{2}{3}\right)^{\frac{3}{2}}} = \frac{2-4}{4\left(\frac{1}{3}\right)^{\frac{3}{2}}} = \frac{-1}{2\left(\frac{1}{3}\right)^{\frac{3}{2}}} < 0$$

Therefore, by second derivative test, $x = \frac{2}{3}$ is a point of local maxima and the local maximum

value of f at $x = \frac{2}{3}$ is

$$f\left(\frac{2}{3}\right) = \frac{2}{3} \sqrt{1 - \frac{2}{3}} = \frac{2}{3} \sqrt{\frac{1}{3}} = \frac{2}{3\sqrt{3}} = \frac{2\sqrt{3}}{9}.$$

We have,

$$f(x) = x^3(2x - 1)^3$$

$$\begin{aligned}\therefore f'(x) &= 3x^2(2x - 1)^3 + 3x^3(2x - 1)^2 \times 2 \\ &= 3x^2(2x - 1)^2(2x - 1 + 2x) \\ &= 3x^2(4x - 1)\end{aligned}$$

For, the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 3x^2(4x - 1) = 0$$

$$\Rightarrow x = 0, \frac{1}{4}$$

At $x = \frac{1}{4}$, $f'(x)$ changes from -ve to +ve

$\therefore x = \frac{1}{4}$ is the point of local minima,

$$\therefore \text{local min value} = f\left(\frac{1}{4}\right) = \frac{-1}{512}.$$

Maxima and Minima Ex 18.2 Q14

We have,

$$f(x) = \frac{x}{2} + \frac{2}{x}, \quad x > 0$$

$$\therefore f'(x) = \frac{1}{2} - \frac{2}{x^2}$$

For the point of local maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{1}{2} - \frac{2}{x^2} = 0$$

$$\Rightarrow x^2 - 4 = 0$$

$$\Rightarrow x = \sqrt{4}, -\sqrt{4}$$

$$\Rightarrow x = 2, -2$$

At $x = 2$, $f'(x)$ changes from -ve to +ve

$\therefore x = 2$ is point of local minima.

$$\therefore \text{local min value} = f(2) = 2.$$

$$g(x) = \frac{1}{x^2 + 2}$$

$$\therefore g'(x) = \frac{-(2x)}{(x^2 + 2)^2}$$


$$g'(x) = 0 \Rightarrow \frac{-2x}{(x^2 + 2)^2} = 0 \Rightarrow x = 0$$

Now, for values close to $x = 0$ and to the left of 0, $g'(x) > 0$. Also, for values close to $x = 0$ and to the right of 0, $g'(x) < 0$.

Therefore, by first derivative test, $x = 0$ is a point of local maxima and the local maximum value of $g(0)$ is $\frac{1}{0+2} = \frac{1}{2}$.



RD Sharma
Solutions
Class 12 Maths
Chapter 18
Ex 18.3



Maxima and Minima 18.3 Q1(i)

$$f(x) = x^4 - 62x^2 + 120x + 9$$

$$\therefore f'(x) = 4x^3 - 124x + 120 = 4(x^3 - 31x + 30)$$

$$f''(x) = 12x^2 - 124 = 4(3x^2 - 31)$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 4(x^3 - 31x + 30) = 0$$

$$\Rightarrow 4(x^3 - 31x + 30) = 0$$

$$\Rightarrow x = 5, 1, -6$$

Now,

$$f''(5) = 176 > 0$$

$$\Rightarrow x = 5 \text{ is point of local minima}$$

$$f''(1) = -112 < 0$$

$$\Rightarrow x = 1 \text{ is point of local maxima}$$

$$f''(-6) = 308 > 0$$

$$\Rightarrow x = -6 \text{ is point of local minima}$$

$$\therefore \text{local max value} = f(1) = 68$$

$$\text{local min value} = f(5) = -316$$

$$\text{and } = f(-6) = -1647.$$

Maxima and Minima 18.3 Q1(ii)

We have,

$$f(x) = x^3 - 6x^2 + 9x + 15$$

$$\begin{aligned}\therefore f'(x) &= 3x^2 - 12x + 9 \\ &= 3(x^2 - 4x + 3) \\ f''(x) &= 6x - 12 \\ &= 6(x - 2)\end{aligned}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 3(x^2 - 4x + 3) = 0$$

$$\Rightarrow 3(x - 3)(x - 1) = 0$$

$$\Rightarrow x = 3, 1$$

Now,

$$f''(3) = 6 > 0$$

$\therefore x = 3$ is point of local minima

$$f''(1) = -6 < 0$$

$\therefore x = 1$ is point of local maxima

$$\therefore \text{local max value} = f(1) = 19$$

$$\text{local min value} = f(3) = 15.$$

Maxima and Minima 18.3 Q1(iii)

We have,

$$f(x) = (x - 1)(x + 2)^2$$

$$\begin{aligned}\therefore f'(x) &= (x + 2)^2 + 2(x - 1)(x + 2) \\ &= (x + 2)(x + 2 + 2x - 2) \\ &= (x + 2)(3x)\end{aligned}$$

$$\begin{aligned}\text{and, } f''(x) &= 3(x + 2) + 3x \\ &= 6x + 6\end{aligned}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 3x(x + 2) = 0$$

$$\Rightarrow x = 0, -2$$

Now,

$$f''(0) = 6 > 0$$

$\therefore x = 0$ is point of local minima

$$f''(-2) = -6 < 0$$

$\therefore x = -2$ is point of local maxima

$$\therefore \text{local max value} = f(-2) = 0$$

$$\text{local min value} = f(0) = -4.$$

Maxima and Minima 18.3 Q1(iv)

We have,

$$f(x) = \frac{2}{x} - \frac{2}{x^2}, \quad x > 0$$

$$\therefore f'(x) = \frac{-2}{x^2} + \frac{4}{x^3}$$

$$\text{and, } f''(x) = \frac{+4}{x^3} - \frac{12}{x^4}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{-2}{x^2} + \frac{4}{x^3} = 0$$

$$\Rightarrow \frac{-2(x-2)}{x^3} = 0$$

$$\Rightarrow x = 2$$

Now,

$$f''(2) = \frac{4}{8} - \frac{12}{6} = \frac{1}{2} - \frac{3}{4} = \frac{-1}{4} < 0$$

$\therefore x = 2$ is point of local maxima

$$\text{local max value} = f(2) = \frac{1}{2}.$$

Maxima and Minima 18.3 Q1(v)

We have,

$$f(x) = xe^x$$

$$\therefore f'(x) = e^x + xe^x = e^x(x+1)$$

$$\begin{aligned} f''(x) &= e^x(x+1) + e^x \\ &= e^x(x+2) \end{aligned}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow e^x(x+1) = 0$$

$$\Rightarrow x = -1$$

Now,

$$f''(-1) = e^{-1} = \frac{1}{e} > 0$$

$\therefore x = -1$ is point of local minima

Hence,

$$\text{local min value} = f(-1) = \frac{-1}{e}.$$

Maxima and Minima 18.3 Q1(vi)

We have,

$$f(x) = \frac{x}{2} + \frac{2}{x}, \quad x > 0$$

$$\therefore f'(x) = \frac{1}{2} - \frac{2}{x^2}$$

$$\text{and, } f''(x) = \frac{4}{x^3}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{1}{2} - \frac{2}{x^2} = 0$$

$$\Rightarrow \frac{x^2 - 4}{2x^2} = 0$$

$$\Rightarrow x = 2, -2$$

Now,

$$f''(2) = \frac{1}{2} > 0$$

$\therefore x = 2$ is point of minima

We will not consider $x = -2$ as $x > 0$

$\therefore$ local min value = $f(2) = 2$.

Maxima and Minima 18.3 Q1(vii)

We have,

$$f(x) = (x+1)(x+2)^{\frac{1}{3}}, \quad x \geq -2$$

$$\therefore f'(x) = (x+2)^{\frac{1}{3}} + \frac{1}{3}(x+1)(x+2)^{-\frac{2}{3}}$$

$$= (x+2)^{-\frac{2}{3}} \left(x+2 + \frac{1}{3}(x+1) \right)$$

$$= \frac{1}{3}(x+2)^{-\frac{2}{3}}(4x+7)$$

$$\text{and, } f''(x) = -\frac{2}{9}(x+2)^{-\frac{5}{3}}(4x+7) + \frac{1}{3}(x+2)^{-\frac{2}{3}} \times 4$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{1}{3}(x+2)^{-\frac{2}{3}}(4x+7) = 0$$

$$\Rightarrow x = -\frac{7}{4}$$

Now,

$$f''\left(-\frac{7}{4}\right) = \frac{4}{3}\left(-\frac{7}{4}+2\right)^{-\frac{2}{3}}$$

$\therefore x = -\frac{7}{4}$ is point of minima

$\therefore$ local min value = $f\left(-\frac{7}{4}\right) = \frac{-3}{4^{\frac{2}{3}}}$.

Maxima and Minima 18.3 Q1(viii)

We have,

$$f(x) = x\sqrt{32-x^2}, -5 \leq x \leq 5$$

$$\therefore f'(x) = \sqrt{32-x^2} + \frac{x}{2\sqrt{32-x^2}} \times (-2x)$$

$$= \frac{2(32-x^2) - 2x^2}{2\sqrt{32-x^2}}$$

$$= \frac{64-4x^2}{2\sqrt{32-x^2}}$$

$$\text{and, } f''(x) = \frac{2\sqrt{32-x^2} \times (-8x) - \frac{2(64-4x^2)}{2\sqrt{32-x^2}} \times (-2x)}{4(32-x^2)}$$

$$= \frac{-4(32-x^2) \times 8x + 4x(64-x^2)}{8(32-x^2)^{\frac{3}{2}}}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{4(16-x^2)}{2\sqrt{32-x^2}} = 0$$

$$\Rightarrow x = \pm 4$$

Now,

$$f''(4) = \frac{4 \times 4(64-16-8 \times 32+8 \times 16)}{8(32-16)^{\frac{3}{2}}} < 0$$

$\therefore x = 4$ is point of maxima

Maxima and Minima 18.3 Q1(ix)

Local Maximum value = $f(4)$

$$= 4\sqrt{32-4^2}$$

$$= 4\sqrt{32-16}$$

$$= 4\sqrt{16}$$

$$= 16$$

Local minimum at $x = -4$;

Local Minimum value = $f(-4)$

$$= -4\sqrt{32-(-4)^2}$$

$$= -4\sqrt{32-16}$$

$$= -4\sqrt{16}$$

$$= -16$$

Maxima and Minima 18.3 Q1(x)

$$f(x) = x + \frac{a^2}{x}$$

$$\therefore f'(x) = 1 - \frac{a^2}{x^2}$$

$$f''(x) = \frac{2a^2}{x^3}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow 1 - \frac{a^2}{x^2} = 0$$

$$\Rightarrow x^2 - a^2 = 0$$

$$\Rightarrow x = \pm a$$

Now,

$$f''(a) = \frac{2}{a} > 0 \text{ as } a > 0$$

$\therefore x = a$ is point of minima

$$f''(-a) = \frac{-2}{a} < 0 \text{ as } a > 0$$

$\therefore x = -a$ is point of maxima

Hence,

$$\text{local max value} = f(-a) = -2a$$

$$\text{local min value} = f(a) = 2a.$$

Maxima and Minima 18.3 Q1(xi)

$$f(x) = x\sqrt{2-x^2}$$

$$\begin{aligned}\therefore f'(x) &= \sqrt{2-x^2} - \frac{2x^2}{2\sqrt{2-x^2}} \\ &= \frac{2(2-x^2) - 2x^2}{2\sqrt{2-x^2}} \\ &= \frac{2-2x^2}{\sqrt{2-x^2}} \\ f''(x) &= \frac{\sqrt{2-x^2}(-4x) + \frac{(2-2x^2)2x}{\sqrt{2-x^2}}}{(\sqrt{2-x^2})^2} \\ &= \frac{-(2-x^2)4x + 4x - 4x^3}{(2-x^2)^{\frac{3}{2}}}\end{aligned}$$

For maxima and minima,

$$\begin{aligned}f'(x) &= 0 \\ \Rightarrow \frac{2(1-x^2)}{\sqrt{2-x^2}} &= 0 \\ \Rightarrow x &= \pm 1\end{aligned}$$

Now,

$$\begin{aligned}f''(1) &< 0 \\ \Rightarrow x = 1 &\text{ is point of local maxima} \\ f''(-1) &> 0 \\ \Rightarrow x = -1 &\text{ is point of local minima}\end{aligned}$$

Hence,

$$\text{local max value} = f(1) = 1$$

$$\text{local min value} = f(-1) = -1.$$

Maxima and Minima 18.3 Q1(xii)

$$f(x) = x + \sqrt{1-x}$$

$$\therefore f'(x) = 1 - \frac{1}{2\sqrt{1-x}} = \frac{2\sqrt{1-x} - 1}{2\sqrt{1-x}}$$

$$\therefore f'(x) = \frac{2\sqrt{1-x} \left(\frac{-1}{\sqrt{1-x}} \right) + \frac{(2\sqrt{1-x} - 1)}{\sqrt{1-x}}}{4(1-x)}$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow \frac{2\sqrt{1-x} - 1}{2\sqrt{1-x}} = 0$$

$$\Rightarrow \sqrt{1-x} = \frac{1}{2}$$

$$\Rightarrow x = 1 - \frac{1}{4} = \frac{3}{4}$$

Now,

$$f''\left(\frac{3}{4}\right) < 0$$

$$\Rightarrow x = \frac{3}{4} \text{ is point of local maxima}$$

Hence,

$$\text{local max value} = f\left(\frac{3}{4}\right) = \frac{5}{4}.$$

Maxima and Minima 18.3 Q2(i)

$$f(x) = (x-1)(x-2)^2$$

$$\begin{aligned} \therefore f'(x) &= (x-2)^2 + 2(x-1)(x-2) \\ &= (x-2)(x-2+2x-2) \\ &= (x-2)(3x-4) \end{aligned}$$

$$f''(x) = (3x-4) + 3(x-2)$$

For maxima and minima,

$$f'(x) = 0$$

$$\Rightarrow (x-2)(3x-4) = 0$$

$$\Rightarrow x = 2, \frac{4}{3}$$

Now,

$$f''(2) > 0$$

$$\therefore x = 2 \text{ is local minima}$$

$$f''\left(\frac{4}{3}\right) = -2 < 0$$

$$\therefore x = \frac{4}{3} \text{ is point of local maxima}$$

$$\therefore \text{local max value} = f\left(\frac{4}{3}\right) = \frac{4}{27}$$

$$\text{local min value} = f(2) = 0.$$

Maxima and Minima 18.3 Q2(ii)

$$f(x) = x\sqrt{1-x}$$

$$\begin{aligned}\therefore f'(x) &= \sqrt{1-x} + \frac{x}{2\sqrt{1-x}}(-1) \\ &= \frac{2(1-x) - x}{2\sqrt{1-x}} \\ &= \frac{2-3x}{2\sqrt{1-x}} \\ f''(x) &= \frac{2\sqrt{1-x}(-3) + \frac{(2-3x)}{\sqrt{1-x}}}{4(1-x)}\end{aligned}$$

For maximum and minimum,

$$f'(x) = 0$$

$$\Rightarrow \frac{2-3x}{2\sqrt{1-x}} = 0$$

$$\Rightarrow x = \frac{2}{3}$$

Now,

$$f''\left(\frac{2}{3}\right) < 0$$

$$\therefore x = \frac{2}{3} \text{ is point of maxima}$$

$$\therefore \text{local max value} = f\left(\frac{2}{3}\right) = \frac{2}{3\sqrt{3}}.$$

Maxima and Minima 18.3 Q2(iii)

$$f(x) = -(x-1)^3(x+1)^2$$

$$\begin{aligned}\therefore f'(x) &= -3(x-1)^2(x+1)^2 - 2(x-1)^3(x+1) \\ &= -(x-1)^2(x+1)(3x+3+2x-2) \\ &= -(x-1)^2(x+1)(5x+1)\end{aligned}$$

$$\therefore f''(x) = -2(x-1)(x+1)(5x+1) - (x-1)^2(5x+1) - 5(x-1)^2(x+1)$$

For maximum and minimum value,

$$f'(x) = 0$$

$$\Rightarrow -(x-1)^2(x+1)(5x+1) = 0$$

$$\Rightarrow x = 1, -1, -\frac{1}{5}$$

Now,

$$f''(1) = 0$$

$\therefore x = 1$ is inflection point

$$f''(-1) = -4 \times -4 = 16 > 0$$

$\therefore x = -1$ is point of minima

$$f''\left(-\frac{1}{5}\right) = -5\left(\frac{36}{25}\right) \times \frac{4}{5} = \frac{-144}{25} < 0$$

$\therefore x = -\frac{1}{5}$ is point of maxima

Hence,

$$\text{local max value} = f\left(-\frac{1}{5}\right) = \frac{3456}{3125}$$

$$\text{local min value} = f(-1) = 0.$$

Maxima and Minima 18.3 Q3

We have,

$$y = a \log x + bx^2 + x$$

$$\therefore \frac{dy}{dx} = \frac{a}{x} + 2bx + 1$$

$$\text{and } \frac{d^2y}{dx^2} = \frac{-a}{x^2} + 2b$$

For maximum and minimum value,

$$\frac{dy}{dx} = 0$$

$$\Rightarrow \frac{a}{x} + 2bx + 1 = 0$$

Given that extreme value exist at $x = 1, 2$

$$\Rightarrow a + 2b = -1 \quad \text{--- (i)}$$

$$\frac{a}{2} + 4b = -1$$

$$\Rightarrow a + 8b = -2 \quad \text{--- (ii)}$$

Solving (i) and (ii), we get

$$a = \frac{-2}{3}, b = \frac{-1}{6}.$$

Maxima and Minima 18.3 Q4

The given function is $f(x) = \frac{\log x}{x}$.

$$f'(x) = \frac{x\left(\frac{1}{x}\right) - \log x}{x^2} = \frac{1 - \log x}{x^2}$$

$$\text{Now, } f'(x) = 0$$

$$\Rightarrow 1 - \log x = 0$$

$$\Rightarrow \log x = 1$$

$$\Rightarrow \log x = \log e$$

$$\Rightarrow x = e$$

$$\begin{aligned}\text{Now, } f''(x) &= \frac{x^2\left(-\frac{1}{x}\right) - (1 - \log x)(2x)}{x^4} \\ &= \frac{-x - 2x(1 - \log x)}{x^4} \\ &= \frac{-3 + 2\log x}{x^3}\end{aligned}$$

$$\text{Now, } f''(e) = \frac{-3 + 2\log e}{e^3} = \frac{-3 + 2}{e^3} = \frac{-1}{e^3} < 0$$

Therefore, by second derivative test, f is the maximum at $x = e$.

Maxima and Minima 18.3 Q5

$$f(x) = \frac{4}{x+2} + x$$

$$\therefore f'(x) = \frac{-4}{(x+2)^2} + 1$$

$$f''(x) = \frac{8}{(x+2)^3}$$

For maximum and minimum value,

$$f'(x) = 0$$

$$\Rightarrow \frac{-4}{(x+2)^2} + 1 = 0$$

$$\Rightarrow (x+2)^2 = 4$$

$$\Rightarrow x^2 + 4x = 0$$

$$\Rightarrow x(x+4) = 0$$

$$x = 0, -4$$

Now,

$$f''(0) = 1 > 0$$

$\therefore x = 0$ is point of minima

$$f''(-4) = -1 < 0$$

$\therefore x = -4$ is point of maxima

$\therefore$ local max value $= f(-4) = -6$

local min value $= f(0) = 2$.

Maxima and Minima 18.3 Q6

We have,

$$y = \tan x - 2x$$

$$\therefore y' = \sec^2 x - 2$$

$$y'' = 2 \sec^2 x \tan x$$

For maximum and minimum value,

$$y' = 0$$

$$\Rightarrow \sec^2 x = 2$$

$$\Rightarrow \sec x = \pm\sqrt{2}$$

$$\Rightarrow x = \frac{\pi}{4}, \frac{3\pi}{4}$$

$$\therefore y''\left(\frac{\pi}{4}\right) = 4 > 0$$

$$\therefore x = \frac{\pi}{4} \text{ is point of minima}$$

$$y''\left(\frac{3\pi}{4}\right) = -4 < 0$$

$$\therefore x = \frac{3\pi}{4} \text{ is point of maxima}$$

Hence,

$$\text{max value} = f\left(\frac{3\pi}{4}\right) = -1 - \frac{3\pi}{2}$$

$$\text{min value} = f\left(\frac{\pi}{4}\right) = 1 - \frac{\pi}{2}$$

Maxima and Minima 18.3 Q7

Consider the function

$$f(x) = x^3 + ax^2 + bx + c$$

$$\text{Then } f'(x) = 3x^2 + 2ax + b$$

It is given that $f(x)$ is maximum at $x = -1$.

$$\therefore f'(-1) = 3(-1)^2 + 2a(-1) + b = 0$$

$$\Rightarrow f'(-1) = 3 - 2a + b = 0 \dots (1)$$

It is given that $f(x)$ is minimum at $x = 3$.

$$\therefore f'(3) = 3(3)^2 + 2a(3) + b = 0$$


$$\Rightarrow f'(3) = 27 + 6a + b = 0 \dots (2)$$

Solving equations (1) and (2), we have,

$$a = -3 \text{ and } b = -9$$

Since $f'(x)$ is independent of constant c , it can be any real number.

RD Sharma
Solutions
Class 12 Maths
Chapter 18
Ex 18.4



Maxima and Minima 18.4 Q1(i)

The given function is $f(x) = 4x - \frac{1}{2}x^2$.

$$\therefore f'(x) = 4 - \frac{1}{2}(2x) = 4 - x$$

Now,

$$f'(x) = 0 \Rightarrow x = 4$$

Then, we evaluate the value of f at critical point $x = 4$ and at the end points of the interval $\left[-2, \frac{9}{2}\right]$.

$$f(4) = 16 - \frac{1}{2}(16) = 16 - 8 = 8$$

$$f(-2) = -8 - \frac{1}{2}(4) = -8 - 2 = -10$$

$$f\left(\frac{9}{2}\right) = 4\left(\frac{9}{2}\right) - \frac{1}{2}\left(\frac{9}{2}\right)^2 = 18 - \frac{81}{8} = 18 - 10.125 = 7.875$$

Hence, we can conclude that the absolute maximum value of f on $\left[-2, \frac{9}{2}\right]$ is 8 occurring at $x = 4$

and the absolute minimum value of f on $\left[-2, \frac{9}{2}\right]$ is -10 occurring at $x = -2$.

Maxima and Minima 18.4 Q1(ii)

The given function is $f(x) = (x-1)^2 + 3$.

$$\therefore f'(x) = 2(x-1)$$

Now,

$$f'(x) = 0 \Rightarrow 2(x-1) = 0 \Rightarrow x = 1$$

Then, we evaluate the value of f at critical point $x = 1$ and at the end points of the interval $[-3, 1]$.

$$f(1) = (1-1)^2 + 3 = 0 + 3 = 3$$

$$f(-3) = (-3-1)^2 + 3 = 16 + 3 = 19$$

Hence, we can conclude that the absolute maximum value of f on $[-3, 1]$ is 19 occurring at $x = -3$ and the minimum value of f on $[-3, 1]$ is 3 occurring at $x = 1$.

Maxima and Minima 18.4 Q1(iii)

$$\text{Let } f(x) = 3x^4 - 8x^3 + 12x^2 - 48x + 25.$$

$$\therefore f'(x) = 12x^3 - 24x^2 + 24x - 48$$

$$= 12(x^3 - 2x^2 + 2x - 4)$$

$$= 12[x^2(x-2) + 2(x-2)]$$

$$= 12(x-2)(x^2+2)$$

Now, $f'(x) = 0$ gives $x = 2$ or $x^2 + 2 = 0$ for which there are no real roots.

Therefore, we consider only $x = 2 \in [0, 3]$.

Now, we evaluate the value of f at critical point $x = 2$ and at the end points of the interval $[0, 3]$.

$$f(2) = 3(16) - 8(8) + 12(4) - 48(2) + 25$$

$$= 48 - 64 + 48 - 96 + 25$$

$$= -39$$

$$f(0) = 3(0) - 8(0) + 12(0) - 48(0) + 25$$

$$= 25$$

$$f(3) = 3(81) - 8(27) + 12(9) - 48(3) + 25$$

$$= 243 - 216 + 108 - 144 + 25 = 16$$

Hence, we can conclude that the absolute maximum value of f on $[0, 3]$ is 25 occurring at $x = 0$ and the absolute minimum value of f at $[0, 3]$ is -39 occurring at $x = 2$.

Maxima and Minima 18.4 Q1(iv)

$$f(x) = (x-2)\sqrt{x-1}$$

$$\Rightarrow f'(x) = \sqrt{x-1} + (x-2) \frac{1}{2\sqrt{x-1}}$$

$$\text{Put } f'(x) = 0$$

$$\Rightarrow \sqrt{x-1} + \frac{x-2}{2\sqrt{x-1}} = 0$$

$$\Rightarrow \frac{2(x-1) + (x-2)}{2\sqrt{x-1}} = 0$$

$$\Rightarrow \frac{3x-4}{2\sqrt{x-1}} = 0$$

$$\Rightarrow x = \frac{4}{3}$$

Now,

$$f(1) = 0$$

$$f\left(\frac{4}{3}\right) = \left(\frac{4}{3} - 2\right)\sqrt{\frac{4}{3} - 1} = \frac{4-6}{3\sqrt{3}} = \frac{-2}{3\sqrt{3}} = \frac{-2\sqrt{3}}{9}$$

$$f(9) = (9-2)\sqrt{9-1} = 7\sqrt{8} = 14\sqrt{2}$$

$\therefore$ The absolute maximum value of $f(x)$ is $14\sqrt{2}$ at $x = 9$ and the absolute minimum value is $\frac{-2\sqrt{3}}{9}$ at $x = \frac{4}{3}$.

Maxima and Minima 18.4 Q2

$$\text{Let } f(x) = 2x^3 - 24x + 107.$$

$$\therefore f'(x) = 6x^2 - 24 = 6(x^2 - 4)$$

Now,

$$f'(x) = 0 \Rightarrow 6(x^2 - 4) = 0 \Rightarrow x^2 = 4 \Rightarrow x = \pm 2$$

We first consider the interval $[1, 3]$.

Then, we evaluate the value of f at the critical point $x = 2 \in [1, 3]$ and at the end points of the interval $[1, 3]$.

$$f(2) = 2(8) - 24(2) + 107 = 16 - 48 + 107 = 75$$

$$f(1) = 2(1) - 24(1) + 107 = 2 - 24 + 107 = 85$$

$$f(3) = 2(27) - 24(3) + 107 = 54 - 72 + 107 = 89$$

Hence, the absolute maximum value of $f(x)$ in the interval $[1, 3]$ is 89 occurring at $x = 3$.

Next, we consider the interval $[-3, -1]$.

Evaluate the value of f at the critical point $x = -2 \in [-3, -1]$ and at the end points of the interval $[-3, -1]$.

$$f(-3) = 2(-27) - 24(-3) + 107 = -54 + 72 + 107 = 125$$

Maxima and Minima 18.4 Q3

$$f(x) = \cos^2 x + \sin x$$

$$\begin{aligned} f'(x) &= 2 \cos x (-\sin x) + \cos x \\ &= -2 \sin x \cos x + \cos x \end{aligned}$$

$$\text{Now, } f'(x) = 0$$

$$\Rightarrow 2 \sin x \cos x = \cos x \Rightarrow \cos x (2 \sin x - 1) = 0$$

$$\Rightarrow \sin x = \frac{1}{2} \text{ or } \cos x = 0$$

$$\Rightarrow x = \frac{\pi}{6}, \text{ or } \frac{\pi}{2} \text{ as } x \in [0, \pi]$$

Now, evaluating the value of f at critical points $x = \frac{\pi}{6}$ and $x = \frac{\pi}{2}$ and at the end points of the interval $[0, \pi]$ (i.e., at $x = 0$ and $x = \pi$), we have:

$$f\left(\frac{\pi}{6}\right) = \cos^2 \frac{\pi}{6} + \sin \frac{\pi}{6} = \left(\frac{\sqrt{3}}{2}\right)^2 + \frac{1}{2} = \frac{5}{4}$$

$$f(0) = \cos^2 0 + \sin 0 = 1 + 0 = 1$$

$$f(\pi) = \cos^2 \pi + \sin \pi = (-1)^2 + 0 = 1$$

$$f\left(\frac{\pi}{2}\right) = \cos^2 \frac{\pi}{2} + \sin \frac{\pi}{2} = 0 + 1 = 1$$

Hence, the absolute maximum value of f is $\frac{5}{4}$ occurring at $x = \frac{\pi}{6}$ and the absolute minimum value of f is 1 occurring at $x = 0, \frac{\pi}{2}, \text{ and } \pi$.

Maxima and Minima 18.4 Q4

We have

$$f(x) = 12x^{\frac{4}{3}} - 6x^{\frac{1}{3}}$$

$$\therefore f'(x) = 16x^{\frac{1}{3}} - \frac{2}{2x^{\frac{2}{3}}} = \frac{2(8x - 1)}{2x^{\frac{2}{3}}}$$

$$\text{Thus, } f'(x) = 0$$

$$\Rightarrow x = \frac{1}{8}$$

Further note that $f'(x)$ is not defined at $x = 0$.

So, the critical points are $x = 0$ and $x = \frac{1}{8}$.

Evaluating the value of f at critical points $x = 0, \frac{1}{8}$ and at end points of the interval $x = -1$ and $x = 1$

$$f(-1) = 12(-1)^{\frac{4}{3}} - 6(-1)^{\frac{1}{3}} = 18$$

$$f(0) = 12(0) - 6(0) = 0$$

$$f\left(\frac{1}{8}\right) = 12\left(\frac{1}{8}\right)^{\frac{4}{3}} - 6\left(\frac{1}{8}\right)^{\frac{1}{3}} = -\frac{9}{4}$$

$$f(1) = 12(1)^{\frac{4}{3}} - 6(1)^{\frac{1}{3}} = 6$$

Hence we conclude that absolute maximum value of f is 18 at $x = -1$

and absolute minimum value of f is $-\frac{9}{4}$ at $x = \frac{1}{8}$.

Maxima and Minima 18.4 Q5

Given,

$$f(x) = 2x^3 - 15x^2 + 36x + 1$$

$$\therefore f'(x) = 6x^2 - 30x + 36 = 6(x^2 - 5x + 6) = 6(x - 2)(x - 3)$$

Note that $f'(x) = 0$ gives $x = 2$ and $x = 3$

We shall now evaluate the value of f at these points and at the end points of the interval $[1, 5]$,

i.e. at $x = 1, 2, 3$ and 5

$$\text{At } x = 1, f(1) = 2(1^3) - 15(1^2) + 36(1) + 1 = 24$$


$$\text{At } x = 2, f(2) = 2(2^3) - 15(2^2) + 36(2) + 1 = 29$$

$$\text{At } x = 3, f(3) = 2(3^3) - 15(3^2) + 36(3) + 1 = 28$$

$$\text{At } x = 5, f(5) = 2(5^3) - 15(5^2) + 36(5) + 1 = 56$$

Thus we conclude that the absolute maximum value of f on $[1, 5]$ is 56, occurring at $x = 5$, and absolute minimum value of f on $[1, 5]$ is 24 which occurs at $x = 1$.

RD Sharma
Solutions
Class 12 Maths
Chapter
Ex 18.5





Maxima and Minima 18.5 Q1

Let x and y be the two numbers.

$$\text{Given that } x + y = 16 \quad \text{---(i)}$$

$$\text{Let } s = x^2 + y^2 \quad \text{---(ii)}$$

From (i) and (ii)

$$s = x^2 + (15 - x)^2$$

$$\begin{aligned}\therefore \frac{ds}{dx} &= 2x + 2(15 - x)(-1) \\ &= 2x - 30 + 2x \\ &= 4x - 30\end{aligned}$$

$$\text{Now, } \frac{ds}{dx} = 0$$

$$\Rightarrow 4x - 30 = 0$$

$$\Rightarrow x = \frac{15}{2}$$

Since,

$$\frac{d^2s}{dx^2} = 4 > 0$$

$$\therefore x = \frac{15}{2} \text{ is the point of local minima.}$$

So, from (i)

$$y = 15 - \frac{15}{2} = \frac{15}{2}$$

Hence, the required numbers are $\frac{15}{2}, \frac{15}{2}$.

Maxima and Minima 18.5 Q2

Let x and y be the two parts of 64.

$$\therefore x + y = 64 \quad \text{---(i)}$$

$$\text{Let } S = x^3 + y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x^3 + (64 - x)^3$$

$$\begin{aligned}\therefore \frac{dS}{dx} &= 3x^2 + 3(64 - x)^2 \times (-1) \\ &= 3x^2 - 3(4096 - 128x + x^2) \\ &= -3(4096 - 128x)\end{aligned}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow -3(4096 - 128x) = 0$$

$$\Rightarrow x = 32$$

Now,

$$\frac{d^2S}{dx^2} = 384 > 0$$

$\therefore x = 32$ is the point of local minima.

Thus, the two parts of 64 are $(32, 32)$.

Maxima and Minima 18.5 Q3

Let x and y be the two numbers, such that, $x, y \geq -2$ and

$$x + y = \frac{1}{2} \quad \text{---(i)}$$

$$\text{Let } S = x + y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x + \left(\frac{1}{2} - x\right)^3$$

$$\begin{aligned}\therefore \frac{dS}{dx} &= 1 + 3\left(\frac{1}{2} - x\right)^2 \times (-1) \\ &= 1 - 3\left(\frac{1}{4} - x + x^2\right) \\ &= \frac{1}{4} + 3x - 3x^2\end{aligned}$$

For maximum and minimum,

$$\begin{aligned}\frac{dS}{dx} &= 0 \\ \Rightarrow \frac{1}{4} + 3x - 3x^2 &= 0 \\ \Rightarrow 1 + 12x - 12x^2 &= 0 \\ \Rightarrow 12x^2 - 12x - 1 &= 0 \\ \Rightarrow x &= \frac{12 \pm \sqrt{144 + 48}}{24} \\ \Rightarrow x &= \frac{1}{2} \pm \frac{8\sqrt{3}}{24} \\ \Rightarrow x &= \frac{1}{2} \pm \frac{1}{\sqrt{3}} \\ \Rightarrow x &= \frac{1}{2} - \frac{1}{\sqrt{3}}, \frac{1}{2} + \frac{1}{\sqrt{3}}\end{aligned}$$

Now,

$$\frac{d^2S}{dx^2} = 3 - 6x$$

$$\begin{aligned}\text{At } x &= \frac{1}{2} - \frac{1}{\sqrt{3}}, \quad \frac{d^2S}{dx^2} = 3 \left(1 - 2\left(\frac{1}{2} - \frac{1}{\sqrt{3}}\right)\right) \\ &= 3 \left(+\frac{2}{\sqrt{3}}\right) = 2\sqrt{3} > 0\end{aligned}$$

$$\therefore x = \frac{1}{2} - \frac{1}{\sqrt{3}} \text{ is point of local minima}$$

$\therefore$ from (i)

$$y = \frac{1}{2} - \left(\frac{1}{2} - \frac{1}{\sqrt{3}}\right) = \frac{1}{\sqrt{3}}$$

Hence, the required numbers are $\frac{1}{2} - \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$.

Maxima and Minima 18.5 Q4

Let x and y be the two parts of 15, such that

$$\therefore x + y = 15 \quad \text{---(i)}$$

$$\text{Also, } S = x^2 y^3 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x^2 (15 - x)^3$$

$$\begin{aligned}\therefore \frac{dS}{dx} &= 2x(15 - x)^3 - 3x^2(15 - x)^2 \\ &= (15 - x)^2 [30x - 2x^2 - 3x^2] \\ &= 5x(15 - x)^2(6 - x)\end{aligned}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 5x(15 - x)^2(6 - x) = 0$$

$$\Rightarrow x = 0, 15, 6$$

Now,

$$\frac{d^2S}{dx^2} = 5(15 - x)^2(6 - x) - 5x \times 2(15 - x)(6 - x) - 5x(15 - x)^2$$

$$\therefore \text{At } x = 0, \frac{d^2S}{dx^2} = 1125 > 0$$

$$\therefore x = 0 \text{ is point of local minima}$$

$$\text{At } x = 15, \frac{d^2S}{dx^2} = 0$$

$$\therefore x = 15 \text{ is an inflection point.}$$

$$\text{At } x = 6, \frac{d^2S}{dx^2} = -2430 < 0$$

$$\therefore x = 6 \text{ is the point of local maxima}$$

Thus the numbers are 6 and 9.

Maxima and Minima 18.5 Q5

Let r and h be the radius and height of the cylinder respectively.

Then, volume (V) of the cylinder is given by,

$$V = \pi r^2 h = 100 \quad (\text{given})$$

$$\therefore h = \frac{100}{\pi r^2}$$

Surface area (S) of the cylinder is given by,

$$S = 2\pi r^2 + 2\pi r h = 2\pi r^2 + \frac{200}{r}$$

$$\therefore \frac{dS}{dr} = 4\pi r - \frac{200}{r^2}, \quad \frac{d^2S}{dr^2} = 4\pi + \frac{400}{r^3}$$

$$\frac{dS}{dr} = 0 \Rightarrow 4\pi r = \frac{200}{r^2}$$

$$\Rightarrow r^3 = \frac{200}{4\pi} = \frac{50}{\pi}$$

$$\Rightarrow r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$$

Now, it is observed that when $r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$, $\frac{d^2S}{dr^2} > 0$.

$\therefore$ By second derivative test, the surface area is the minimum when the radius of the cylinder

is $\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ cm.

$$\text{When } r = \left(\frac{50}{\pi}\right)^{\frac{1}{3}}, \quad h = \frac{100}{\pi \left(\frac{50}{\pi}\right)^{\frac{2}{3}}} = \frac{2 \times 50}{(50)^{\frac{2}{3}} (\pi)^{1-\frac{2}{3}}} = 2 \left(\frac{50}{\pi}\right)^{\frac{1}{3}} \text{ cm.}$$

Hence, the required dimensions of the can which has the minimum surface area is given by

radius = $\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ cm and height = $2 \left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ cm.

We are given that the bending moment M at a distance x from one end of the beam is given by

$$(i) \quad M = \frac{WL}{2}x - \frac{W}{2}x^2$$

$$\therefore \quad \frac{dM}{dx} = \frac{WL}{2} - Wx$$

For maxima and minima,

$$\frac{dM}{dx} = 0 \Rightarrow \quad \frac{WL}{2} - Wx = 0 \Rightarrow \quad x = \frac{L}{2}$$

Now,

$$\frac{d^2M}{dx^2} = -W < 0$$

$$\therefore \quad x = \frac{L}{2} \text{ is point of local maxima.}$$

$$(ii) \quad M = \frac{Wx}{3} - \frac{Wx^3}{3L^2}$$

$$\therefore \quad \frac{dM}{dx} = \frac{W}{3} - \frac{Wx^2}{L^2}$$

For maxima and minima,

$$\frac{dM}{dx} = 0 \Rightarrow \quad \frac{W}{3} - \frac{Wx^2}{L^2} = 0 \Rightarrow \quad x = \frac{L}{\sqrt{3}}$$

Now,

$$\frac{d^2M}{dx^2} = -\frac{2xW}{L^2}$$

$$\text{At } x = \frac{L}{\sqrt{3}}, \quad \frac{d^2M}{dx^2} = -\frac{2W}{\sqrt{3}L} < 0$$

$$\therefore \quad x = \frac{L}{\sqrt{3}} \text{ is point of local maxima}$$

$$\Rightarrow \quad \begin{aligned} \frac{d^2s}{dx^2} &= -\frac{\sqrt{2}r}{\frac{r^2}{2}} \\ &= \frac{2\sqrt{2}}{r} < 0 \end{aligned}$$

$$\therefore \quad x = \frac{r}{\sqrt{2}} \text{ is the point of local maxima}$$

From (i)

$$y = \frac{r}{\sqrt{2}}$$

Hence, $x = \frac{r}{\sqrt{2}}, y = \frac{r}{\sqrt{2}}$ is the required number.

Let a piece of length l be cut from the given wire to make a square.

Then, the other piece of wire to be made into a circle is of length $(28 - l)$ m.

Now, side of square $= \frac{l}{4}$.

Let r be the radius of the circle. Then, $2\pi r = 28 - l \Rightarrow r = \frac{1}{2\pi}(28 - l)$.

The combined areas of the square and the circle (A) is given by,

$$\begin{aligned} A &= (\text{side of the square})^2 + r^2 \\ &= \frac{l^2}{16} + \pi \left[\frac{1}{2\pi}(28 - l) \right]^2 \\ &= \frac{l^2}{16} + \frac{1}{4\pi}(28 - l)^2 \\ \therefore \frac{dA}{dl} &= \frac{2l}{16} + \frac{2}{4\pi}(28 - l)(-1) = \frac{l}{8} - \frac{1}{2\pi}(28 - l) \\ \frac{d^2A}{dl^2} &= \frac{1}{8} + \frac{1}{2\pi} > 0 \\ \text{Now, } \frac{dA}{dl} = 0 &\Rightarrow \frac{l}{8} - \frac{1}{2\pi}(28 - l) = 0 \\ \Rightarrow \frac{\pi l - 4(28 - l)}{8\pi} &= 0 \\ \Rightarrow (\pi + 4)l - 112 &= 0 \\ \Rightarrow l &= \frac{112}{\pi + 4} \end{aligned}$$

Thus, when $l = \frac{112}{\pi + 4}$, $\frac{d^2A}{dl^2} > 0$.

$\therefore$ By second derivative test, the area (A) is the minimum when $l = \frac{112}{\pi + 4}$.

Hence, the combined area is the minimum when the length of the wire in making the square is $\frac{112}{\pi + 4}$ cm while the length of the wire in making the circle is $28 - \frac{112}{\pi + 4} = \frac{28\pi}{\pi + 4}$ cm.

Let the wire of length 20 m be cut into x cm and y cm and bent into a square and equilateral triangle, so that the sum of area of square and triangle is minimum.

Now,

$$x + y = 20 \quad \text{---(i)}$$

$$x = 4l \text{ and } y = 3a$$

Let s = sum of area of square and triangle

$$s = l^2 + \frac{\sqrt{3}}{4}a^2 \quad \text{---(ii)}$$

$$\left[\because \text{area of equilateral } \Delta = \frac{\sqrt{3}}{4}(\text{one side})^2 \right]$$

We have, $4l + 3a = 20$

$$\Rightarrow 4l = 20 - 3a$$

$$\Rightarrow l = \frac{20 - 3a}{4}$$

From (i), we have,

$$s = \left(\frac{20 - 3a}{4} \right)^2 + \frac{\sqrt{3}}{4}a^2$$

$$\frac{ds}{da} = 2 \left(\frac{20 - 3a}{4} \right) \left(\frac{-3}{4} \right) + 2a \times \frac{\sqrt{3}}{4}$$

To find the maximum or minimum, $\frac{ds}{da} = 0$

$$\Rightarrow 2 \left(\frac{20 - 3a}{4} \right) \left(\frac{-3}{4} \right) + 2a \times \frac{\sqrt{3}}{4} = 0$$

$$\Rightarrow -3(20 - 3a) + 4a\sqrt{3} = 0$$

$$\Rightarrow -60 + 9a + 4a\sqrt{3} = 0$$

$$\Rightarrow 9a + 4a\sqrt{3} = 60$$

$$\Rightarrow a(9 + 4\sqrt{3}) = 60$$

$$\Rightarrow a = \frac{60}{9 + 4\sqrt{3}}$$

Differentiating once again, we have,

$$\frac{d^2s}{da^2} = \frac{9 + 4\sqrt{3}}{8} > 0$$

Thus, the sum of the areas of the square and triangle is minimum when $a = \frac{60}{9 + 4\sqrt{3}}$

We know that, $l = \frac{20 - 3a}{4}$

$$\Rightarrow l = \frac{20 - 3 \left(\frac{60}{9 + 4\sqrt{3}} \right)}{4}$$

$$\Rightarrow l = \frac{180 + 80\sqrt{3} - 180}{4(9 + 4\sqrt{3})}$$

$$\Rightarrow l = \frac{20\sqrt{3}}{9 + 4\sqrt{3}}$$

Let r be the radius of the circle and a be the side of the square.

Then, we have:

$$2\pi r + 4a = k \text{ (where } k \text{ is constant)}$$

$$\Rightarrow a = \frac{k - 2\pi r}{4}$$

The sum of the areas of the circle and the square (A) is given by,

$$A = \pi r^2 + a^2 = \pi r^2 + \frac{(k - 2\pi r)^2}{16}$$

$$\therefore \frac{dA}{dr} = 2\pi r + \frac{2(k - 2\pi r)(-2\pi)}{16} = 2\pi r - \frac{\pi(k - 2\pi r)}{4}$$

$$\text{Now, } \frac{dA}{dr} = 0$$

$$\Rightarrow 2\pi r = \frac{\pi(k - 2\pi r)}{4}$$

$$8r = k - 2\pi r$$

$$\Rightarrow (8 + 2\pi)r = k$$

$$\Rightarrow r = \frac{k}{8 + 2\pi} = \frac{k}{2(4 + \pi)}$$

$$\text{Now, } \frac{d^2A}{dr^2} = 2\pi + \frac{\pi^2}{2} > 0$$

$$\therefore \text{ When } r = \frac{k}{2(4 + \pi)}, \frac{d^2A}{dr^2} > 0.$$

$$\therefore \text{ The sum of the areas is least when } r = \frac{k}{2(4 + \pi)}.$$

$$\text{When } r = \frac{k}{2(4 + \pi)}, a = \frac{k - 2\pi \left[\frac{k}{2(4 + \pi)} \right]}{4} = \frac{k(4 + \pi) - k}{4(4 + \pi)} = \frac{4k}{4(4 + \pi)} = \frac{k}{4 + \pi} = 2r.$$

Hence, it has been proved that the sum of their areas is least when the side of the square is double the radius of the circle.

ABC is a right angled triangle. Hypotenuse $h = AC = 5$ cm.

Let x and y one the other two side of the triangle.

$$\therefore x^2 + y^2 = 25 \quad \text{---(i)}$$

$$\therefore \text{Area of } \triangle ABC = \frac{1}{2} BC \times AB$$

$$\Rightarrow S = \frac{1}{2} xy \quad \text{---(ii)}$$

From (i) and (ii)

$$S = \frac{1}{2} x \sqrt{25 - x^2}$$

$$\begin{aligned} \therefore \frac{ds}{dx} &= \frac{1}{2} \left[\sqrt{25 - x^2} - \frac{2x^2}{2\sqrt{25 - x^2}} \right] \\ &= \frac{1}{2} \frac{[25 - x^2 - x^2]}{\sqrt{25 - x^2}} \\ &= \frac{1}{2} \left[\frac{25 - 2x^2}{\sqrt{25 - x^2}} \right] \end{aligned}$$

For maxima and minima,

$$\frac{ds}{dx} = 0$$

$$\Rightarrow \frac{1}{2} \left[\frac{25 - 2x^2}{\sqrt{25 - x^2}} \right] = 0$$

$$\Rightarrow x = 5\sqrt{2}$$

Now,

$$\frac{d^2s}{dx^2} = \frac{1}{2} \frac{\sqrt{25 - x^2} \times (-4x) + \frac{(25 - 2x^2) 2x}{2\sqrt{25 - x^2}}}{(25 - x^2)}$$

$$\text{At } x = \frac{5}{\sqrt{2}}, \frac{d^2s}{dx^2} = \frac{1}{2} \left[\frac{-\frac{25}{\sqrt{2}} \times \frac{5}{\sqrt{2}} + 0}{\frac{25}{2}} \right]$$

$$= -\frac{5}{2} < 0$$

$$\therefore x = \frac{5}{\sqrt{2}} \text{ is a point local maxima,}$$

ABC is a given triangle with $AB = a, BC = b$ and $\angle ABC = \theta$.

AD is perpendicular to BC .

$$\therefore BD = a \sin \theta$$

Now,

$$\text{Area of } \triangle ABC = \frac{1}{2} \times BC \times AD$$

$$\Rightarrow A = \frac{1}{2} b \times a \sin \theta \quad \text{---(i)}$$

$$\therefore \frac{dA}{d\theta} = \frac{1}{2} ab \cos \theta$$

For maxima and minima,

$$\frac{dA}{d\theta} = 0$$

$$\Rightarrow \frac{1}{2} ab \cos \theta = 0$$

$$\Rightarrow \cos \theta = 0$$

$$\Rightarrow \theta = \frac{\pi}{2}$$

Now,

$$\frac{d^2A}{d\theta^2} = -\frac{1}{2} ab \sin \theta$$

$$\text{At } \theta = \frac{\pi}{2}, \quad \frac{d^2A}{d\theta^2} = -\frac{1}{2} ab < 0$$

$$\therefore \theta = \frac{\pi}{2} \text{ is point of local maxima}$$

$$\therefore \text{Maximum area of } \Delta = \frac{1}{2} ab \sin \frac{\pi}{2} = \frac{1}{2} ab.$$

Let the side of the square to be cut off be x cm. Then, the length and the breadth of the box will be $(18 - 2x)$ cm each and the height of the box is x cm.

Therefore, the volume $V(x)$ of the box is given by,

$$V(x) = x(18 - 2x)^2$$

$$\begin{aligned}\therefore V'(x) &= (18 - 2x)^2 - 4x(18 - 2x) \\ &= (18 - 2x)[18 - 2x - 4x] \\ &= (18 - 2x)(18 - 6x) \\ &= 6 \times 2(9 - x)(3 - x) \\ &= 12(9 - x)(3 - x)\end{aligned}$$

$$\begin{aligned}\text{And, } V''(x) &= 12[-(9 - x) - (3 - x)] \\ &= -12(9 - x + 3 - x) \\ &= -12(12 - 2x) \\ &= -24(6 - x)\end{aligned}$$

Maximum volume is $V_{x=3} = 3 \times (18 - 2 \times 3)^2$

$$\Rightarrow V = 3 \times 12^2$$

$$\Rightarrow V = 3 \times 144$$

$$\Rightarrow V = 432 \text{ cm}^3$$

Maxima and Minima 18.5 Q13

Let the side of the square to be cut off be x cm. Then, the height of the box is x , the length is $45 - 2x$, and the breadth is $24 - 2x$.

Therefore, the volume $V(x)$ of the box is given by,

$$\begin{aligned}V(x) &= x(45 - 2x)(24 - 2x) \\ &= x(1080 - 90x - 48x + 4x^2) \\ &= 4x^3 - 138x^2 + 1080x \\ \therefore V'(x) &= 12x^2 - 276x + 1080 \\ &= 12(x^2 - 23x + 90) \\ &= 12(x - 18)(x - 5) \\ V''(x) &= 24x - 276 = 12(2x - 23)\end{aligned}$$

$$\text{Now, } V'(x) = 0 \Rightarrow x = 18 \text{ and } x = 5$$

It is not possible to cut off a square of side 18 cm from each corner of the rectangular sheet.

Thus, x cannot be equal to 18.

$$\therefore x = 5$$

$$\text{Now, } V''(5) = 12(10 - 23) = 12(-13) = -156 < 0$$

$\therefore$ By second derivative test, $x = 5$ is the point of maxima.

Hence, the side of the square to be cut off to make the volume of the box maximum possible is 5 cm.

Maxima and Minima 18.5 Q14

Let l , b , and h represent the length, breadth, and height of the tank respectively.

Then, we have height $(h) = 2$ m

$$\text{Volume of the tank} = 8\text{m}^3$$

$$\text{Volume of the tank} = l \times b \times h$$

$$\therefore 8 = l \times b \times 2$$

$$\Rightarrow lb = 4 \Rightarrow b = \frac{4}{l}$$

$$\text{Now, area of the base} = lb = 4$$

$$\text{Area of the 4 walls } (A) = 2h(l + b)$$

$$\therefore A = 4\left(l + \frac{4}{l}\right)$$

$$\Rightarrow \frac{dA}{dl} = 4\left(1 - \frac{4}{l^2}\right)$$

$$\text{Now, } \frac{dA}{dl} = 0$$

$$\Rightarrow 1 - \frac{4}{l^2} = 0$$

$$\Rightarrow l^2 = 4$$

$$\Rightarrow l = \pm 2$$

However, the length cannot be negative.

Therefore, we have $l = 4$.

$$\therefore b = \frac{4}{l} = \frac{4}{2} = 2$$

$$\text{Now, } \frac{d^2 A}{dl^2} = \frac{32}{l^3}$$

$$\text{When } l = 2, \frac{d^2 A}{dl^2} = \frac{32}{8} = 4 > 0.$$

Thus, by second derivative test, the area is the minimum when $l = 2$.

We have $l = b = h = 2$.

$$\therefore \text{Cost of building the base} = \text{Rs } 70 \times (lb) = \text{Rs } 70 (4) = \text{Rs } 280$$


$$\text{Cost of building the walls} = \text{Rs } 2h(l + b) \times 45 = \text{Rs } 90 (2) (2 + 2)$$

$$= \text{Rs } 8 (90) = \text{Rs } 720$$

$$\text{Required total cost} = \text{Rs } (280 + 720) = \text{Rs } 1000$$

Hence, the total cost of the tank will be Rs 1000.

Radius of the semicircular opening = $\frac{x}{2}$



It is given that the perimeter of the window is 10 m.

$$\therefore x + 2y + \frac{\pi x}{2} = 10$$

$$\Rightarrow x \left(1 + \frac{\pi}{2} \right) + 2y = 10$$

$$\Rightarrow 2y = 10 - x \left(1 + \frac{\pi}{2} \right)$$

$$\Rightarrow y = 5 - x \left(\frac{1}{2} + \frac{\pi}{4} \right)$$

$\therefore$ Area of the window (A) is given by,

$$A = xy + \frac{\pi}{2} \left(\frac{x}{2} \right)^2$$

$$= x \left[5 - x \left(\frac{1}{2} + \frac{\pi}{4} \right) \right] + \frac{\pi}{8} x^2$$

$$= 5x - x^2 \left(\frac{1}{2} + \frac{\pi}{4} \right) + \frac{\pi}{8} x^2$$

$$\therefore \frac{dA}{dx} = 5 - 2x \left(\frac{1}{2} + \frac{\pi}{4} \right) + \frac{\pi}{4} x$$

$$= 5 - x \left(1 + \frac{\pi}{2} \right) + \frac{\pi}{4} x$$

$$\therefore \frac{d^2 A}{dx^2} = - \left(1 + \frac{\pi}{2} \right) + \frac{\pi}{4} = -1 - \frac{\pi}{4}$$

$$\text{Now, } \frac{dA}{dx} = 0$$

$$\Rightarrow 5 - x \left(1 + \frac{\pi}{2} \right) + \frac{\pi}{4} x = 0$$

$$\Rightarrow 5 - x - \frac{\pi}{4} x = 0$$

$$\Rightarrow x \left(1 + \frac{\pi}{4} \right) = 5$$

$$\Rightarrow x = \frac{5}{\left(1 + \frac{\pi}{4} \right)} = \frac{20}{\pi + 4}$$

Thus, when $x = \frac{20}{\pi + 4}$ then $\frac{d^2 A}{dx^2} < 0$.


Therefore, by second derivative test, the area is the maximum when length $x = \frac{20}{\pi + 4}$ m.

Now,

$$y = 5 - \frac{20}{\pi + 4} \left(\frac{2 + \pi}{4} \right) = 5 - \frac{5(2 + \pi)}{\pi + 4} = \frac{10}{\pi + 4} \text{ m}$$

Hence, the required dimensions of the window to admit maximum light is given

by length $= \frac{20}{\pi + 4}$ m and breadth $= \frac{10}{\pi + 4}$ m.



The perimeter of the window = 12 m

$$\Rightarrow (l + 2b) + (l + l) = 12$$

$$\Rightarrow 3l + 2b = 12 \quad \text{----- (i)}$$

Let S = Area of the rectangle + Area of the equilateral Δ

From (i),

$$S = l \left(\frac{12 - 3l}{2} \right) + \frac{\sqrt{3}}{4} l^2$$

$$\therefore \frac{dS}{dl} = 6 - 3l + \frac{\sqrt{3}}{2} l = 6 - \sqrt{3} \left(\sqrt{3} - \frac{1}{2} \right) l$$

For maxima and minima,

$$\frac{dS}{dl} = 0$$

$$\Rightarrow 6 - \sqrt{3} \left(\sqrt{3} - \frac{1}{2} \right) l = 0$$

$$\Rightarrow l = \frac{6}{\sqrt{3} \left(\sqrt{3} - \frac{1}{2} \right)} = \frac{12}{6 - \sqrt{3}}$$

$$\text{Now, } \frac{d^2S}{dl^2} = -\sqrt{3} \left(\sqrt{3} - \frac{1}{2} \right) = -3 + \frac{\sqrt{3}}{2} < 0$$


$$\therefore l = \frac{12}{6 - \sqrt{3}} \text{ is the point of local maxima}$$

From (i),

$$b = \frac{12 - 3l}{2} = \frac{12 - 3 \left(\frac{12}{6 - \sqrt{3}} \right)}{2} = \frac{24 - 6\sqrt{3}}{6 - \sqrt{3}}$$

A sphere of fixed radius (R) is given.

Let r and h be the radius and the height of the cylinder respectively.



From the given figure, we have $h = 2\sqrt{R^2 - r^2}$.

The volume (V) of the cylinder is given by,

$$\begin{aligned} V &= \pi r^2 h = 2\pi r^2 \sqrt{R^2 - r^2} \\ \therefore \frac{dV}{dr} &= 4\pi r \sqrt{R^2 - r^2} + \frac{2\pi r^2 (-2r)}{2\sqrt{R^2 - r^2}} \\ &= 4\pi r \sqrt{R^2 - r^2} - \frac{2\pi r^3}{\sqrt{R^2 - r^2}} \\ &= \frac{4\pi r (R^2 - r^2) - 2\pi r^3}{\sqrt{R^2 - r^2}} \\ &= \frac{4\pi r R^2 - 6\pi r^3}{\sqrt{R^2 - r^2}} \end{aligned}$$

$$\text{Now, } \frac{dV}{dr} = 0 \Rightarrow 4\pi r R^2 - 6\pi r^3 = 0$$

$$\Rightarrow r^2 = \frac{2R^2}{3}$$

$$\begin{aligned}\text{Now, } \frac{d^2V}{dr^2} &= \frac{\sqrt{R^2 - r^2} (4\pi R^2 - 18\pi r^2) - (4\pi r R^2 - 6\pi r^3) \frac{(-2r)}{2\sqrt{R^2 - r^2}}}{(R^2 - r^2)} \\ &= \frac{(R^2 - r^2)(4\pi R^2 - 18\pi r^2) + r(4\pi r R^2 - 6\pi r^3)}{(R^2 - r^2)^{\frac{3}{2}}} \\ &= \frac{4\pi R^4 - 22\pi r^2 R^2 + 12\pi r^4 + 4\pi r^2 R^2}{(R^2 - r^2)^{\frac{3}{2}}}\end{aligned}$$

Now, it can be observed that at $r^2 = \frac{2R^2}{3}$, $\frac{d^2V}{dr^2} < 0$.


$\therefore$ The volume is the maximum when $r^2 = \frac{2R^2}{3}$.

When $r^2 = \frac{2R^2}{3}$, the height of the cylinder is $2\sqrt{R^2 - \frac{2R^2}{3}} = 2\sqrt{\frac{R^2}{3}} = \frac{2R}{\sqrt{3}}$.

Hence, the volume of the cylinder is the maximum when the height of the cylinder is $\frac{2R}{\sqrt{3}}$.

Maxima and Minima 18.5 Q18

Let $EFGH$ be a rectangle inscribed in a semi-circle with radius r .



Let l and b are the length and width of rectangle.

In $\triangle OHE$

$$HE^2 = OE^2 - OH^2$$

$$\Rightarrow HE = b = \sqrt{r^2 - \left(\frac{l}{2}\right)^2} \quad \text{---(i)}$$

Let S = Area of rectangle

$$= lb = l \times \sqrt{r^2 - \left(\frac{l}{2}\right)^2}$$

$$\therefore S = \frac{1}{2} l \sqrt{4r^2 - l^2}$$

$$\begin{aligned} \therefore \frac{ds}{dl} &= \frac{1}{2} \left[\sqrt{4r^2 - l^2} - \frac{l^2}{\sqrt{4r^2 - l^2}} \right] \\ &= \frac{1}{2} \left[\frac{4r^2 - l^2 - l^2}{\sqrt{4r^2 - l^2}} \right] \\ &= \frac{2r^2 - l^2}{\sqrt{4r^2 - l^2}} \end{aligned}$$

For maxima and minima,

$$\begin{aligned} \frac{ds}{dl} &= 0 \\ \Rightarrow \frac{2r^2 - l^2}{\sqrt{4r^2 - l^2}} &= 0 \\ \Rightarrow l &= \pm \sqrt{2}r \end{aligned}$$

Also,

$$\frac{d^2s}{dl^2} = 0 \text{ at } l = \sqrt{2}r$$

So, the dimension of the rectangle

$$l = \sqrt{2}r, \quad b = \sqrt{r^2 - \left(\frac{l}{2}\right)^2} = \frac{r}{\sqrt{2}}$$

$$\begin{aligned} \text{Area of rectangle} = lb &= \sqrt{2}r \times \frac{r}{\sqrt{2}} \\ &= r^2. \end{aligned}$$

Let r and h be the radius and the height (altitude) of the cone respectively.

Then, the volume (V) of the cone is given as:

$$V = \frac{1}{3}\pi r^2 h \Rightarrow h = \frac{3V}{r^2}$$

The surface area (S) of the cone is given by,

$S = \pi r l$ (where l is the slant height)

$$\begin{aligned} &= \pi r \sqrt{r^2 + h^2} \\ &= \pi r \sqrt{r^2 + \frac{9V^2}{r^4}} = \frac{\pi \sqrt{9r^6 + V^2}}{r^2} \\ &= \frac{1}{r} \sqrt{\pi^2 r^6 + 9V^2} \end{aligned}$$

$$\begin{aligned} \therefore \frac{dS}{dr} &= \frac{r \cdot \frac{6\pi^2 r^5}{2\sqrt{\pi^2 r^6 + 9V^2}} - \sqrt{\pi^2 r^6 + 9V^2}}{r^2} \\ &= \frac{3\pi^2 r^6 - \pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \\ &= \frac{2\pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \end{aligned}$$

$$\begin{aligned} &= \frac{2\pi^2 r^6 - 9V^2}{r^2 \sqrt{\pi^2 r^6 + 9V^2}} \\ \text{Now, } \frac{dS}{dr} &= 0 \Rightarrow 2\pi^2 r^6 = 9V^2 \Rightarrow r^6 = \frac{9V^2}{2\pi^2} \end{aligned}$$

Thus, it can be easily verified that when $r^6 = \frac{9V^2}{2\pi^2}$, $\frac{d^2S}{dr^2} > 0$.


$\therefore$ By second derivative test, the surface area of the cone is the least when $r^6 = \frac{9V^2}{2\pi^2}$.

$$\text{When } r^6 = \frac{9V^2}{2\pi^2}, h = \frac{3V}{\pi r^2} = \frac{3}{\pi r^2} \left(\frac{2\pi^2 r^6}{9} \right)^{\frac{1}{2}} = \frac{3}{\pi r^2} \cdot \frac{\sqrt{2}\pi r^3}{3} = \sqrt{2}r.$$

Hence, for a given volume, the right circular cone of the least curved surface has an altitude equal to $\sqrt{2}$ times the radius of the base.

We have a cone, which is inscribed in a sphere.

Let v be the volume of greatest cone ABC . It is obvious that, for maximum volume the axis of the cone must be along the diameter of sphere.



Let $OD = x$ and $AO = OB = R$

$$\Rightarrow BD = \sqrt{R^2 - x^2} \text{ and } AD = R + x$$

Now,

$$\begin{aligned} v &= \frac{1}{3} \pi r^2 h \\ &= \frac{1}{3} \pi BD^2 \times AD \\ &= \frac{1}{3} \pi (R^2 - x^2) \times (R + x) \end{aligned}$$

$$\begin{aligned} \therefore \frac{dv}{dx} &= \frac{\pi}{3} [-2x(R + x) + R^2 - x^2] \\ &= \frac{\pi}{3} [R^2 - 2xR - 3x^2] \end{aligned}$$

For maximum and minimum

$$\begin{aligned} \frac{dv}{dx} &= 0 \\ \Rightarrow \frac{\pi}{3} [R^2 - 2xR - 3x^2] &= 0 \\ \Rightarrow \frac{\pi}{3} [(R - 3x)(R + x)] &= 0 \\ \Rightarrow R - 3x = 0 \text{ or } x = -R \\ \Rightarrow x = \frac{R}{3} \end{aligned}$$

$\left[\because x = -R \text{ is not possible as, } x = -R \text{ will make the altitude } 0 \right]$

Now,

$$\begin{aligned} \frac{d^2v}{dx^2} &= \frac{\pi}{3} [-2R - 6x] \\ \text{At } x = \frac{R}{3}, \frac{d^2v}{dx^2} &= \frac{\pi}{3} [-2R - 2R] \\ &= \frac{-4\pi R}{3} < 0 \end{aligned}$$

$\therefore x = \frac{R}{3}$ is the point of local maxima.

$$\text{Volume of the cone} = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow V = \frac{1}{3} \pi r^2 h$$

Squaring both the sides, we have,

$$\begin{aligned} V^2 &= \left(\frac{1}{3} \pi r^2 h \right)^2 \\ &= \frac{1}{9} \pi^2 r^4 h^2 \dots (1) \end{aligned}$$

$$\Rightarrow \pi^2 r^4 h^2 = \frac{9V^2}{r^2} \dots (2)$$

Consider the curved surface area of the cone.

Thus,

$$C = \pi r l$$

Squaring both the sides, we have,


$$C^2 = \pi^2 r^2 l^2$$


We know that $l^2 = r^2 + h^2$

$$\Rightarrow C^2 = \pi^2 r^2 (r^2 + h^2)$$

$$\Rightarrow C^2 = \pi^2 r^4 + \pi^2 r^2 h^2$$

$$\Rightarrow C^2 = \pi^2 r^4 + \frac{9V^2}{r^2} \dots (\text{From equation (2)})$$





ABC is an isosceles triangle such that $AB = AC$.
 The vertical angle $\angle BAC = 2\theta$
 Triangle is inscribed in the circle with center O and radius a.

Draw AM perpendicular to BC.
 $\therefore \Delta ABC$ is an isosceles triangle the circumcentre of the circle will lie on the perpendicular from A to BC.

Let O be the circumcentre.
 $\angle BOC = 2 \times 2\theta = 4\theta$ [Using central angle theorem]
 $\angle COM = 2\theta$ [$\therefore \Delta OMB$ and ΔOMC are congruent triangles]
 $OA = OB = OC = a$ [Radius of the circle]

In ΔOMC ,
 $CM = a \sin 2\theta$ and $OM = a \cos 2\theta$
 $BC = 2CM$...[Perpendicular from the center bisects the chord]
 $BC = 2a \sin 2\theta$ (1)
 Height of $\Delta ABC = AM = AO + OM$
 $AM = a + a \cos 2\theta$ (2)

Area of ΔABC is,

$$A = \frac{1}{2} \times BC \times AM$$

Differentiating equation (3) with respect to θ

$$\frac{dA}{d\theta} = a^2 \left(2 \cos 2\theta + \frac{1}{2} \times 4 \cos 4\theta \right)$$

$$\frac{dA}{d\theta} = 2a^2 (\cos 2\theta + \cos 4\theta)$$

Differentiating again with respect to θ

$$\frac{d^2A}{d\theta^2} = 2a^2 (-2 \sin 2\theta - 4 \sin 4\theta)$$

For maximum value of area equating $\frac{dA}{d\theta} = 0$

$$2a^2 (\cos 2\theta + \cos 4\theta) = 0$$

$$\cos 2\theta + \cos 4\theta = 0$$

$$\cos 2\theta + 2 \cos^2 2\theta - 1 = 0$$

$$(2 \cos 2\theta - 1)(2 \cos 2\theta + 1) = 0$$

$$\cos 2\theta = \frac{1}{2} \text{ or } \cos 2\theta = -1$$

$$2\theta = \frac{\pi}{3} \text{ or } 2\theta = \pi$$

$$\theta = \frac{\pi}{6} \text{ or } \theta = \frac{\pi}{2}$$

If $2\theta = \pi$ it will not form a triangle.

$$\therefore \theta = \frac{\pi}{6}$$

Also $\frac{d^2A}{d\theta^2}$ is negative for $\theta = \frac{\pi}{6}$.

Thus the area of the triangle is maximum when $\theta = \frac{\pi}{6}$.

Maxima

Here, $ABCD$ is a rectangle with width $AB = x$ cm and length $AD = y$ cm.

The rectangle is rotated about AD . Let v be the volume of the cylinder so formed.

$$\therefore v = \pi r^2 y \quad \text{---(i)}$$

Again,

$$\text{Perimeter of } ABCD = 2(l + b) = 2(x + y) \quad \text{---(ii)}$$

$$\Rightarrow 36 = 2(x + y)$$

$$\Rightarrow y = 18 - x \quad \text{---(iii)}$$

From (i) and (ii), we get

$$v = \pi r^2 (18 - x) = \pi (18x^2 - x^3)$$

$$\Rightarrow \frac{dv}{dx} = \pi (36x - 3x^2)$$

For maxima or minima, we have,

$$\frac{dv}{dx} = 0$$

$$\Rightarrow \pi (36x - 3x^2) = 0$$

$$\Rightarrow 3\pi (12x - x^2) = 0$$

$$\Rightarrow x(12 - x) = 0$$

$$\Rightarrow x = 0 \text{ (Not possible) or } 12$$

$$\therefore x = 12 \text{ cm}$$

From (iii)

$$y = 18 - 12 = 6 \text{ cm}$$

Now,

$$\frac{d^2v}{dx^2} = \pi (36 - 6x)$$

$$\text{At } (x = 12, y = 6) \frac{d^2v}{dx^2} = \pi (36 - 72) = -36\pi < 0$$


$$\therefore (x = 12, y = 6) \text{ is the point of local maxima,}$$

Hence,

The dimension of rectangle, which wiout maximum value, when revolved about one of its side is width = 12 cm and length = 6 cm.

Maxima and Minima 18.5 Q24

Let r and h be the radius of the base of cone and height of the cone respectively.



Let $OD = x$

It is obvious that the axis of cone must be along the diameter of sphere for maximum volume of cone.

Now,

$$\begin{aligned}\text{In } \triangle BOD, BD &= \sqrt{R^2 - x^2} \\ &= \sqrt{144 - x^2}\end{aligned}$$

$$AD = AO + OD = R + x = 12 + x$$

$$v = \text{volume of cone} = \frac{1}{3} \pi r^2 h$$

$$\begin{aligned}\Rightarrow v &= \frac{1}{3} \pi BD^2 \times AD \\ &= \frac{1}{3} \pi (144 - x^2) (12 + x) \\ &= \frac{1}{3} \pi (1728 + 144x - 12x^2 - x^3)\end{aligned}$$

$$\therefore \frac{dv}{dx} = \frac{1}{3} \pi (144 - 24x - 3x^2)$$

For maximum and minimum of v ,

$$\frac{dv}{dx} = 0$$

$$\Rightarrow \frac{1}{3} \pi (144 - 24x - 3x^2) = 0$$

$$\Rightarrow x = -12, 4$$

$x = -12$ is not possible

$$\therefore x = 4$$

Now,

$$\frac{d^2v}{dx^2} = \frac{\pi}{3} (-24 - 6x)$$

$$\begin{aligned}\text{At } x = 4, \frac{d^2v}{dx^2} &= -2\pi (4 + x) \\ &= -2\pi \times 8 = -16\pi < 0\end{aligned}$$

$\therefore x = 4$ is point of local maxima.

Hence,

$$\begin{aligned}\text{Height of cone of maximum volume} &= R + x \\ &= 12 + 4 \\ &= 16 \text{ cm.}\end{aligned}$$

We have, a closed cylinder whose volume $v = 2156 \text{ cm}^3$

Let r and h be the radius and the height of the cylinder. Then,

$$\therefore v = \pi r^2 h = 2156 \quad \text{---(i)}$$

$$\text{Total surface area} = S = 2\pi rh + 2\pi r^2$$

$$\Rightarrow S = 2\pi r(h + r) \quad \text{---(ii)}$$

From (i) and (ii)

$$S = \frac{2156 \times 2}{r} + 2\pi r^2$$

$$\therefore \frac{ds}{dr} = -\frac{4312}{r^2} + 4\pi r$$

For maximum and minimum

$$\frac{ds}{dr} = 0$$

$$\Rightarrow \frac{-4312 + 4\pi r^3}{r^2} = 0$$

$$\Rightarrow r^3 = \frac{4312}{4\pi}$$

$$\Rightarrow r = 7$$

Now,

$$\frac{d^2s}{dr^2} = \frac{8624}{r^3} + 4\pi > 0 \text{ for } r = 7.$$

$\therefore r = 7$ is the point of local minima


Hence,

The total surface area of closed cylinder will be minimum at $r = 7 \text{ cm}$.

Maxima and Minima 18.5 Q26

Let r be the radius of the base of the cylinder and h be the height of the cylinder.

$$\therefore LM = h.$$



Let $R = 5\sqrt{3}$ cm be the radius of the sphere.

It is obvious, that for maximum volume of cylinder $ABCD$, the axis of cylinder must be along the diameter of sphere.

Let $OL = x$

$\therefore h = 2x$

Now,

$$\begin{aligned}\text{In } \triangle AOL, AL &= \sqrt{AO^2 - OL^2} \\ &= \sqrt{75 - x^2}\end{aligned}$$

Now,

$$v = \text{volume of cylinder} = \pi r^2 h$$

$$\begin{aligned}\Rightarrow v &= \pi AL^2 \times ML \\ &= \pi (75 - x^2) \times 2x\end{aligned}$$

For maxima and minima of v , we must have,

$$\begin{aligned}\frac{dv}{dx} &= \pi [150 - 6x^2] = 0 \\ \Rightarrow x &= 5 \text{ cm}\end{aligned}$$

$$\text{Also, } \frac{d^2v}{dx^2} = -12\pi x$$

$$\text{At } x = 5, \frac{d^2v}{dx^2} = -60\pi x < 0$$

$\therefore x = 5$ is point of local maxima.

Hence,

$$\text{The maximum volume of cylinder is} = \pi (75 - 25) \times 10 = 500\pi \text{ cm}^3.$$

Maxima and Minima 18.5 Q27

Let x and y be two positive numbers with

$$x^2 + y^2 = r^2 \quad \text{--- (i)}$$

$$\text{Let } S = x + y \quad \text{--- (ii)}$$

$$\therefore S = x + \sqrt{r^2 - x^2} \quad \text{from (ii)}$$

$$\therefore \frac{dS}{dx} = 1 - \frac{x}{\sqrt{r^2 - x^2}}$$

For maxima and minima,

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 1 - \frac{x}{\sqrt{r^2 - x^2}} = 0$$

$$\Rightarrow x = \sqrt{r^2 - x^2}$$

$$\Rightarrow 2x^2 = r^2$$

$$\Rightarrow x = \frac{r}{\sqrt{2}}, \frac{r}{\sqrt{2}}$$

$\therefore x$ & y are positive numbers

$$\therefore x = \frac{r}{\sqrt{2}}$$

$$\text{Also, } \frac{d^2S}{dx^2} = \frac{-\left(\sqrt{r^2 - x^2} + \frac{x^2}{\sqrt{r^2 - x^2}}\right)}{r^2 - x^2}$$

$$\text{At, } x = \frac{r}{\sqrt{2}}, \frac{d^2S}{dx^2} = - \left[\frac{\frac{r^2}{\sqrt{2}} + \frac{\frac{r^2}{2}}{\frac{r}{\sqrt{2}}}}{\frac{r^2}{2}} \right] < 0$$

Since $\frac{d^2S}{dx^2} < 0$, the sum is largest when $x = y = \frac{r}{\sqrt{2}}$

The given equation of parabola is

$$x^2 = 4y \quad \text{---(i)}$$

Let $P(x, y)$ be the nearest point on (i) from the point $A(0, 5)$

Let S be the square of the distance of P from A .

$$\therefore S = x^2 + (y - 5)^2 \quad \text{---(ii)}$$

From (i),

$$S = 4y + (y - 5)^2$$

$$\Rightarrow \frac{dS}{dy} = 4 + 2(y - 5)$$

For maxima or minima, we have

$$\frac{dS}{dy} = 0$$

$$\Rightarrow 4 + 2(y - 5) = 0$$

$$\Rightarrow 2y = 6$$

$$\Rightarrow y = 3$$

From (i)

$$x^2 = 12$$

$$\therefore x = \pm 2\sqrt{3}$$

$$\Rightarrow P = (2\sqrt{3}, 3) \text{ and } P' = (-2\sqrt{3}, 3)$$

Now,

$$\frac{d^2S}{dy^2} = 2 > 0$$

$\therefore P$ and P' are the point of local minima

Hence, the nearest points are $P(2\sqrt{3}, 3)$ and $P'(-2\sqrt{3}, 3)$.

Let $P(x, y)$ be a point on
 $y^2 = 4x$ ---(i)

Let S be the square of the distance between $A(2, -8)$ and P .

$\therefore S = (x - 2)^2 + (y + 8)^2$ ---(ii)

Using (i),

$$S = \left(\frac{y^2}{4} - 2\right)^2 + (y + 8)^2$$

$$\therefore \frac{dS}{dy} = 2\left(\frac{y^2}{4} - 2\right) \times \frac{y}{2} + 2(y + 8)$$

$$= \frac{y^3 - 8y}{4} + 2y + 16$$

$$= \frac{y^3}{4} + 16$$

For maxima and minima,

$$\frac{dS}{dy} = 0$$

$$\Rightarrow \frac{y^3}{4} + 16 = 0$$

$$\Rightarrow y = -4$$

Now,

$$\frac{d^2S}{dy^2} = \frac{3y^2}{4}$$

At $y = -4$, $\frac{d^2S}{dy^2} = 12 > 0$

$\therefore y = -4$ is the point of local minima

From (i)

$$x = \frac{y^2}{4} = 4$$

Thus, the required point is $(4, -4)$ nearest to $(2, -8)$.

Let $P(x, y)$ be a point on the curve,
 $x^2 = 8y$ ---(i)

Let $A = (2, 4)$ be a point and

let $S =$ square of the distance between P and A

$$\therefore S = (x - 2)^2 + (y - 4)^2 \quad \text{---(ii)}$$

Using (i), we get

$$S = (x - 2)^2 + \left(\frac{x^2}{8} - 4\right)^2$$

$$\begin{aligned} \therefore \frac{dS}{dy} &= 2(x - 2) + 2\left(\frac{x^2}{8} - 4\right) \times \frac{2x}{8} \\ &= 2(x - 2) + \frac{(x^2 - 32)x}{16} \end{aligned}$$

$$\begin{aligned} \text{Also, } \frac{d^2S}{dx^2} &= 2 + \frac{1}{16}[x^2 - 32 + 2x^2] \\ &= 2 + \frac{1}{16}[3x^2 - 32] \end{aligned}$$

For maxima and minima,

$$\begin{aligned} \frac{dS}{dx} &= 0 \\ \Rightarrow 2(x - 2) + \frac{x(x^2 - 32)}{16} &= 0 \\ \Rightarrow 32x - 64 + x^3 - 32x &= 0 \\ \Rightarrow x^3 - 64 &= 0 \\ \Rightarrow x &= 4 \end{aligned}$$

Now,

$$\text{At } x = 4, \frac{d^2S}{dx^2} = 2 + \frac{1}{16}[16 \times 3 - 32] = 2 + 1 = 3 > 0$$

$\therefore x = 4$ is point of local minima

From (i)

$$y = \frac{x^2}{8} = 2$$

Thus, $P(4, 2)$ is the nearest point.

Let $P(x, y)$ be a point on the curve $x^2 = 2y$ which is closest to $A(0, 5)$

Let S = square of the length of AP

$$\Rightarrow S = x^2 + (y - 5)^2 \quad \text{---(ii)}$$

Using (i),

$$S = 2y + (y - 5)^2$$

$$\therefore \frac{dS}{dy} = 2 + 2(y - 5)$$

For maxima and minima,

$$\frac{dS}{dy} = 0$$

$$\Rightarrow 2 + 2y - 10 = 0$$

$$\Rightarrow y = 4$$

Now,

$$\frac{d^2S}{dy^2} = 2 > 0$$

$\therefore y = 4$ is the point of local minima

From (i)

$$r = \pm 2\sqrt{2}$$

Hence, $(\pm 2\sqrt{2}, 4)$ is the closest point on the curve to $A(0, 5)$.

The given equations are

$$y = x^2 + 7x + 2 \quad \text{---(i)}$$

$$\text{and } y = 3x - 3 \quad \text{---(ii)}$$

Let $P(x, y)$ be the point on parabola (i) which is closest to the line (ii)

Let S be the perpendicular distance from P to the line (ii).

$$\begin{aligned} \therefore S &= \frac{|y - 3x + 3|}{\sqrt{1^2 + (-3)^2}} \\ \Rightarrow S &= \frac{|x^2 + 7x + 2 - 3x + 3|}{\sqrt{10}} \quad \text{---(iii)} \\ \Rightarrow \frac{dS}{dx} &= \frac{2x + 4}{\sqrt{10}} \end{aligned}$$

For maxima or minima, we have

$$\begin{aligned} \frac{dS}{dx} &= 0 \\ \Rightarrow \frac{2x + 4}{\sqrt{10}} &= 0 \\ \Rightarrow x &= -2 \end{aligned}$$

From (i)

$$y = 4 - 14 + 2 = -8$$

Now,

$$\frac{d^2S}{dx^2} = \frac{2}{\sqrt{10}} > 0$$

$\therefore (x = -2, y = -8)$ is the point of local minima,

Hence,

The closest point on the parabola to the line $y = 3x - 3$ is $(-2, -8)$.

Let $P(x, y)$ be a point on the curve $y^2 = 2x$ which is minimum distance from the point $A(1, 4)$.

Let

S = square of the length of AP

$$S = (x - 1)^2 + (y - 4)^2$$

Using this equation, we have

$$S = x^2 + 1 - 2x + y^2 + 16 - 8y$$

$$S = x^2 - 2x + 2x + 17 - 8y$$

$$S = \frac{y^4}{4} - 8y + 17 \quad \left[\text{Since } x = \frac{y^2}{2} \right]$$

$$\frac{dS}{dy} = y^3 - 8$$

For maxima and minima, we have

$$\frac{dS}{dy} = 0$$

$$y^3 - 8 = 0$$

$$y^3 = 8$$

$$y = 2$$

Now,

$$\frac{d^2S}{dy^2} = 3y^2$$

$$\frac{d^2S}{dy^2} = 12 > 0$$

$\therefore y = 2$ is minimum point

We have

$$x = \frac{y^2}{2}$$

$$= \frac{4}{2}$$

$$= 2$$

Hence, $(2, 2)$ is at a minimum distance from the point $(1, 4)$.

The given equation of curve is

$$y = x^3 + 3x^2 + 2x - 27 \quad \text{--- (i)}$$

Slope of (i)

$$m = \frac{dy}{dx} = -3x^2 + 6x + 2 \quad \text{--- (ii)}$$

Now,

$$\frac{dm}{dx} = -6x + 6$$

and $\frac{d^2m}{dx^2} = -6 < 0$

For maxima and minima,

$$\frac{dm}{dx} = 0$$

$$\Rightarrow -6x + 6 = 0$$

$$\Rightarrow x = 1$$

$$\therefore \frac{d^2m}{dx^2} = -6 < 0$$

$\therefore x = 1$ is point of local maxima

Hence, maximum slope = $-3 + 6 + 2 = 5$

Maxima and Minima 18.5 Q35

We have,

Cost of producing x radio sets is Rs. $\frac{x^2}{4} + 35x + 25$

Selling price of x radio is Rs. $x \left(50 - \frac{x}{2} \right)$

So,

Profit on x radio sets is

$$P = \text{Rs} \left(50x - \frac{x^2}{2} - \frac{x^2}{4} - 35x - 25 \right)$$

$$\begin{aligned} \therefore \frac{dP}{dx} &= 50 - x - \frac{x}{2} - 35 \\ &= 15 - \frac{3}{2}x \end{aligned}$$

For maxima and minima,

$$\frac{dP}{dx} = 0$$

$$\Rightarrow 15 - \frac{3}{2}x = 0$$

$$\Rightarrow x = 10$$

Also,

$$\frac{d^2P}{dx^2} = \frac{-3}{2} < 0$$

$\therefore x = 10$ is the point of local maxima

Hence, the daily output should be 10 radio sets.

Maxima and Minima 18.5 Q35

We have,

Cost of producing x radio sets is Rs. $\frac{x^2}{4} + 35x + 25$

Selling price of x radio is Rs. $x \left(50 - \frac{x}{2} \right)$

So,

Profit on x radio sets is

$$P = \text{Rs} \left(50x - \frac{x^2}{2} - \frac{x^2}{4} - 35x - 25 \right)$$

$$\begin{aligned}\therefore \frac{dP}{dx} &= 50 - x - \frac{x}{2} - 35 \\ &= 15 - \frac{3}{2}x\end{aligned}$$

For maxima and minima,

$$\frac{dP}{dx} = 0$$

$$\Rightarrow 15 - \frac{3}{2}x = 0$$

$$\Rightarrow x = 10$$

Also,

$$\frac{d^2P}{dx^2} = \frac{-3}{2} < 0$$

$\therefore x = 10$ is the point of local maxima

Hence, the daily output should be 10 radio sets.

Maxima and Minima 18.5 Q36

Let $S(x)$ be the selling price of x items and let $C(x)$ be the cost price of x items.

$$\text{Then, we have } S(x) = \left(5 - \frac{x}{100}\right)x = 5x - \frac{x^2}{100}$$

$$\text{and } C(x) = \frac{x}{5} + 500$$

Thus, the profit function $P(x)$ is given by

$$P(x) = S(x) - C(x) = 5x - \frac{x^2}{100} - \frac{x}{5} - 500 = \frac{24}{5}x - \frac{x^2}{100} - 500$$

$$\therefore P'(x) = \frac{24}{5} - \frac{x}{50}$$

$$\text{Now, } P'(x) = 0$$

$$\Rightarrow \frac{24}{5} - \frac{x}{50} = 0$$

$$\Rightarrow x = \frac{24}{5} \times 50 = 240$$

$$\text{Also } P''(x) = -\frac{1}{50}$$

$$\text{So, } P''(240) = -\frac{1}{50} < 0$$

Thus, $x = 240$ is a point of maxima.

Hence, the manufacturer can earn maximum profit, if he sells 240 items.

Let l be the length of side of square base of the tank and h be the height of tank.

Then,

$$\text{Volume of tank } (v) = l^2 h$$

$$\text{Total surface area } (s) = l^2 + 4lh$$

Since the tank holds a given quantity of water the volume (v) is constant.

$$\therefore v = l^2 h \quad \text{---(i)}$$

Also, cost of lining with lead will be least if the total surface area is least.

So we need to minimise the surface area.

$$\therefore S = l^2 + 4lh \quad \text{---(ii)}$$

Now,

From (i) and (ii)

$$S = l^2 + \frac{4v}{l}$$

$$\therefore \frac{ds}{dl} = 2l - \frac{4v}{l^2}$$

For maximum and minimum

$$\frac{ds}{dl} = 0$$

$$\Rightarrow 2l - \frac{4v}{l^2} = 0$$

$$\Rightarrow 2l^3 - 4v = 0$$

$$\Rightarrow l^3 = 2v = 2l^2 h$$

$$\Rightarrow l^2[l - 2h] = 0$$

$$\Rightarrow l = 0 \text{ or } 2h$$

$l = 0$ is not possible.

$$\therefore l = 2h$$

Now,


$$\frac{d^2s}{dl^2} = 2 + \frac{8v}{l^3}$$

$$\text{At } l = 2h, \frac{d^2s}{dl^2} > 0 \quad \text{for all } h.$$

$$\therefore l = 2h \text{ is point of local minima}$$

$$\therefore S \text{ is minimum when } l = 2h$$

Let $ABCDEFGH$ be a box of constant volume c . We are given that the box is twice as long as its width.



$$\begin{aligned}\therefore \quad & \text{Let } BF = x \\ \Rightarrow & AB = 2x\end{aligned}$$

Cost of material of top and front side = $3 \times$ cost of material of the bottom of the box.

$$\begin{aligned}\Rightarrow & 2x \times x + xh + xh + 2xh + 2xh = 3 \times 2x^2 \\ \Rightarrow & 2x^2 + 2xh + 4xh = 6x^2 \\ \Rightarrow & 4x^2 - 6xh = 0 \\ \Rightarrow & 2x(2x - 3h) = 0 \\ \Rightarrow & x = \frac{3h}{2} \text{ or } h = \frac{2x}{3}\end{aligned}$$

$$\text{Volume of box} = 2x \times x \times h$$

$$\begin{aligned}\Rightarrow & c = 2x^2h \\ \Rightarrow & h = \frac{c}{2x^2} \quad \text{---(ii)}\end{aligned}$$

Now,

$$\begin{aligned}S &= \text{Surface area of box} = 2(2x^2 + 2xh + xh) \\ \Rightarrow & S = 2(2x^2 + 3xh)\end{aligned}$$

From (i)

$$\begin{aligned}S &= 2\left(2x^2 + \frac{3xc}{2x^2}\right) \\ \Rightarrow & S = 2\left(2x^2 + \frac{3c}{2x}\right)\end{aligned}$$

For maxima and minima,

$$\begin{aligned}\frac{dS}{dx} &= 2\left(4x - \frac{3c}{2x^2}\right) = 0 \\ \Rightarrow & 8x^3 - 3c = 0 \\ \Rightarrow & x = \left(\frac{3c}{8}\right)^{\frac{1}{3}}\end{aligned}$$

Now,

$$\frac{d^2s}{dx^2} = 2 \left(4 + 3 \frac{c}{x^3} \right) > 0 \text{ as } x = \left(\frac{3c}{8} \right)^{\frac{1}{3}}$$

$$x = \left(\frac{3c}{8} \right)^{\frac{1}{3}} \text{ is point of local minima}$$

∴ Most economic dimension will be

$$x = \text{width} = \left(\frac{3c}{8} \right)^{\frac{1}{3}}$$

$$2x = \text{length} = 2 \left(\frac{3c}{8} \right)^{\frac{1}{3}}$$

$$h = \text{height} = \frac{2x}{3} = \frac{2}{3} \left(\frac{3c}{8} \right)^{\frac{1}{3}}.$$

Maxima and Minima 18.5 Q39

Let s be the sum of the surface areas of a sphere and a cube.

$$\therefore s = 4\pi r^2 + 6l^2 \quad \text{---(i)}$$

Let v = volume of sphere + volume of cube

$$\Rightarrow v = \frac{4}{3}\pi r^3 + l^3 \quad \text{---(ii)}$$

From (i)

$$l = \sqrt{\frac{s - 4\pi r^2}{6}}$$

$$\therefore v = \frac{4}{3}\pi r^3 + \left(\frac{s - 4\pi r^2}{6}\right)^{\frac{3}{2}}$$

$$\therefore \frac{dv}{dr} = 4\pi r^2 + \frac{3}{2} \left(\frac{s - 4\pi r^2}{6}\right)^{\frac{1}{2}} \times \left(\frac{-4\pi}{6}\right)^{\frac{1}{2}}$$

For maxima and minima,

$$\frac{dv}{dr} = 0$$

$$\Rightarrow 4\pi r^2 = \frac{\pi}{6} (s - 4\pi r^2)^{\frac{1}{2}} \times 2r = 0$$

$$\Rightarrow 2r\pi [2r - l] = 0$$

$$\therefore r = 0, \quad \frac{l}{2}$$

Now,

$$\frac{d^2v}{dr^2} = 8\pi r - \frac{2\pi}{\sqrt{6}} \left[(s - 4\pi r^2) \right]^{\frac{1}{2}} - \frac{8\pi r^2}{2(s - 4\pi r^2)^{\frac{1}{2}}}$$

$$\text{At } r = \frac{l}{2}$$

$$\frac{d^2v}{dr^2} = \pi \frac{l}{2} - \frac{2\pi}{\sqrt{6}} \left[\sqrt{6}l - \frac{8\pi \frac{l^2}{4}}{2\sqrt{6}l} \right] = 4\pi l - \frac{2\pi}{\sqrt{6}} \left[\frac{12l^2 - 2\pi l^2}{2\sqrt{6}l} \right]$$

Let ABCDEF be a half cylinder with rectangular base and semi-circular ends.

Here AB = height of the cylinder

AB = h

Let r be the radius of the cylinder.

Volume of the half cylinder is $V = \frac{1}{2} \pi r^2 h$

$$\Rightarrow \frac{2V}{\pi r^2} = h$$

$\therefore$ TSA of the half cylinder is

S = LSA of the half cylinder + area of two semi-circular ends + area of the rectangle (base)

$$S = \pi r h + \frac{\pi r^2}{2} + \frac{\pi r^2}{2} + h \times 2r$$

$$S = (\pi r + 2r)h + \pi r^2$$

$$S = (\pi r + 2r) \frac{2V}{\pi r^2} + \pi r^2$$

$$S = (\pi + 2) \frac{2V}{\pi r} + \pi r^2$$

Differentiate S wrt r we get,

$$\frac{ds}{dr} = \left[(\pi + 2) \times \frac{2V}{\pi} \left(\frac{-1}{r^2} \right) + 2\pi r \right]$$

For maximum and minimum values of S, we have $\frac{ds}{dr} = 0$

$$\Rightarrow (\pi + 2) \times \frac{2V}{\pi} \left(\frac{-1}{r^2} \right) + 2\pi r = 0$$

$$\Rightarrow (\pi + 2) \times \frac{2V}{\pi r^2} = 2\pi r$$

But $2r = D$

$$\therefore h : D = \pi : \pi + 2$$

Differentiate $\frac{ds}{dr}$ wrt r we get,

$$\frac{d^2s}{dr^2} = (\pi + 2) \times \frac{2V}{\pi} \times \frac{2}{r^3} + 2\pi > 0$$


Thus S will be minimum when h : D is $\pi : \pi + 2$.

Height of the cylinder : Diameter of the circular end

$$\pi : \pi + 2$$

Maxima and Minima 18.5 Q41

Let ABCD be the cross-sectional area of the beam which is cut from a circular log of radius a.



$$\therefore AO = a \Rightarrow AC = 2a$$

Let x be the width of log and y be the depth of log $ABCD$

Let S be the strength of the beam according to the question,

$$S = xy^2 \quad \text{---(i)}$$

In $\triangle ABC$

$$x^2 + y^2 = (2a)^2$$

$$\Rightarrow y = (2a)^2 - x^2 \quad \text{---(ii)}$$

From (i) and (ii), we get

$$S = x \left((2a)^2 - x^2 \right)$$

$$\Rightarrow \frac{dS}{dx} = (4a^2 - x^2) - 2x^2$$

$$\Rightarrow \frac{dS}{dx} = 4a^2 - 3x^2$$

For maxima or minima

$$\frac{dS}{dx} = 0$$

$$\Rightarrow 4a^2 - 3x^2 = 0$$

$$\Rightarrow x^2 = \frac{4a^2}{3}$$

$$\therefore x = \frac{2a}{\sqrt{3}}$$

From (ii),

$$y^2 = 4a^2 - \frac{4a^2}{3} = \frac{8a^2}{3}$$

$$\therefore y = 2a \times \sqrt{\frac{2}{3}}$$

Now,

$$\frac{d^2S}{dx^2} = -6x$$

$$\text{At } x = \frac{2a}{\sqrt{3}}, y = \sqrt{\frac{2}{3}} 2a, \quad \frac{d^2S}{dx^2} = -\frac{12a}{\sqrt{3}} < 0$$

$$\therefore \left(x = \frac{2a}{\sqrt{3}}, y = \sqrt{\frac{2}{3}} 2a \right) \text{ is the point of local maxima.}$$

Hence,

$$\text{The dimension of strongest beam is width } = x = \frac{2a}{\sqrt{3}} \text{ and depth } = y = \sqrt{\frac{2}{3}} 2a.$$

Maxima and Minima 18.5 Q42

Let l be a line through the point $P(1, 4)$ that cuts the x -axis and y -axis.

Now, equation of l is

$$y - 4 = m(x - 1)$$

$$\therefore x\text{-Intercept is } \frac{m-4}{m} \text{ and } y\text{-Intercept is } 4-m$$

$$\text{Let } S = \frac{m-4}{m} + 4 - m$$

$$\therefore \frac{dS}{dm} = +\frac{4}{m^2} - 1$$

For maxima and minima,

$$\frac{dS}{dm} = 0$$

$$\Rightarrow \frac{4}{m^2} - 1 = 0$$

$$\Rightarrow m = \pm 2$$

Now,

$$\frac{d^2S}{dm^2} = -\frac{8}{m^3}$$

$$\text{At } m = 2, \frac{d^2S}{dm^2} = -1 < 0$$

$$m = -2, \frac{d^2S}{dm^2} = 1 > 0$$

$$\therefore m = -2 \text{ is point of local minima.}$$

$\therefore$ least value of sum of intercept is


$$\begin{aligned} & \frac{m-4}{m} + 4 - m \\ &= 3 + 6 = 9 \end{aligned}$$

Maxima and Minima 18.5 Q43

The area of the page $PQRS$ is 150 cm^2

Also, $AB + CD = 3 \text{ cm}$

$EF + GH = 2 \text{ cm}$



Let x and y be the combined width of margin at the top and bottom and the sides respectively.

$\therefore x = 3 \text{ cm}$ and $y = 2 \text{ cm}$.

Now, area of printed matter = area of $P'Q'R'S'$

$$\Rightarrow A = P'Q' \times Q'R'$$

$$\Rightarrow A = (b - y)(l - x)$$

$$\Rightarrow A = (b - 2)(l - 3) \quad \text{---(i)}$$

Also,

$$\text{Area of } PQRS = 150 \text{ cm}^2$$

$$\Rightarrow lb = 150 \quad \text{---(ii)}$$

From (i) and (ii)

$$A = (b - 2) \left(\frac{150}{b} - 3 \right)$$

$\therefore$ For maximum and minimum,

$$\frac{dA}{db} = \left(\frac{150}{b} - 3 \right) + (b - 2) \left(-\frac{150}{b^2} \right) = 0$$

$$\Rightarrow \frac{(150 - 3b)}{b} + (-150) \frac{(b - 2)}{b^2} = 0$$

$$\Rightarrow 150b - 3b^2 - 150b + 300 = 0$$

$$\Rightarrow -3b^2 + 300 = 0$$

$$\Rightarrow b = 10$$

From (ii)

$$l = 15$$

Now,

$$\frac{d^2A}{db^2} = \frac{-150}{b^2} - 150 \left[-\frac{1}{b^2} + \frac{4}{b^3} \right]$$

At $b = 10$

$$\begin{aligned}\frac{d^2A}{db^2} &= -\frac{15}{10} - 150\left[-\frac{1}{100} + \frac{4}{1000}\right] \\ &= -1.5 - .15[-10 + 4] \\ &= -1.5 + .9 \\ &= -0.6 < 0\end{aligned}$$

$\therefore b = 10$ is point of local maxima.

Hence,

The required dimension will be $l = 15$ cm, $b = 10$ cm.

Maxima and Minima 18.5 Q44

The space s described in time t by a moving particle is given by

$$s = t^5 - 40t^3 + 30t^2 + 80t - 250$$

$$\therefore \text{velocity} = \frac{ds}{dt} = 5t^4 - 120t^2 + 60t + 80$$

$$\text{Acceleration} = a = \frac{d^2s}{dt^2} = 20t^3 - 240t + 60t \quad \text{---(i)}$$

Now,

$$\frac{da}{dt} = 60t^2 - 240$$

For maxima and minima,

$$\frac{da}{dt} = 0$$

$$\Rightarrow 60t^2 - 240 = 0$$

$$\Rightarrow 60(t^2 - 4) = 0$$

$$\Rightarrow t = 2$$

Now,

$$\frac{d^2a}{dt^2} = 120t$$

$$\text{At } t = 2, \frac{d^2a}{dt^2} = 240 > 0$$

$\therefore t = 2$ is point of local minima

Hence, minimum acceleration is $160 - 480 + 60 = -260$.

Maxima and Minima 18.5 Q45

We have,

$$\text{Distance, } s = \frac{t^4}{4} - 2t^3 + 4t^2 - 7$$

$$\text{Velocity, } v = \frac{ds}{dt} = t^3 - 6t^2 + 8t$$

$$\text{Acceleration, } a = \frac{d^2s}{dt^2} = 3t^2 - 12t + 8$$

For velocity to be maximum and minimum,

$$\frac{dv}{dt} = 0$$

$$\Rightarrow 3t^2 - 12t + 8 = 0$$

$$\Rightarrow t = \frac{12 \pm \sqrt{144 - 96}}{6}$$

$$= 2 \pm \frac{4\sqrt{3}}{6}$$

$$\therefore t = 2 + \frac{2}{\sqrt{3}}, 2 - \frac{2}{\sqrt{3}}$$

Now,

$$\frac{d^2v}{dt^2} = 6t - 12$$

$$\text{At } t = 2 - \frac{2}{\sqrt{3}}, \frac{d^2v}{dt^2} = 6\left(2 - \frac{2}{\sqrt{3}}\right) - 12 = \frac{-12}{\sqrt{3}} < 0$$

$$t = 2 + \frac{2}{\sqrt{3}}, \frac{d^2v}{dt^2} = 6\left(2 + \frac{2}{\sqrt{3}}\right) - 12 = \frac{12}{\sqrt{3}} > 0$$

$$\therefore \text{At } t = 2 - \frac{2}{\sqrt{3}}, \text{ velocity is maximum}$$

For acceleration to be maximum and minimum

$$\frac{da}{dt} = 0$$

$$\Rightarrow 6t - 12 = 0$$

$$\Rightarrow t = 2$$

Now,

$$\frac{d^2a}{dt^2} = 6 > 0$$

$$\therefore \text{At, } t = 2 \text{ Acceleration is minimum.}$$

