


RD Sharma
Solutions
Class 11 Maths
Chapter 11
Ex 11.1



Trigonometric Equations Ex 11.1 Q1(i)

We have,

$$\sin \theta = \frac{1}{2}$$

$$\Rightarrow \sin \theta = \sin \frac{\pi}{6} \quad \left[\because \sin \frac{\pi}{6} = \frac{1}{2} \right]$$

$\Rightarrow$ the general solution is

$$\theta = n\pi + (-1)^n \frac{\pi}{6}, n \in \mathbb{Z} \quad \left[\because \text{if } \sin \theta = \sin \alpha \Rightarrow \theta = n\pi + (-1)^n \alpha \right]$$

Trigonometric Equations Ex 11.1 Q1(ii)

We have,

$$\cos \theta = -\frac{\sqrt{3}}{2}$$

$$\Rightarrow \cos \theta = \cos \left(\pi + \frac{\pi}{6} \right)$$

$$\Rightarrow \cos \theta = \cos \frac{7\pi}{6} \quad \left[\because \cos \frac{7\pi}{6} = -\frac{\sqrt{3}}{2} \right]$$

$\therefore$ the general solution is

$$\theta = 2n\pi \pm \frac{7\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q1(iii)

$$\csc \theta = -\sqrt{2}$$

$$\Rightarrow \frac{1}{\sin \theta} = -\sqrt{2}$$

$$\Rightarrow \sin \theta = -\frac{1}{\sqrt{2}}$$

$$\Rightarrow \sin \theta = \sin \left(\pi + \frac{\pi}{4} \right)$$

$$\Rightarrow \sin \theta = \sin \frac{5\pi}{4} \text{ or } \sin \theta = \sin \left(-\frac{\pi}{4} \right)$$

$$\because \sin(-\theta) = -\sin \theta.$$

$$\therefore \theta = n\pi + (-1)^{n+1} \frac{\pi}{4}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q1(iv)

We have,

$$\sec \theta = \sqrt{2}$$

$$\Rightarrow \frac{1}{\cos \theta} = \sqrt{2}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \Rightarrow \cos \theta = \cos \frac{\pi}{4}$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{4}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q1(v)

We have,

$$\tan \theta = \frac{-1}{\sqrt{3}}$$

$$\Rightarrow \tan \theta = \tan \left(-\frac{\pi}{6} \right)$$

$$\Rightarrow \tan \theta = \tan \left(-\frac{\pi}{6} \right) \quad \left[\because \tan(-\theta) = -\tan \theta \right]$$

$$\Rightarrow \theta = n\pi + \left(-\frac{\pi}{6} \right), n \in \mathbb{Z}$$

$$\text{or } \theta = n\pi - \frac{\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q1(vi)

We have,

Copykitab
 Same textbooks, knock away

$$\begin{aligned}\sqrt{3} \sec \theta &= 2 \\ \Rightarrow \frac{1}{\cos \theta} &= \frac{2}{\sqrt{3}} \\ \Rightarrow \cos \theta &= \frac{\sqrt{3}}{2} \\ \Rightarrow \cos \theta &= \cos \left(\frac{\pi}{6} \right) \\ \Rightarrow \theta &= 2n\pi \pm \frac{\pi}{6}, n \in \mathbb{Z}\end{aligned}$$

Trigonometric Equations Ex 11.1 Q2(i)

We have,

$$\begin{aligned}\sin 2\theta &= \frac{\sqrt{3}}{2} \\ \Rightarrow \sin 2\theta &= \sin \left(\frac{\pi}{3} \right) \\ \Rightarrow 2\theta &= n\pi + (-1)^n \frac{\pi}{3}, n \in \mathbb{Z}\end{aligned}$$

$$\theta = \frac{n\pi}{2} + (-1)^n \frac{\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(ii)

We have,

$$\begin{aligned}\cos 3\theta &= \frac{1}{2} \\ \Rightarrow \cos 3\theta &= \cos \left(\frac{\pi}{3} \right) \\ \Rightarrow 3\theta &= 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z} \\ \Rightarrow \theta &= 2n\frac{\pi}{3} \pm \frac{\pi}{9}, n \in \mathbb{Z}\end{aligned}$$

Trigonometric Equations Ex 11.1 Q2(iii)

$$\begin{aligned}\sin 9\theta &= \sin \theta \\ \sin 9\theta - \sin \theta &= 0\end{aligned}$$

Apply $\sin A - \sin B$ formula

$$\sin A - \sin B = 2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right)$$

$$\sin 9\theta - \sin \theta = 2 \cos 5\theta \sin 4\theta = 0$$

$$\cos 5\theta \sin 4\theta = 0$$

$$\Rightarrow \cos 5\theta = 0 \text{ (or) } \sin 4\theta = 0$$

$$5\theta = \frac{(2n+1)\pi}{2} \text{ (or) } 4\theta = n\pi$$

$$\theta = \left\{ \frac{(2n+1)\pi}{10} \right\} \text{ (or) } \theta = \left\{ \frac{n\pi}{4} \right\} \text{ where } n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(vi)

We have,

$$\begin{aligned}\sin 2\theta &= \cos 3\theta \\ \Rightarrow \cos 3\theta &= \sin 2\theta \\ \Rightarrow \cos 3\theta &= \cos \left(\frac{\pi}{2} - 2\theta \right) \quad \left[\because \cos \left(\frac{\pi}{2} - \theta \right) = \sin \theta \right] \\ \Rightarrow 3\theta &= 2n\pi \pm \left(\frac{\pi}{2} - 2\theta \right), n \in \mathbb{Z} \\ \Rightarrow \text{either}\end{aligned}$$

$$5\theta = 2n\pi + \frac{\pi}{2}, n \in \mathbb{Z} \text{ or } \theta = 2n\pi - \frac{\pi}{2}, n \in \mathbb{Z}$$

$$\Rightarrow 5\theta = (4n+1)\frac{\pi}{2}, n \in \mathbb{Z} \text{ or } \theta = (4n-1)\frac{\pi}{2}$$

$$\Rightarrow \theta = (4n+1)\frac{\pi}{10}, n \in \mathbb{Z} \text{ or } \theta = (4n-1)\frac{\pi}{2}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(v)

We have,

$$\tan \theta + \cot 2\theta = 0$$

$$\tan \theta = -\cot 2\theta$$

$$\Rightarrow \cot 2\theta = -\tan \theta$$

$$\Rightarrow \tan 2\theta = -\cot \theta$$

$$\Rightarrow \tan 2\theta = -\tan \left(\frac{\pi}{2} - \theta \right)$$

$$\Rightarrow \tan 2\theta = \tan \left(\theta - \frac{\pi}{2} \right)$$

$$\Rightarrow 2\theta = n\pi + \left(\theta - \frac{\pi}{2} \right), n \in \mathbb{Z}$$

$$\Rightarrow \theta = n\pi - \frac{\pi}{2}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(vi)

We have,

$$\tan 3\theta = \cot \theta$$

$$\Rightarrow \tan 3\theta = \tan \left(\frac{\pi}{2} - \theta \right) \quad \left[\because \tan \left(\frac{\pi}{2} - \theta \right) = \cot \theta \right]$$

$$\Rightarrow 3\theta = n\pi + \frac{\pi}{2} - \theta, n \in \mathbb{Z}$$

$$\Rightarrow 4\theta = n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$$

$$\Rightarrow \theta = \frac{n\pi}{4} + \frac{\pi}{8}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(vii)

We have,

$$\tan 2\theta, \tan \theta = 1$$

$$\Rightarrow \tan 2\theta = \frac{1}{\tan \theta}$$

$$\Rightarrow \tan 2\theta = \cot \theta$$

$$\Rightarrow \tan 2\theta = \tan \left(\frac{\pi}{2} - \theta \right)$$

$$\Rightarrow 2\theta = n\pi + \frac{\pi}{2} - \theta, n \in \mathbb{Z}$$

$$\Rightarrow 3\theta = n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$$

$$\Rightarrow \theta = \frac{n\pi}{3} + \frac{\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(viii)

$$\tan m\theta + \cot n\theta = 0$$

$$\sin m\theta \sin n\theta + \cos m\theta \cos n\theta = 0$$

$$\cos(m-n)\theta = 0$$

$$(m-n)\theta = \left(\frac{2k+1}{2} \right) \pi$$

$$\theta = \left(\frac{2k+1}{2(m-n)} \right) \pi, k \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(ix)

We have,

$$\tan p\theta = \cot q\theta$$

$$\Rightarrow \tan p\theta = \tan \left(\frac{\pi}{2} - q\theta \right)$$

$$\Rightarrow p\theta = n\pi + \left(\frac{\pi}{2} - q\theta \right), n \in \mathbb{Z}$$

$$\Rightarrow (p+q)\theta = n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$$

$$\Rightarrow (p+q)\theta = (2n+1) \frac{\pi}{2}, n \in \mathbb{Z}$$

$$\Rightarrow \theta = \frac{(2n+1) \pi}{(p+q) 2}, n \in \mathbb{Z}$$

Copykitab
Same textbooks, click away

Trigonometric Equations Ex 11.1 Q2(x)

$$\sin 2x + \cos x = 0$$

$$2 \sin x \cos x + \cos x = 0$$

$$\cos x (2 \sin x + 1) = 0$$

$$\cos x = 0 \text{ or } 2 \sin x + 1 = 0$$

$$x = (4m-1)\frac{\pi}{2} \text{ or } \sin x = \frac{-1}{2}$$

$$x = (4m-1)\frac{\pi}{2} \text{ or } x = (4n-1)\frac{\pi}{6}, \quad m, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(xi)

We have,

$$\sin \theta = \tan \theta$$

$$\Rightarrow \sin \theta = \frac{\sin \theta}{\cos \theta}$$

$$\Rightarrow \sin \theta = \frac{\sin \theta}{\cos \theta} = 0$$

$$\Rightarrow \sin \theta (\cos \theta - 1) = 0$$

$$\Rightarrow \text{either } \sin \theta = 0 \quad \text{or } \cos \theta - 1 = 0$$

$$\Rightarrow \theta = n\pi, n \in \mathbb{Z} \quad \text{or } \cos \theta = 1$$

$$\Rightarrow \cos \theta = \cos 0^\circ$$

$$\theta = 2m\pi, m \in \mathbb{Z}$$

Thus,

$$\theta = n\pi, n \in \mathbb{Z} \quad \text{or } \theta = 2m\pi, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q2(xii)

$$\cos(2x) = -\sin(3x)$$

$$= -\cos\left(\frac{\pi}{2} - 3x\right)$$

$$= \cos\left(\frac{\pi}{2} + 3x\right)$$

$$\Rightarrow 2n\pi + 2x = \frac{\pi}{2} + 3x$$

$$x = (4m-1)\frac{\pi}{2}, m \in \mathbb{Z}$$

or

$$\Rightarrow 2n\pi - 2x = \frac{\pi}{2} + 3x$$

$$x = (4n-1)\frac{\pi}{10}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q3(i)

We have,

$$\sin^2 \theta - \cos \theta = \frac{1}{4}$$

$$\Rightarrow 1 - \cos^2 \theta - \cos \theta = \frac{1}{4} \quad \left[\because \sin^2 \theta = 1 - \cos^2 \theta \right]$$

$$\Rightarrow \cos^2 \theta + \cos \theta - \frac{3}{4} = 0$$

$$\Rightarrow 4\cos^2 \theta + 4\cos \theta - 3 = 0$$

$$\Rightarrow 4\cos^2 \theta + 6\cos \theta - 2\cos \theta - 3 = 0 \quad \left[\text{factorize it} \right]$$

$$\Rightarrow 2\cos \theta (2\cos \theta + 3) - 1(\cos \theta + 3) = 0$$

$$\Rightarrow (2\cos \theta - 1)(2\cos \theta + 3) = 0$$

$\Rightarrow$ either

$$2\cos \theta - 1 = 0 \quad \text{or } 2\cos \theta + 3 = 0$$

$$\Rightarrow \cos \theta = \frac{1}{2} \quad \text{or } \cos \theta = -\frac{3}{2} \quad \left[\text{This is not possible as } -1 < \cos \theta < 1 \right]$$

$$\Rightarrow \cos \theta = \cos \frac{\pi}{3}$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q3(ii)

We have,

$$2\cos^2\theta - 5\cos\theta + 2 = 0$$

$$\Rightarrow 2\cos^2\theta - 4\cos\theta - \cos\theta + 2 = 0 \quad [\text{use factorization}]$$

$$\Rightarrow 2\cos\theta(\cos\theta - 2) - 1(\cos\theta - 2) = 0$$

$$\Rightarrow (2\cos\theta - 1)(\cos\theta - 2) = 0$$

 $\Rightarrow$ either

$$2\cos\theta - 1 = 0 \quad \text{or} \quad \cos\theta - 2 = 0$$

$$\Rightarrow \cos\theta = \frac{1}{2} \quad \text{or} \quad \cos\theta = 2$$

$$\Rightarrow \cos\theta = \cos\frac{\pi}{3} \quad [\text{This is not possible as } -1 < \cos\theta < 1]$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$$

Thus,

$$\theta = 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q3(iii)

We have,

$$2\sin^2x + \sqrt{3}\cos x + 1 = 0$$

$$\Rightarrow 2(1 - \cos^2x) + \sqrt{3}\cos x + 1 = 0$$

$$\Rightarrow 2\cos^2x - \sqrt{3}\cos x - 3 = 0$$

factorise it, we get,

$$\Rightarrow 2\cos^2x - 2\sqrt{3}\cos x + \sqrt{3}\cos x - 3 = 0$$

$$\Rightarrow 2\cos x(\cos x - \sqrt{3}) + \sqrt{3}(\cos x - \sqrt{3}) = 0$$

$$\Rightarrow (2\cos x + \sqrt{3})(\cos x - \sqrt{3}) = 0$$

 $\Rightarrow$ either

$$\cos x = -\frac{\sqrt{3}}{2} \quad \text{or} \quad \cos x = \sqrt{3} \quad [\text{This is not possible as } -1 < \cos x < 1]$$

$$\Rightarrow \cos x = \cos\left(\pi - \frac{\pi}{6}\right)$$

$$\Rightarrow \cos x = \cos\frac{5\pi}{6}$$

$$\Rightarrow x = 2n\pi \pm \frac{5\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q3(vi)

We have,

$$4\sin^2\theta - 8\cos\theta + 1 = 0$$

$$\Rightarrow 4(1 - \cos^2\theta) - 8\cos\theta + 1 = 0$$

$$\Rightarrow 4\cos^2\theta + 8\cos\theta - 5 = 0$$

factorise it, we get,

$$\Rightarrow 4\cos^2\theta + 10\cos\theta - 2\cos\theta - 5 = 0$$

$$\Rightarrow 2\cos\theta(2\cos\theta + 5) - 1(2\cos\theta + 5) = 0$$

$$\Rightarrow (2\cos\theta - 1)(2\cos\theta + 5) = 0$$

$$\text{either } 2\cos\theta - 1 = 0 \quad \text{or} \quad 2\cos\theta + 5 = 0$$

$$\Rightarrow \cos\theta = \frac{1}{2} \quad \text{or} \quad \cos\theta = -\frac{5}{2} \quad [\text{This is not possible as } -1 < \cos\theta < 1]$$

$$\Rightarrow \cos\theta = \cos\frac{\pi}{3}$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q3(v)

We have,

$$\tan^2x + (1 - \sqrt{3})\tan x - \sqrt{3} = 0$$

$$\Rightarrow \tan^2x + \tan x - \sqrt{3}\tan x - \sqrt{3} = 0$$

$$\Rightarrow \tan x(\tan x + 1) - \sqrt{3}(\tan x + 1) = 0$$

$$\Rightarrow (\tan x - \sqrt{3})(\tan x + 1) = 0$$

$$\begin{aligned} \Rightarrow & \text{either} \\ & \tan x = \sqrt{3} \quad \text{or} \quad \tan x = -1 \\ \Rightarrow & \tan x = \tan \frac{\pi}{3} \quad \text{or} \quad \tan x = -\tan \frac{\pi}{4} \\ \Rightarrow & x = n\pi + \frac{\pi}{3}, n \in \mathbb{Z} \quad \text{or} \quad x = m\pi - \frac{\pi}{4}, m \in \mathbb{Z} \\ \therefore x = n\pi + \frac{\pi}{3} \quad \text{or} \quad m\pi - \frac{\pi}{4}, n, m \in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q3(vi)

$$\begin{aligned} 3\cos^2 \theta - 2\sqrt{3}\sin \theta \cos \theta - 3\sin^2 \theta &= 0 \\ \sqrt{3}\cos^2 \theta - 2\sin \theta \cos \theta - \sqrt{3}\sin^2 \theta &= 0 \quad (\text{Dividing by } \sqrt{3}) \\ \sqrt{3}\cos^2 \theta + \sin \theta \cos \theta - 3\sin \theta \cos \theta - \sqrt{3}\sin^2 \theta &= 0 \\ \cos \theta (\sqrt{3}\cos \theta + \sin \theta) - \sqrt{3}\sin \theta (\sqrt{3}\cos \theta + \sin \theta) &= 0 \\ (\sqrt{3}\cos \theta + \sin \theta)(\cos \theta - \sqrt{3}\sin \theta) &= 0 \\ \sqrt{3}\cos \theta + \sin \theta &= 0 \quad \text{or} \quad \cos \theta - \sqrt{3}\sin \theta = 0 \\ \tan \theta = -\sqrt{3} = -\tan \frac{\pi}{3} \quad \text{or} \quad \tan \theta = \frac{1}{\sqrt{3}} = \tan \frac{\pi}{6} \\ \theta = n\pi - \frac{\pi}{3} \quad \text{or} \quad \theta = m\pi + \frac{\pi}{6} \\ n, m &\in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q3(vii)

We have,

$$\begin{aligned} \cos 4\theta &= \cos 2\theta \\ \Rightarrow \cos 4\theta - \cos 2\theta &= 0 \\ \Rightarrow 2\sin \theta \cdot \sin 3\theta &= 0 \\ \Rightarrow \text{either} \\ \sin \theta &= 0 \quad \text{or} \quad \sin 3\theta = 0 \\ \Rightarrow \theta = n\pi, n \in \mathbb{Z} \quad \text{or} \quad 3\theta = m\pi, m \in \mathbb{Z} \end{aligned}$$

Thus,

$$\theta = n\pi \quad \text{or} \quad m\frac{\pi}{3}, n, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(i)

$$\begin{aligned} \cos \theta + \cos 2\theta + \cos 3\theta &= 0 \\ \Rightarrow \cos 2\theta + 2\cos 2\theta \cdot \cos \theta &= 0 \quad \left[\because \cos \theta + \cos 3\theta = 2\cos 2\theta \cdot \cos \theta \right] \\ \Rightarrow \cos 2\theta (1 + 2\cos \theta) &= 0 \\ \text{either} \\ \cos 2\theta &= 0 \quad \text{or} \quad 1 + 2\cos \theta = 0 \\ \Rightarrow 2\theta = (2n+1)\frac{\pi}{4}, n \in \mathbb{Z} \quad \text{or} \quad \cos \theta &= -\frac{1}{2} \\ \Rightarrow \theta = (2n+1)\frac{\pi}{4}, n \in \mathbb{Z} \quad \text{or} \quad \cos \theta &= +\cos \left(\pi - \frac{\pi}{3} \right) \\ &\text{or} \quad \cos \theta = \cos 2\frac{\pi}{3} \\ &\text{or} \quad \theta = 2n\pi \pm \frac{2\pi}{3}, n \in \mathbb{Z} \end{aligned}$$

Thus,

$$\theta = (2n+1)\frac{\pi}{4}, \quad \text{or} \quad \left(2n\pi \pm \frac{2\pi}{3} \right), n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(ii)

$$\begin{aligned} \cos \theta + \cos 3\theta - \cos 2\theta &= 0 \\ \Rightarrow 2\cos 2\theta \cdot \cos \theta - \cos 2\theta &= 0 \\ \Rightarrow \cos 2\theta (2\cos \theta - 1) &= 0 \\ \text{either} \\ \cos 2\theta &= 0 \quad \text{or} \quad 2\cos \theta = 1 \\ \Rightarrow 2\theta = (2n+1)\frac{\pi}{2}, n \in \mathbb{Z} \quad \text{or} \quad \cos \theta &= \frac{1}{2} = \cos \frac{\pi}{3} \\ \Rightarrow \theta = (2n+1)\frac{\pi}{4}, n \in \mathbb{Z} \quad \text{or} \quad \theta &= 2m\pi \pm \frac{\pi}{3}, m \in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q4(iii)

$$\begin{aligned} \sin \theta + \sin 5\theta &= \sin 3\theta \\ \Rightarrow 2\sin 3\theta \cdot \cos 2\theta - \sin 3\theta &= 0 \quad \left[\because \sin C + \sin D = 2\sin \frac{C+D}{2} \cdot \cos \frac{C-D}{2} \right] \end{aligned}$$

Wop/kitab
Same textbooks, knock away

$$\begin{aligned}
 &\Rightarrow \sin 3\theta [2 \cos 2\theta - 1] = 0 \\
 &\Rightarrow \text{either} \\
 &\quad \sin 3\theta = 0 \quad \text{or} \quad 2 \cos 2\theta - 1 = 0 \\
 &\Rightarrow 3\theta = n\pi, n \in \mathbb{Z} \quad \text{or} \quad \cos 2\theta = \frac{1}{2} = \cos \frac{\pi}{3} \\
 &\Rightarrow \theta = \frac{n\pi}{3}, n \in \mathbb{Z} \quad \text{or} \quad 2\theta = 2m\pi \pm \frac{\pi}{3}, m \in \mathbb{Z} \\
 &\quad \text{or} \quad \theta = m\pi \pm \frac{\pi}{6}
 \end{aligned}$$

Thus,

$$\theta = \frac{n\pi}{3} \quad \text{or} \quad m\pi \pm \frac{\pi}{6}, n, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(vi)

We have,

$$\begin{aligned}
 &\cos \theta \cdot \cos 2\theta \cdot \cos 3\theta = \frac{1}{4} \\
 &\Rightarrow 2 \cos \theta \cdot \cos 3\theta \cdot \cos 2\theta = \frac{1}{2} \\
 &\Rightarrow (\cos 4\theta + \cos 2\theta) \cos 2\theta = \frac{1}{2} \\
 &\Rightarrow (2 \cos^2 2\theta - 1 + \cos 2\theta) \cos 2\theta = \frac{1}{2} \\
 &\Rightarrow 2 \cos^3 2\theta + \cos^2 2\theta - \cos 2\theta = \frac{1}{2} \\
 &\Rightarrow 4 \cos^2 2\theta + 2 \cos^2 2\theta - 2 \cos 2\theta - 1 = 0 \\
 &\Rightarrow 2 \cos^2 2\theta (2 \cos \theta + 1) - 1 (2 \cos 2\theta + 1) = 0 \\
 &\Rightarrow (2 \cos^2 2\theta - 1) (2 \cos 2\theta + 1) = 0
 \end{aligned}$$

either

$$\begin{aligned}
 &2 \cos^2 2\theta - 1 = 0 \quad \text{or} \quad \Rightarrow 2 \cos 2\theta + 1 = 0 \\
 &\Rightarrow \cos 4\theta = 0 \quad \text{or} \quad \Rightarrow \cos 2\theta = -\frac{1}{2} \\
 &\Rightarrow 4\theta = (2n+1) \frac{\pi}{2} \quad \text{or} \quad \Rightarrow \cos 2\theta = \cos 2 \frac{\pi}{3} \\
 &\Rightarrow \theta = (2n+1) \frac{\pi}{8} \quad \text{or} \quad \Rightarrow 2\theta = 2m\pi \pm 2 \frac{\pi}{3} \\
 &\quad \Rightarrow \theta = m\pi \pm \frac{\pi}{3}
 \end{aligned}$$

Thus,

$$\theta = (2n+1) \frac{\pi}{8} \quad \text{or} \quad \theta = m\pi \pm \frac{\pi}{3}, m, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(v)

We have,

$$\begin{aligned}
 &\cos \theta + \sin \theta = \cos 2\theta + \sin 2\theta \\
 &\Rightarrow \cos \theta - \cos 2\theta = \sin 2\theta - \sin \theta \\
 &\Rightarrow 2 \sin \frac{3\theta}{2} \cdot \sin \frac{\theta}{2} = 2 \cos \frac{3\theta}{2} \cdot \sin \frac{\theta}{2} \\
 &\Rightarrow 2 \sin \frac{\theta}{2} \left(\sin \frac{3\theta}{2} - \cos \frac{3\theta}{2} \right) = 0 \\
 &\Rightarrow 2 \sin \frac{\theta}{2} \left(\sin \frac{3\theta}{2} - \cos \frac{3\theta}{2} = 0 \right)
 \end{aligned}$$

either

$$\begin{aligned}
 &\sin \frac{\theta}{2} = 0 \quad \text{or} \quad \sin \frac{3\theta}{2} - \cos \frac{3\theta}{2} = 0 \\
 &\Rightarrow \frac{\theta}{2} = n\pi, n \in \mathbb{Z} \quad \text{or} \quad \tan \frac{3\theta}{2} = 1 = \tan \frac{\pi}{4} \\
 &\Rightarrow \theta = 2n\pi, n \in \mathbb{Z} \quad \text{or} \quad \frac{3\theta}{2} = n\pi + \frac{\pi}{4} \\
 &\quad \text{or} \quad \theta = 2n \frac{\pi}{3} + \frac{\pi}{3.2}, n \in \mathbb{Z}
 \end{aligned}$$

Thus,

$$\theta = 2n\pi \quad \text{or} \quad 2n \frac{\pi}{3} + \frac{\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(vi)

We have,

$$\begin{aligned}
 &\sin \theta + \sin 2\theta + \sin 3\theta = 0 \\
 &\Rightarrow \sin 2\theta + 2 \sin 2\theta \cdot \cos \theta = 0 \\
 &\Rightarrow \sin 2\theta + (1 + 2 \cos \theta) = 0 \\
 &\Rightarrow \text{either}
 \end{aligned}$$

$$\begin{aligned} \sin 2\theta &= 0 & \text{or } 1 + 2\cos \theta &= 0 \\ \Rightarrow 2\theta &= n\pi, n \in \mathbb{Z} & \text{or } \cos \theta &= -\frac{1}{2} = \cos\left(\pi - \frac{\pi}{3}\right) \\ \Rightarrow \theta &= \frac{n\pi}{2}, n \in \mathbb{Z} & \text{or } \theta &= 2m\pi \pm \frac{2\pi}{3}, m \in \mathbb{Z} \end{aligned}$$

Thus,

$$\theta = \frac{n\pi}{2}, n \in \mathbb{Z} \quad \text{or} \quad \theta = 2m\pi \pm \frac{2\pi}{3}, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(vii)

Given, $\sin x + \sin 2x + \sin 3x + \sin 4x = 0$

$$(\sin 4x + \sin 2x) + (\sin 3x + \sin x) = 0$$

Using, $(\sin A + \sin B)$ formula $\Rightarrow$

$$2\sin\left[\frac{(4x+2x)}{2}\right]\cos\left[\frac{(4x-2x)}{2}\right] + 2\sin\left[\frac{(3x+x)}{2}\right]\cos\left[\frac{(3x-x)}{2}\right] = 0$$

$$2\sin 3x \cos x + 2\sin 2x \cos x = 0$$

$$2\cos x (\sin 3x + \sin 2x) = 0$$

$$2\cos x \left(2\sin\left[\frac{(3x+2x)}{2}\right]\cos\left[\frac{(3x-2x)}{2}\right]\right) = 0$$

$$4\cos x \sin \frac{5x}{2} \cos \frac{x}{2} = 0$$

$$\cos x = 0; \sin \frac{5x}{2} = 0; \cos \frac{x}{2} = 0$$

$$x = \frac{(2n+1)\pi}{2}; \frac{5x}{2} = m\pi; \frac{x}{2} = \frac{(2r+1)\pi}{2}$$

$$x = \frac{(2n+1)\pi}{2}; x = \frac{2m\pi}{5}; x = (2r+1)\pi, m, r, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(viii)

We have,

$$\sin 3\theta - \sin \theta = 4\cos^2 \theta - 2$$

$$\Rightarrow 2\cos 2\theta \cdot \sin \theta = 2(2\cos^2 \theta - 1)$$

$$\Rightarrow 2\cos 2\theta \cdot \sin \theta = 2\cos 2\theta \quad \left[\because \cos 2\theta = 2\cos^2 \theta - 1\right]$$

$$\Rightarrow 2\cos 2\theta (\sin \theta - 1) = 0$$

either

$$\cos 2\theta = 0 \quad \text{or} \quad \sin \theta - 1 = 0$$

$$\Rightarrow 2\theta = (2n+1)\frac{\pi}{2}, n \in \mathbb{Z} \quad \text{or} \quad \sin \theta = 1 = \sin \frac{\pi}{2}$$

$$\Rightarrow \theta = (2n+1)\frac{\pi}{4}, n \in \mathbb{Z} \quad \text{or} \quad \theta = m\pi + (-1)^m \frac{\pi}{2}, m \in \mathbb{Z}$$

Thus,

$$\theta = (2n+1)\frac{\pi}{4}, n \in \mathbb{Z} \quad \text{or} \quad m\pi + (-1)^m \frac{\pi}{2}, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q4(viii)

$$\sin 2x - \sin 4x + \sin 6x = 0$$

$$(\sin 2x + \sin 6x) - \sin 4x = 0$$

$$2\sin\left(\frac{8x}{2}\right)\cos\left(\frac{4x}{2}\right) - \sin 4x = 0$$

$$2\sin 4x \cdot \cos 2x - \sin 4x = 0$$

$$\sin 4x (2\cos 2x - 1) = 0$$

$$\sin 4x = 0 \quad \text{or} \quad 2\cos 2x - 1 = 0$$

$$4x = n(\pi) \quad \text{or} \quad \cos 2x = 1/2$$

$$x = \left[\frac{n\pi}{4}\right] \quad \text{or} \quad \cos 2x = \cos \left[\frac{\pi}{3}\right]$$

$$x = \left[\frac{n\pi}{4}\right] \quad \text{or} \quad x = n(\pi) \pm \left[\frac{\pi}{6}\right]$$

Trigonometric Equations Ex 11.1 Q5(i)

$$\tan x + \tan 2x + \frac{(\tan x + \tan 2x)}{1 - \tan x \cdot \tan 2x} = 0$$

$$[\tan x + \tan 2x] \left[1 + \frac{1}{1 - \tan x \cdot \tan 2x}\right] = 0$$

$$\tan x + \tan 2x (2 - \tan x \cdot \tan 2x) = 0$$

$$\tan x = \tan(-2x) \quad \text{or} \quad \tan x \cdot \tan 2x = 2$$

$$x = n\pi - 2x \text{ or } \tan x \cdot \frac{2 \tan x}{1 - \tan^2 x} = 2$$

$$3x = n\pi \text{ or } \frac{2 \tan^2 x}{1 - \tan^2 x} = 2$$

$$3x = n\pi \text{ or } 2 \tan^2 x = 2 - 2 \tan^2 x$$

$$3x = n\pi \text{ or } 4 \tan^2 x = 2$$

$$x = \frac{n\pi}{3} \text{ or } \tan^2 x = 1/2$$

$$x = \frac{n\pi}{3} \text{ or } x = m\pi \pm \tan^{-1}\left(\frac{1}{\sqrt{2}}\right), \quad n, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q5(ii)

$$\tan \theta + \tan 2\theta = \tan(\theta + 2\theta)$$

$$\tan \theta + \tan 2\theta - \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \tan 2\theta} = 0$$

$$[\tan \theta + \tan 2\theta] \left[1 - \frac{1}{1 - \tan \theta \tan 2\theta} \right] = 0$$

$$[\tan \theta + \tan 2\theta] \left[\frac{1 - \tan \theta \tan 2\theta - 1}{1 - \tan \theta \tan 2\theta} \right] = 0$$

$$[\tan \theta + \tan 2\theta] \left[\frac{-\tan \theta \tan 2\theta}{1 - \tan \theta \tan 2\theta} \right] = 0$$

$$\tan \theta = 0 \text{ or } \tan 2\theta = 0 \text{ or } \tan \theta + \tan 2\theta = 0$$

$$\theta = n\pi \text{ or } \frac{n\pi}{2} \text{ or } \tan \theta \left[\frac{1 - \tan^2 \theta + 2}{1 - \tan^2 \theta} \right] = 0$$

$$\theta = n\pi \text{ or } \frac{n\pi}{2} \text{ or } \tan \theta = \pm \sqrt{3}$$

$$\theta = m\pi \text{ or } \frac{n\pi}{3}, \quad m, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q5(iii)

We have,

$$\tan 3\theta + \tan \theta = 2 \tan 2\theta$$

$$\Rightarrow \tan 3\theta - \tan 2\theta = \tan 2\theta - \tan \theta$$

$$\Rightarrow \tan 3\theta - \tan 2\theta = \tan 2\theta - \tan \theta$$

$$\Rightarrow 2 \sin^2 \theta \sin 2\theta = 0$$

$\Rightarrow$ either

$$\sin \theta = 0 \quad \text{or} \quad \sin 2\theta = 0$$

$$\Rightarrow \theta = n\pi, n \in \mathbb{Z} \quad \text{or} \quad 2\theta = m\pi, m \in \mathbb{Z}$$

$$\Rightarrow \theta = n\pi, n \in \mathbb{Z} \quad \text{or} \quad \theta = m \frac{\pi}{2}, m \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q6(i)

We have,

$$\sin \theta + \cos \theta = \sqrt{2}$$

$$\Rightarrow \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta = 1$$

$$\Rightarrow \sin \frac{\pi}{4} \sin \theta + \cos \frac{\pi}{4} \cos \theta = 1 \quad \left[\because \cos \frac{\pi}{4} = \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}} \right]$$

$$\Rightarrow \cos \left(\theta - \frac{\pi}{4} \right) = \cos 0^\circ$$

$$\Rightarrow \theta - \frac{\pi}{4} = 2n\pi, n \in \mathbb{Z}$$

$$\Rightarrow \theta = 2n\pi + \frac{\pi}{4}, n \in \mathbb{Z}$$

$$\therefore \theta = \left(8n + 1 \right) \frac{\pi}{4}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q6(ii)

$$\sqrt{3} \cos \theta + \sin \theta = 1$$

Divide both side by 2, we get

$$\frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta = \frac{1}{2}$$

$$\Rightarrow \cos \frac{\pi}{6} \cos \theta + \sin \frac{\pi}{6} \sin \theta = \frac{1}{2} \quad \left[\because \sin \frac{\pi}{6} = \frac{1}{2}, \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \right]$$

$$\begin{aligned} \Rightarrow \quad \cos\left(\theta - \frac{\pi}{6}\right) &= \cos \frac{\pi}{3} \\ \Rightarrow \quad \theta - \frac{\pi}{6} &= 2n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z} \\ \Rightarrow \quad \theta &= 2n\pi \pm \frac{\pi}{3} + \frac{\pi}{6}, n \in \mathbb{Z} \\ \Rightarrow \quad \theta &= (4n+1)\frac{\pi}{6} \quad \text{or} \quad (12m-1)\frac{\pi}{6}, n, m \in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q6(iii)

We have,

$$\sin \theta + \cos \theta = 1$$

divide both side by $\sqrt{2}$, we get,

$$\begin{aligned} \Rightarrow \quad \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta &= \frac{1}{\sqrt{2}} \\ \Rightarrow \quad \sin \frac{\pi}{4} \sin \theta + \cos \frac{\pi}{4} \cos \theta &= \frac{1}{\sqrt{2}} \\ \Rightarrow \quad \cos\left(\theta - \frac{\pi}{4}\right) &= \cos \frac{\pi}{4} \\ \Rightarrow \quad \theta - \frac{\pi}{4} &= 2n\pi \pm \frac{\pi}{4}, n \in \mathbb{Z} \\ \Rightarrow \quad \theta &= 2n\pi + \frac{\pi}{2} \quad \text{or} \quad 2n\pi, n \in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q6(iv)

We have,

$$\cos \sec \theta = 1 + \cot \theta$$

$$\Rightarrow \quad \frac{1}{\sin \theta} = 1 + \frac{\cos \theta}{\sin \theta}$$

$$\Rightarrow \quad 1 = \sin \theta + \cos \theta$$

Divide both side by $\sqrt{2}$, we get,

$$\begin{aligned} \Rightarrow \quad \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta &= \frac{1}{\sqrt{2}} \\ \Rightarrow \quad \sin \frac{\pi}{4} \sin \theta + \cos \frac{\pi}{4} \cos \theta &= \frac{1}{\sqrt{2}} \\ \Rightarrow \quad \cos\left(\theta - \frac{\pi}{4}\right) &= \cos \frac{\pi}{4} \\ \Rightarrow \quad \theta - \frac{\pi}{4} &= 2n\pi \pm \frac{\pi}{4}, n \in \mathbb{Z} \\ \therefore \theta &= \left(2n\pi + \frac{\pi}{2}\right) \quad \text{or} \quad 2n\pi, n \in \mathbb{Z} \end{aligned}$$

Trigonometric Equations Ex 11.1 Q6(v)

$$(\sqrt{3}-1)\cos \theta + (\sqrt{3}+1)\sin \theta = 2$$

Divide on both sides by $2\sqrt{2}$

$$\frac{(\sqrt{3}-1)}{2\sqrt{2}}\cos \theta + \frac{(\sqrt{3}+1)}{2\sqrt{2}}\sin \theta = \frac{1}{\sqrt{2}}$$

$$\sin\left(\theta + \tan^{-1}\left(\frac{\sqrt{3}-1}{\sqrt{3}+1}\right)\right) = \sin \frac{\pi}{4}$$

$$\theta = 2n\pi + \frac{\pi}{3} \text{ or } 2n\pi - \frac{\pi}{6}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q7(i)

$$\cot x + \tan x = 2$$

$$2 \sin x \cos x = 1$$

$$\sin 2x = 1$$

$$2x = \frac{(2n+1)\pi}{2}$$

$$x = \frac{(2n+1)\pi}{4}, n \in \mathbb{Z}$$

Trigonometric Equations Ex 11.1 Q7(ii)

$$2\sin^2 \theta = 3\cos \theta$$

$$\begin{aligned}
 2 - 2\cos^2 \theta &= 3\cos \theta \\
 2\cos^2 \theta + 3\cos \theta - 2 &= 0 \\
 2\cos^2 \theta + 4\cos \theta - \cos \theta - 2 &= 0 \\
 (\cos \theta + 2)(2\cos \theta - 1) &= 0 \\
 \cos \theta &= -2 \text{ or } \cos \theta = 0.5 \\
 \cos \theta &= -2, \text{ never possible} \\
 \cos \theta &= 0.5, \theta = 60, 300
 \end{aligned}$$

Trigonometric Equations Ex 11.1 Q7(iii)

$$\sec x \cos 5x + 1 = 0$$

$$\frac{\cos 5x + \cos x}{\cos x} = 0 \Rightarrow \cos x \neq 0$$

$$2\cos 3x \cos 2x = 0$$

$$\cos 3x = 0 \text{ or } \cos 2x = 0$$

$$3x = \frac{\pi}{2} \text{ or } 2x = \frac{\pi}{2}$$

$$x = \frac{\pi}{4}, \frac{\pi}{6}$$

Trigonometric Equations Ex 11.1 Q7(iv)

$$2\sin^2 \theta + 5 - 6 = 0$$

$$\sin^2 \theta = \frac{1}{2}$$

$$\sin \theta = \pm \frac{1}{\sqrt{2}}$$

$$\theta = n\pi \pm \frac{\pi}{4}$$

Trigonometric Equations Ex 11.1 Q7(v)

$$\sin x - 3\sin 2x + \sin 3x = \cos x - 3\cos 2x + \cos 3x$$

$$(\sin x + \sin 3x) - 3\sin 2x - (\cos x + \cos 3x) + 3\cos 2x = 0$$

$$2\sin 2x \cos x - 3\sin 2x - 2\cos 2x \cos x + 3\cos 2x = 0$$

$$\sin 2x(2\cos x - 3) - \cos 2x(2\cos x - 3) = 0$$

$$(2\cos x - 3)(\sin 2x - \cos 2x) = 0$$

$$\cos x = \frac{3}{2} \text{ or } \sin 2x - \cos 2x = 0$$

$$\text{but } \cos x \in [-1, 1] \Rightarrow \cos x \neq \frac{3}{2}$$

$$\sin 2x = \cos 2x$$

$$\tan 2x = 1$$

$$2x = n\pi + \frac{\pi}{4}$$

$$x = \frac{n\pi}{2} + \frac{\pi}{8}$$