


RD Sharma
Solutions
Class 11 Maths
Chapter 14
Ex 14.1



Quadratic Equations Ex 14.1 Q1

$$x^2 + 1 = 0$$

$$\Rightarrow x^2 + i^2 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow (x + i)(x - i) = 0 \quad [a^2 - b^2 = (a + b)(a - b)]$$

$$\Rightarrow x = i, -i$$

Quadratic Equations Ex 14.1 Q2

$$9x^2 + 4 = 0$$

$$\Rightarrow (3x)^2 - (2i)^2 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow (3x + 2i)(3x - 2i) = 0$$

$$\Rightarrow 3x + 2i = 0 \quad \text{or} \quad 3x - 2i = 0$$

$$\Rightarrow x = \frac{-2}{3}i \quad \text{or} \quad x = \frac{2}{3}i$$

$$\therefore x = \frac{-2}{3}i, \frac{2}{3}i$$

Quadratic Equations Ex 14.1 Q3

$$x^2 + 2x + 5 = 0$$

Now, completing the squares, we get

$$(x + 1)^2 + 4 = 0$$

$$\Rightarrow (x + 1)^2 - 2i^2 = 0$$

$$\Rightarrow (x + 1 + 2i)(x + 1 - 2i) = 0$$

$$\Rightarrow (x + 1 + 2i) = 0 \quad \text{or} \quad (x + 1 - 2i) = 0$$

$$\therefore x = -1 - 2i, \quad -1 + 2i$$

Quadratic Equations Ex 14.1 Q4

$$4x^2 - 12x + 25 = 0$$

Now, completing the squares, we get

$$(2x - 3)^2 + 16 = 0$$

$$\Rightarrow (2x - 3)^2 - 4i^2 = 0$$

$$\Rightarrow (2x - 3 + 4i)(2x - 3 - 4i) = 0$$

$$\Rightarrow (2x - 3 + 4i) = 0 \quad \text{or} \quad (2x - 3 - 4i) = 0$$

$$\therefore x = \frac{3}{2} + 2i, \quad \frac{3}{2} - 2i$$

Quadratic Equations Ex 14.1 Q5

$$x^2 + x + 1 = 0$$

Now, completing the squares, we get

$$\begin{aligned} & \left(x + \frac{1}{2}\right)^2 + \frac{3}{4} = 0 \\ \Rightarrow & \left(x + \frac{1}{2}\right)^2 - \left(\frac{\sqrt{3}}{2}i\right)^2 = 0 \\ \Rightarrow & \left(x + \frac{1}{2} + \frac{\sqrt{3}}{2}i\right)\left(x + \frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 0 \\ \Rightarrow & \left(x + \frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = 0 \quad \text{or} \quad \left(x + \frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 0 \\ \therefore x = & \frac{-1}{2} + \frac{\sqrt{3}}{2}i, \quad \frac{-1}{2} - \frac{\sqrt{3}}{2}i \end{aligned}$$

Quadratic Equations Ex 14.1 Q6

$$4x^2 + 1 = 0$$

$$\begin{aligned} \Rightarrow & (2x)^2 - i^2 = 0 \quad [\because i^2 = -1] \\ \Rightarrow & (2x + i)(2x - i) = 0 \\ \Rightarrow & \text{either } 2x + i = 0 \quad \text{or} \quad 2x - i = 0 \\ \Rightarrow & x = \frac{-i}{2} \quad \text{or} \quad x = \frac{i}{2} \end{aligned}$$

$$\therefore x = \frac{-i}{2}, \frac{i}{2}$$

Quadratic Equations Ex 14.1 Q7

$$x^2 - 4x + 7 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\text{where } D = b^2 - 4ac = (-4)^2 - 4.1.7 = -12$$

from (A)

$$x = \frac{-(-4) \pm \sqrt{-12}}{2}$$

$$= \frac{4 \pm 2\sqrt{3}i}{2}$$

$$= 2 \pm \sqrt{3}i$$

$$\therefore x = 2 + \sqrt{3}i, 2 - \sqrt{3}i$$

Quadratic Equations Ex 14.1 Q8

$$x^2 + 2x + 2 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned}\text{where } D &= b^2 - 4ac \\ &= 2^2 - 4.1.2 \\ &= 4 - 8 \\ &= -4\end{aligned}$$

from (A)

$$\begin{aligned}x &= \frac{-2 \pm \sqrt{-4}}{2} \\ &= \frac{-2 \pm 2i}{2} \\ &= -1 \pm i\end{aligned}$$

$$\therefore x = -1 + i, \quad -1 - i$$

Quadratic Equations Ex 14.1 Q9

$$5x^2 - 6x + 2 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned}\text{where } D &= b^2 - 4ac \\ &= (-6)^2 - 4.5.2 \\ &= 36 - 40 \\ &= -4\end{aligned}$$

from (A)

$$\begin{aligned}x &= \frac{-(-6) \pm \sqrt{-4}}{2.5} \\ &= \frac{6 \pm 2i}{10} \\ &= \frac{3 \pm i}{5}\end{aligned}$$

$$\therefore x = \frac{3}{5} + \frac{i}{5}, \quad \frac{3}{5} - \frac{i}{5}$$

Quadratic Equations Ex 14.1 Q10

$$21x^2 + 9x + 1 = 0$$

Comparing the given equation with the general form

$$ax^2 + bx + c = 0, \text{ we get } a = 21, b = 9, c = 1$$

Substituting a and b in,

$$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \text{ and } \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$\alpha = \frac{-9 + \sqrt{81 - 84}}{42} \text{ and } \beta = \frac{-9 - \sqrt{81 - 84}}{42}$$

$$\Rightarrow \alpha = \frac{-9 + \sqrt{-3}}{42} \text{ and } \beta = \frac{-9 - \sqrt{-3}}{42}$$

$$\Rightarrow \alpha = \frac{-9 + i\sqrt{3}}{42} \text{ and } \beta = \frac{-9 - i\sqrt{3}}{42}$$

$$\text{The roots are } x = \frac{-9}{42} \pm \frac{i\sqrt{3}}{42}$$

Quadratic Equations Ex 14.1 Q11

$$x^2 - x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\text{where } D = b^2 - 4ac$$

$$= (-1)^2 - 4.1.1$$

$$= 1 - 4$$

$$= -3$$

from (A)

$$\therefore x = \frac{-(-1) \pm \sqrt{-3}}{2}$$

$$= \frac{1 \pm \sqrt{3}i}{2}$$

$$\therefore x = \frac{1}{2} + \frac{\sqrt{3}}{2}i, \quad \frac{1}{2} - \frac{\sqrt{3}}{2}i$$

Quadratic Equations Ex 14.1 Q12

$$x^2 + x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned} \text{where } D &= b^2 - 4ac \\ &= 1^2 - 4.1.1 \\ &= 1 - 4 \\ &= -3 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-1 \pm \sqrt{-3}}{2} \\ &= \frac{-1 \pm \sqrt{3}i}{2} \end{aligned}$$

$$\therefore x = \frac{-1}{2} + \frac{\sqrt{3}}{2}i, \quad \frac{-1}{2} - \frac{\sqrt{3}}{2}i$$

Quadratic Equations Ex 14.1 Q13

$$17x^2 - 8x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned} \text{where } D &= b^2 - 4ac \\ &= (-8)^2 - 4.17.1 \\ &= 64 - 68 \\ &= -4 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-(-8) \pm \sqrt{-4}}{2.17} \\ &= \frac{8 \pm 2i}{34} \\ &= \frac{4 \pm i}{17} \end{aligned}$$

$$\therefore x = \frac{4}{17} + \frac{i}{17}, \quad \frac{4}{17} - \frac{i}{17}$$

Quadratic Equations Ex 14.1 Q14

$$27x^2 - 10x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= (-10)^2 - 4.27.1 \\ &= 100 - 108 \\ &= -8 \end{aligned}$$

from (A)

$$x = \frac{-(-10) \pm \sqrt{-8}}{54}$$

$$= \frac{10 \pm 2\sqrt{2}i}{54}$$

$$= \frac{5 \pm \sqrt{2}i}{27}$$

$$\therefore x = \frac{5}{27} + \frac{\sqrt{2}i}{27}, \quad \frac{5}{27} - \frac{\sqrt{2}i}{27}$$

Quadratic Equations Ex 14.1 Q15

$$17x^2 + 28x + 12 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= (28)^2 - 4.17.12 \\ &= 784 - 816 \\ &= -32 \end{aligned}$$

from (A)

$$x = \frac{-28 \pm \sqrt{-32}}{2.17}$$

$$= \frac{-28 \pm 4\sqrt{2}i}{34}$$

$$\therefore x = \frac{-14 \pm 2\sqrt{2}i}{17}$$

Quadratic Equations Ex 14.1 Q16

$$21x^2 - 28x + 10 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= (-28)^2 - 4.21.10 \\ &= 784 - 840 \\ &= -56 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-(-28) \pm \sqrt{-56}}{2.21} \\ &= \frac{28 \pm 2\sqrt{14}i}{42} \end{aligned}$$

$$\therefore x = \frac{2}{3} \pm \frac{\sqrt{14}}{21}i$$

Quadratic Equations Ex 14.1 Q17

$$8x^2 - 9x + 3 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= (-9)^2 - 4.8.3 \\ &= 81 - 96 \\ &= -15 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-b \pm \sqrt{D}}{2a} \\ &= \frac{-(-9) \pm \sqrt{-15}}{2.8} \\ &= \frac{9 \pm \sqrt{15}i}{16} \end{aligned}$$

Thus

$$\therefore x = \frac{9 \pm \sqrt{15}i}{16}$$

Quadratic Equations Ex 14.1 Q18

$$13x^2 + 7x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned} \text{where } D &= b^2 - 4ac \\ &= 7^2 - 4.13.1 \\ &= 49 - 52 \\ &= -3 \end{aligned}$$

Thus, from (A)

$$\begin{aligned} x &= \frac{-7 \pm \sqrt{-3}}{2.13} \\ &= \frac{-7 \pm \sqrt{3}i}{26} \end{aligned}$$

Thus

$$\therefore x = \frac{-7}{26} \pm \frac{\sqrt{3}}{26}i$$

Quadratic Equations Ex 14.1 Q19

$$2x^2 + x + 1 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\begin{aligned} \text{where } D &= b^2 - 4ac \\ &= 1^2 - 4.2.1 \\ &= 1 - 8 \\ &= -7 \end{aligned}$$

Thus, from (A)

$$\begin{aligned} x &= \frac{-1 \pm \sqrt{-7}}{2.2} \\ &= \frac{-1 \pm \sqrt{7}i}{4} \end{aligned}$$

Thus

$$\therefore x = \frac{-1}{4} \pm \frac{\sqrt{7}}{4}i$$

Quadratic Equations Ex 14.1 Q20

$$\sqrt{3}x^2 - \sqrt{2}x + 3\sqrt{3} = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= (-\sqrt{2})^2 - 4.\sqrt{3}.3\sqrt{3} \\ &= 2 - 36 \\ &= -34 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-(-\sqrt{2}) \pm \sqrt{-34}}{2.\sqrt{3}} \\ &= \frac{\sqrt{2} \pm \sqrt{34}i}{2\sqrt{3}} \end{aligned}$$

Thus

$$\therefore x = \frac{\sqrt{2} \pm \sqrt{34}i}{2\sqrt{3}}$$

Quadratic Equations Ex 14.1 Q21

$$\sqrt{2}x^2 + x + \sqrt{2} = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

where $D = b^2 - 4ac$

$$\begin{aligned} &= 1^2 - 4.\sqrt{2}.\sqrt{2} \\ &= 1 - 8 \\ &= -7 \end{aligned}$$

from (A)

$$\begin{aligned} x &= \frac{-1 \pm \sqrt{-7}}{2.\sqrt{2}} \\ &= \frac{-1 \pm \sqrt{7}i}{2\sqrt{2}} \end{aligned}$$

Thus

$$\therefore x = \frac{-1 \pm \sqrt{7}i}{2\sqrt{2}}$$

Quadratic Equations Ex 14.1 Q22

$$x^2 + x + \frac{1}{\sqrt{2}} = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\text{where } D = b^2 - 4ac$$

$$= 1^2 - 4 \cdot 1 \cdot \frac{1}{\sqrt{2}}$$

$$= 1 - 2\sqrt{2}$$

from (A)

$$x = \frac{-1 \pm \sqrt{-(2\sqrt{2} - 1)}}{2}$$

$$= \frac{-1 \pm \sqrt{2\sqrt{2} - 1}i}{2}$$

Thus,

$$\therefore x = \frac{-1 \pm \sqrt{2\sqrt{2} - 1}i}{2}$$

Quadratic Equations Ex 14.1 Q23

$$x^2 + \frac{x}{\sqrt{2}} + 1 = 0 \quad \Rightarrow \quad \sqrt{2}x^2 + x + \sqrt{2} = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$D = b^2 - 4ac$$

$$= 1^2 - 4 \cdot \sqrt{2} \cdot \sqrt{2}$$

$$= 1 - 8$$

$$= -7$$

from (A)

$$x = \frac{-1 \pm \sqrt{-7}}{2\sqrt{2}}$$

$$= \frac{-1 \pm \sqrt{7}i}{2\sqrt{2}}$$

Thus,

$$\therefore x = \frac{-1 \pm \sqrt{7}i}{2\sqrt{2}}$$

Quadratic Equations Ex 14.1 Q24

$$\sqrt{5}x^2 + x + \sqrt{5} = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\text{where } D = b^2 - 4ac$$

$$= 1^2 - 4 \cdot \sqrt{5} \cdot \sqrt{5}$$

$$= 1 - 20$$

$$= -19$$

from (A)

$$x = \frac{-1 \pm \sqrt{-19}}{2 \cdot \sqrt{5}}$$

$$= \frac{-1 \pm \sqrt{19}i}{2\sqrt{5}}$$

Thus,

$$\therefore x = \frac{-1 \pm \sqrt{19}i}{2\sqrt{5}}$$

Quadratic Equations Ex 14.1 Q25

$$-x^2 + x - 2 = 0$$

We will apply discriminant rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \dots\dots\dots (A)$$

$$\text{where } D = b^2 - 4ac$$

$$= 1^2 - 4 \cdot (-1) \cdot (-2)$$

$$= 1 - 8$$

$$= -7$$

from (A)

$$x = \frac{-1 \pm \sqrt{-7}}{2 \cdot (-1)}$$

$$= \frac{-1 \pm \sqrt{7}i}{-2}$$

Thus,

$$\therefore x = \frac{-1 \pm \sqrt{7}i}{-2}$$

Quadratic Equations Ex 14.1 Q26

We will apply discriminate rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \quad \dots\dots(A)$$

Where $D = b^2 - 4ac$

$$= (-2)^2 - 4(1)\left(\frac{3}{2}\right)$$

$$= 4 - 6$$

$$= -2$$

From (A)

$$x = \frac{-(-2) \pm \sqrt{-2}}{2(1)}$$

$$= \frac{2 \pm i\sqrt{2}}{2}$$

$$= 1 \pm \frac{i}{\sqrt{2}}$$

Thus,

$$\therefore x = 1 \pm \frac{i}{\sqrt{2}}$$

Quadratic Equations Ex 14.1 Q27

We will apply discriminate rule,

$$x = \frac{-b \pm \sqrt{D}}{2a} \quad \dots\dots(A)$$

Where $D = b^2 - 4ac$

$$= (-4)^2 - 4(3)\left(\frac{20}{3}\right)$$

$$= 16 - 80$$

$$= -64$$

From (A)

$$x = \frac{-(-4) \pm \sqrt{-64}}{2(3)}$$


$$= \frac{4 \pm i8}{6}$$

$$= \frac{2}{3} \pm \frac{4i}{3}$$

Thus,

$$\therefore x = \frac{2}{3} \pm \frac{4i}{3}$$

RD Sharma
Solutions
Class 11 Maths
Chapter 14
Ex 14.2



Quadratic Equations Ex 14.2 Q1(i)

$$x^2 + 10ix - 21 = 0$$

$$\Rightarrow x^2 + 10ix + 21i^2 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow x^2 + 7ix + 3ix + 21i^2 = 0$$

$$\Rightarrow x(x + 7i) + 3i(x + 7i) = 0$$

$$\Rightarrow (x + 3i)(x + 7i) = 0$$

$$\therefore x = -3i, -7i$$

Quadratic Equations Ex 14.2 Q1(ii)

$$x^2 + (1 - 2i)x - 2i = 0$$

$$\Rightarrow x^2 + x - 2i - 2i = 0$$

$$\Rightarrow x(x + 1) - 2i(x + 1) = 0$$

$$\Rightarrow (x - 2i)(x + 1) = 0$$

$$\Rightarrow x = 2i, -1$$

Quadratic Equations Ex 14.2 Q1(iii)

$$x^2 - (2\sqrt{3} + 3i)x + 6\sqrt{3}i = 0$$

$$\Rightarrow x^2 - 2\sqrt{3}x - 3ix + 6\sqrt{3}i = 0$$

$$\Rightarrow x(x - 2\sqrt{3}) - 3i(x - 2\sqrt{3}) = 0$$

$$\Rightarrow (x - 3i)(x - 2\sqrt{3}) = 0$$

$$\Rightarrow x = 3i, 2\sqrt{3}$$

Quadratic Equations Ex 14.2 Q1(iv)

$$6x^2 - 17ix - 12 = 0$$

$$\Rightarrow 6x^2 - 17ix + 12i^2 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow 6x^2 - 9ix - 8ix + 12i^2 = 0$$

$$\Rightarrow 3x(2x - 3i) - 4i(2x - 3i) = 0$$

$$\Rightarrow (3x - 4i)(2x - 3i) = 0$$

$$\Rightarrow x = \frac{4}{3}i \quad \text{or} \quad \frac{3}{2}i$$

Quadratic Equations Ex 14.2 Q2(i)

$$x^2 - (3\sqrt{2} + 2i)x + 6\sqrt{2}i = 0$$

$$\Rightarrow x^2 - 3\sqrt{2}x - 2ix + \sqrt{2}i = 0$$

$$\Rightarrow x(x - 3\sqrt{2}) - 2i(x - 3\sqrt{2}) = 0$$

$$\Rightarrow (x - 2i)(x - 3\sqrt{2}) = 0$$

$$\Rightarrow x = 2i \quad \text{or} \quad 3\sqrt{2}$$

Quadratic Equations Ex 14.2 Q2(ii)

$$x^2 - (5 - i)x + (18 + i) = 0$$

$$\Rightarrow x^2 - 5x - ix + 18 + i = 0$$

$$\Rightarrow x^2 - (3 - 4i)x - (2 + 3i)x + (18 + i) = 0$$

$$\Rightarrow x(x - (3 - 4i)) - (2 + 3i)(x - (3 - 4i)) = 0$$

$$\Rightarrow (x - (2 + 3i))(x - (3 - 4i)) = 0$$

$$\Rightarrow x = 2 + 3i \quad \text{or} \quad 3 - 4i$$

Quadratic Equations Ex 14.2 Q2(iii)

$$(2 + i)x^2 - (5 - i)x + 2(1 - i) = 0$$

$$\Rightarrow (2 + i)x^2 - 2x - (3 - i)x + 2(1 - i) = 0$$

$$\Rightarrow x[2 + i)x - 2] - (1 - i)[(2 + i)x - 2] = 0$$

$$\Rightarrow [x - (1 - i)][(2 + i)x - 2] = 0$$

$$\text{either } [x - (1 - i)] = 0 \quad \text{or} \quad [(2 + i)x - 2] = 0$$

$$\Rightarrow x = 1 - i \quad \text{or} \quad x = \frac{2}{2 + i}$$

$$\Rightarrow x = 1 - i \quad \text{or} \quad x = \frac{2 \times 2 - i}{(2 + i)(2 - i)}$$

$$\text{or } x = \frac{4 - 2i}{4 + 1} = \frac{4}{5} - \frac{2}{5}i$$

Thus,

$$x = 1 - i, \quad \frac{4}{5} - \frac{2}{5}i$$

Quadratic Equations Ex 14.2 Q2(iv)

$$x^2 - (2+i)x - (1-7i) = 0$$

$$\Rightarrow x^2 - (2+i)x - (1-7i) = 0$$

$$\Rightarrow x^2 - (3-i)x + (1-2i)x - (1-7i) = 0$$

$$\Rightarrow x(x - (3-i)) + (1-2i)(x - (3-i)) = 0$$

$$\Rightarrow [x + (1-2i)][x - (3-i)] = 0$$

$$\Rightarrow x = -1+2i, \quad 3-i$$

Quadratic Equations Ex 14.2 Q2(v)

$$ix^2 - 4x - 4i = 0$$

$$\Rightarrow ix^2 + 4i^2x + 4i^3 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow x^2 + 4ix + 4i^2 = 0$$

$$\Rightarrow x^2 + 2ix + 2ix + 4i^2 = 0$$

$$\Rightarrow x(x + 2i) + 2i(x + 2i) = 0$$

$$\Rightarrow (x + 2i)(x + 2i)$$

$$\therefore x = -2i, \quad -2i$$

Quadratic Equations Ex 14.2 Q2(vi)

$$x^2 + 4ix - 4 = 0$$

$$\Rightarrow x^2 + 4ix + 4i^2 = 0 \quad [\because i^2 = -1]$$

$$\Rightarrow x^2 + 2ix + 2ix + 4i^2 = 0$$

$$\Rightarrow x(x + 2i) + 2i(x + 2i) = 0$$

$$\Rightarrow (x + 2i)(x + 2i) = 0$$

$$\Rightarrow x = -2i, \quad -2i$$

Quadratic Equations Ex 14.2 Q2(vii)

$$2x^2 + \sqrt{15}ix - i = 0$$

Comparing the given equation with the general form

$$ax^2 + bx + c = 0, \text{ we get } a = 2, b = \sqrt{15}i, c = -i$$

Substituting a and b in,

$$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$\alpha = \frac{-\sqrt{15}i + \sqrt{-15 + 8i}}{4} \quad \text{and} \quad \beta = \frac{-\sqrt{15}i - \sqrt{-15 + 8i}}{4}$$

$$\text{Let } \sqrt{-15 + 8i} = a + bi$$

$$\Rightarrow -15 + 8i = (a + bi)^2$$

$$\Rightarrow -15 + 8i = a^2 - b^2 + 2abi$$

$$\Rightarrow a^2 - b^2 = -15 \quad \text{and} \quad 2abi = 8i$$

$$\text{Now } (a^2 + b^2)^2 = (a^2 - b^2)^2 + 4a^2b^2$$

$$\Rightarrow (a^2 + b^2)^2 = (-15)^2 + 64 = 289$$

$$\Rightarrow a^2 + b^2 = 17$$

Solving $a^2 - b^2 = -15$ and $a^2 + b^2 = 17$, we get

$$a^2 = 1 \quad \text{and} \quad b^2 = 16$$

$$\Rightarrow a = \pm 1 \quad \text{and} \quad b = \pm 4$$

$$\Rightarrow a = \pm 1 \text{ and } b = \pm 4$$

$$\Rightarrow a = 1, b = 4 \text{ or } a = -1, b = -4$$

$$\therefore \sqrt{-15+8i} = 1+4i, -1-4i$$

$$\text{When } \sqrt{-15+8i} = 1+4i$$

$$\alpha = \frac{-\sqrt{15}i + 1 + 4i}{4} = \frac{1 + (4 - \sqrt{15})i}{4}$$

$$\text{and } \beta = \frac{-\sqrt{15}i - (1 + 4i)}{4} = \frac{-1 - (4 + \sqrt{15})i}{4}$$

$$\text{When } \sqrt{-15+8i} = -1-4i$$

$$\alpha = \frac{-\sqrt{15}i - 1 - 4i}{4} = \frac{-1 - (4 + \sqrt{15})i}{4}$$

$$\text{and } \beta = \frac{-\sqrt{15}i - (-1 - 4i)}{4} = \frac{1 + (4 - \sqrt{15})i}{4}$$

Quadratic Equations Ex 14.2 Q2(viii)

$$x^2 - x + (1+i) = 0$$

$$x^2 - x + (1+i) = 0$$

$$x^2 - ix - (1-i)x + i(1-i) = 0$$

$$(x-i)(x-(1-i)) = 0$$

$$x = i, 1-i$$

Quadratic Equations Ex 14.2 Q2(ix)

We will apply discriminate rule on $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Now,

$$ix^2 - x + 12i = 0$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(i)(12i)}}{2i}$$

$$= \frac{1 \pm \sqrt{1 + 48}}{2i}$$

$$= \frac{1 \pm \sqrt{49}}{2i}$$

$$= \frac{1 \pm 7}{2i}$$

$$= \frac{8}{2i}, \frac{-6}{2i}$$

$$= \frac{4}{i}, -\frac{3}{i}$$

$$= -4i, 3i$$

Quadratic Equations Ex 14.2 Q2(x)

We will apply discriminate rule on $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Now,

$$x^2 - (3\sqrt{2} - 2i)x - \sqrt{2}i = 0$$

$$x = \frac{(3\sqrt{2} - 2i) \pm \sqrt{[-(3\sqrt{2} - 2i)]^2 - 4(1)(-\sqrt{2}i)}}{2(1)}$$

$$= \frac{(3\sqrt{2} - 2i) \pm \sqrt{(3\sqrt{2} - 2i)^2 + 4\sqrt{2}i}}{2}$$

$$= \frac{3\sqrt{2} - 2i}{2} \pm \frac{4 - \sqrt{2}i}{2}$$

Quadratic Equations Ex 14.2 Q2(xi)

$$x^2 - (\sqrt{2} + i)x + \sqrt{2}i = 0$$

$$x^2 - \sqrt{2}x - ix + \sqrt{2}i = 0$$

$$x(x - \sqrt{2}) - i(x - \sqrt{2}) = 0$$

$$(x - i)(x - \sqrt{2}) = 0$$

$$x = i, \sqrt{2}$$

Quadratic Equations Ex 14.2 Q2(xii)

$$2x^2 - (3 + 7i)x + (9i - 3) = 0$$

$$2x^2 - 3x - 7ix + (9i - 3) = 0$$

$$(2x - 3 - i)(x - 3i) = 0$$

$$\left(x - \frac{3+i}{2}\right)(x - 3i) = 0$$

$$x = \frac{3+i}{2}, 3i$$

Kopykitab
Same textbooks, click away