
RD Sharma Solutions apter 6

Ex 6.1 Andrew and a september 19 and a sep Class 11 Maths

To obtain the graph of  $y = 3 \sin x$  we first draw the graph of  $y = \sin x$  in the interval  $\lceil 0,2\pi \rceil$ . The maximum and minimum values are 3 and - 3 respectively.



We have,

$$y = 2\sin\left(x - \frac{\pi}{4}\right)$$

$$\Rightarrow \qquad (y-0) = 2\sin\left(x - \frac{\pi}{4}\right)$$


Shifting the origin at  $\left(\frac{\pi}{4},0\right)$ , we have

$$X = X + \frac{\pi}{4} \text{ and } y = Y + 0$$

Substituting these values in (i), we get

$$Y = 2 \sin X$$

Thus we draw the graph of  $Y = 2\sin X$  and shift it by  $\frac{\pi}{4}$  to the right to get the required graph.



We have,

$$y = 2\sin(2x - 1)$$

$$\Rightarrow (y-0) = 2 \sin 2 \left(x - \frac{1}{2}\right)$$


Shifting the origin at  $\left(\frac{1}{2},0\right)$ , we have

$$X = X + \frac{1}{2}$$
 and  $y = Y + 0$ 

Substituting these values in (i), we get

$$Y = 2 \sin 2X$$

Thus we draw the graph of  $Y = 2 \sin 2X$  and shift it by 1/2 to the right to get the required graph.



We have,

$$y=3\sin\left(3x+1\right)$$

$$\Rightarrow \qquad (y-0) = 3\sin 3\left(x + \frac{1}{3}\right)$$


Shifting the origin at  $\left(-\frac{1}{3},0\right)$ , we have

$$X = X - \frac{1}{3}$$
 and  $y = Y + 0$ 

Substituting these values in (i), we get

$$Y = 3 \sin 3X$$

Thus we draw the graph of  $Y = 3 \sin 3X$  and shift it by 1/3 to the left to get the required graph.



We have,

$$y = 3\sin\left(2x - \frac{\pi}{4}\right)$$


$$\Rightarrow \qquad (y-0) = 3\sin 2\left(x - \frac{\pi}{8}\right)$$


Shifting the origin at  $\left(\frac{\pi}{8}, 0\right)$ , we have

$$X = X + \frac{\pi}{8} \text{ and } y = Y + 0$$

Substituting these values in (i), we get

$$Y = 3 \sin 2X$$





We have,

$$y = \sin\left(x + \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \sin\left(x + \frac{\pi}{4}\right)$$


Shifting the origin at  $\left(-\frac{\pi}{4},0\right)$ , we obtain

$$x=X-\frac{\pi}{4},\ y=Y+0$$

Substituting these values in (i), we get

Thus we draw the graph of  $Y = \sin X$  and shift it by  $\frac{\pi}{4}$  to the left to get the required graph

To obtain the graph of  $y = \sin 3x$  we first draw the graph of  $y = \sin x$  in the interval  $[0,2\pi]$  and then divide the x-coordinates of the points where it crosses x-axis by 3.





RD Sharma
Solutions
Class 11 Maths
Chapter 6
Ex 6.2

We have,

$$y = \cos\left(x + \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \cos\left(x + \frac{\pi}{4}\right)$$
 Shifting the origin at  $\left(-\frac{\pi}{4}, 0\right)$ , we obtain


$$X = X - \frac{\pi}{4}, \ y = Y + 0$$

Substituting these values in (i), we get

$$Y = \cos X$$
.

Thus we draw the graph of  $Y = \cos X$  and shift it by  $\frac{\pi}{4}$  to the left to get the required graph.

---(i)



We have,

$$y = \cos\left(x - \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \cos\left(x - \frac{\pi}{4}\right)$$

Shifting the origin at 
$$(\frac{\pi}{-}, 0)$$
, we obtain


Shifting the origin at  $\left(\frac{\pi}{4},0\right)$ , we obtain

$$x = X + \frac{\pi}{4}, y = Y + 0$$

Substituting these values in (i), we get

$$Y = \cos X$$
.

Thus we draw the graph of  $Y = \cos X$  and shift it by  $\frac{\pi}{4}$  to the right to get the required graph.



We have,

$$y = 3\cos(2x - 1)$$

$$\Rightarrow \qquad (y-0) = 3\cos 2\left(x-\frac{1}{2}\right)$$


Shifting the origin at  $\left(\frac{1}{2},0\right)$ , we have

$$x = X + \frac{1}{3} \text{ and } y = Y + 0$$

Substituting these values in (i), we get

$$Y = 3\cos 2X$$

Thus we draw the graph of  $Y = 3\cos 2X$  and shift it by 1/2 to the right to get the required graph.



We have,

$$y = 2\cos\left(x - \frac{\pi}{2}\right)$$


$$\Rightarrow y - 0 = 2\cos\left(x - \frac{\pi}{2}\right)$$

Shifting the origin at  $\left(\frac{\pi}{2},0\right)$ , we obtain


$$x=X+\frac{\pi}{2}\,,\ y=Y+0$$

Substituting these values in (i), we get

$$Y = 2\cos X$$



## Chapter 6 Graphs of Trigonometric Functions Ex 6.2 Q2



We have,

1

$$y = \cos 2\left(x - \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \cos 2\left(x - \frac{\pi}{4}\right)$$

---(i)


Shifting the origin at  $\left(\frac{\pi}{4},0\right)$ , we obtain

$$x=X+\frac{\pi}{4},\ y=Y+0$$

Substituting these values in (i), we get


$$Y = \cos 2X$$
.

Thus we draw the graph of  $Y = \cos 2X$  and shift it by  $\frac{\pi}{4}$  to the right to get the required graph.



To obtain the graph of  $y = \cos \frac{x}{2}$  we first draw the graph of  $y = \cos x$  in the interval

 $\left[0,2\pi\right]$  and then divide the x-coordinates of the points where it crosses x-axis by 1/2.



RD Sharma Solutions apter 6

Ex. 6.3

Little levilled less little levilled le Class 11 Maths

We know that

$$y = sin^2 x = \frac{1 - \cos 2x}{2} = \frac{1}{2} - \frac{1}{2} \cos 2x$$

We have,

$$y = \frac{1}{2} - \frac{1}{2}\cos 2x$$

$$\Rightarrow \qquad y - \frac{1}{2} = -\frac{1}{2} \cos 2x$$

Shifting the origin at  $\left(0, -\frac{1}{2}\right)$ , we obtain

$$x = X, y = Y + \frac{1}{2}$$

Substituting these values in (i), we get

 $Y = -\frac{1}{2}\cos 2X.$ 

Thus we draw the graph of  $Y = \cos 2X$ , adjust the maximum and minimum values to 1/2 and -1/2

and shift it by  $\frac{1}{2}$  up to get the required graph.

We know that

$$y = \cos^2 x = \frac{1 + \cos 2x}{2} = \frac{1}{2} + \frac{1}{2}\cos 2x$$

We have,

$$y = \frac{1}{2} + \frac{1}{2}\cos 2x$$

$$y - \frac{1}{2} = \frac{1}{2}\cos 2x \qquad ---(i)$$


Shifting the origin at  $\left(0, -\frac{1}{2}\right)$ , we obtain

$$x=X\,,\ y=Y+\frac{1}{2}$$

Substituting these values in (i), we get

$$Y = -\frac{1}{2}\cos 2X.$$

Thus we draw the graph of  $Y = \cos 2X$ , adjust the maximum and minimum values to 1/2 and -1/2 and shift it by  $\frac{1}{2}$  down to get the required graph.



## Chapter 6 Graphs of Trigonometric Functions Ex 6.3 Q3

We have,

$$y = \sin^2\left(x - \frac{\pi}{4}\right)$$

$$\Rightarrow y - 0 = \sin^2\left(x - \frac{\pi}{4}\right)$$


Shifting the origin at  $\left(\frac{\pi}{4},0\right)$ , we obtain

$$X = X + \frac{\pi}{4}, \quad Y = Y + 0$$

Substituting these values in (i), we get


$$Y = \sin^2 X$$
.

Thus we draw the graph of  $Y = \sin^2 X$  and shift it by  $\frac{\pi}{4}$  to the right to get the required graph.




## Chapter 6 Graphs of Trigonometric Functions Ex 6.3 Q4


To obtain the graph of y = tan 2x we first draw the graph of y = tan x in the interval  $\left(\frac{\pi}{2}\right)$  and then divide the x-coordinates of the points where it crosses x-axis by 2.



To obtain the graph of  $y = 2 \tan 3x$  we first draw the graph of  $y = \tan x$  in the interval  $\left(-\frac{\pi}{2}, -\frac{\pi}{2}\right)$  and then divide the x-coordinates of the points where it crosses x-axis by 3. We then stretch the graph vertically by a factor of 2.



To obtain the graph of  $y=2\cot 2x$  we first draw the graph of  $y=\cot x$  in the interval  $(0,\pi)$  and then divide the x-coordinates of the points where it crosses x-axis by 2. We then stretch the graph vertically by a factor of 2. d We



## Chapter 6 Graphs of Trigonometric Functions Ex 6.3 Q7

$$y = \cos 2\left(x - \frac{\pi}{6}\right)$$

$$\Rightarrow \qquad y - 0 = \cos 2\left(x - \frac{\pi}{6}\right) \qquad ---(i)$$


Shifting the origin at  $\left(\frac{\pi}{6},0\right)$ , we obtain

$$X = X + \frac{\pi}{6}, \quad y = Y + 0$$

Substituting these values in (i), we get

$$Y = \cos 2X$$
.

Thus we draw the graph of  $Y = \cos 2X$  and shift it by  $\frac{\pi}{6}$  to the right to get the required graph.



We know that

$$y = \sin^2 x = \frac{1 - \cos 2x}{2} = \frac{1}{2} - \frac{1}{2} \cos 2x$$

$$y = \frac{1}{2} - \frac{1}{2}\cos 2x$$

$$\Rightarrow \qquad y - \frac{1}{2} = -\frac{1}{2}\cos 2x$$

---(i)


Shifting the origin at  $\left(0, -\frac{1}{2}\right)$ , we obtain

$$X = X, y = Y + \frac{1}{2}$$


$$Y = -\frac{1}{2}\cos 2X.$$

 $x=X,\ y=Y+\frac{1}{2}$  Substituting these values in (i), we get  $Y=-\frac{1}{2}\cos 2X.$  Thus we draw the graph of  $Y=\cos 2X$ , adjust the maximum and minimum values to 1/2 and -1/2 

and shift it by  $\frac{1}{2}$  up to get the required graph.



# Chapter 6 Graphs of Trigonometric Functions Ex 6.3 Q9



## Chapter 6 Graphs of Trigonometric Functions Ex 6.3 Q10

